

# First results of asymmetry with QADC signals of single $\text{PbWO}_4$ crystal

---

Cosenza INFN Group

( L. La Rotonda, E. Meoni, A. Policicchio, G.  
Susinno,  
T. Venturelli)

# Data set

We have started to study QADC data taken by single  $\text{PbWO}_4$  crystal at different angles with electrons of 50 GeV both **without Pb** in front to the crystal and **with different mm of Pb** in front to the crystal.

The goal is computed: asymmetry versus angle for different shower depths and the fraction of Cerenkov component (as previous year)

This study is related to second paper : "Separation of crystal signals into Cerenkov and scintillation components".

## 1. Stand-alone measurements of single $\text{PbWO}_4$ crystals (room temperature)

(Data relevant for paper #2)

Runs 112 - 135, logbook page 16

Beam: 50 GeV  $e^-$ ,  $\theta = 0^\circ$

### • Angular scan of the crystal response (electrons)

Runs 402 - 433, logbook pages 33-34

Beam: 50 GeV  $e^-$ ,  $\theta = -60^\circ$  to  $+60^\circ$

Analysed

### • Angular scan of the crystal response (mips)

Runs 497 - 532, logbook pages 38-39

Beam: 70 GeV  $\pi^-$ ,  $\theta = -75^\circ$  to  $+75^\circ$

To be analysed

### • Angular scans of the response to developing showers (50 GeV $e^- + \text{Pb}$ )

10 mm Pb upstream, Runs 256 - 278,  $\theta = -35^\circ$  to  $+35^\circ$ , logbook page 25

20 mm Pb upstream, Runs 284 - 309,  $\theta = -35^\circ$  to  $+35^\circ$ , logbook page 26

35 mm Pb upstream, Runs 310 - 336,  $\theta = -45^\circ$  to  $+45^\circ$ , logbook page 28

4 mm Pb upstream, Runs 434 - 457,  $\theta = -35^\circ$  to  $+35^\circ$ , logbook page 35

8 mm Pb upstream, Runs 458 - 460,  $\theta = -30^\circ$  to  $+30^\circ$ , logbook page 35


14 mm Pb upstream, Runs 461 - 463,  $\theta = -30^\circ$  to  $+30^\circ$ , logbook page 35

25 mm Pb upstream, Runs 466 - 487,  $\theta = -35^\circ$  to  $+35^\circ$ , logbook p. 35,37


Analysed

To be analysed


# 30 GeV electron showers in the early stage





ADC spectra after pedestal subtraction



Selected only events with ADC counts larger than 3 pedestal RMS




We have equalized ADC signals of the two channels at 0° degrees



The ADC signal increases with the angle, because increases the track length.


For positive angles the CH25 signal is larger than CH24 one because CH25 collects a larger Cerenkov component (vice versa for negative ones), except for positive angles 55° and 60°.

But at fixed positive and negative corresponding angles, the difference between the signal of the 2 CHs is not equal, this is evident also in asymmetry slope.



**Asymmetry**  $\frac{\text{ADC}_{\text{mean}}(\text{CH25}) - \text{ADC}_{\text{mean}}(\text{CH24})}{\text{ADC}_{\text{mean}}(\text{CH24}) + \text{ADC}_{\text{mean}}(\text{CH25})}$

$\text{ADC}_{\text{mean}}$  is the mean of the ADC spectra shown before




At positive angles the slope is similar to the one observed last year

At negative angles slope quite different to be investigated!


We have computed asymmetry also event by event at a fixed angle

$$\text{Asymmetry} = \frac{\text{ADC}_{\text{coun}}(\text{CH25}) - \text{ADC}_{\text{coun}}(\text{CH24})}{\text{ADC}_{\text{coun}}(\text{CH24}) + \text{ADC}_{\text{coun}}(\text{CH25})}$$




Taken mean value of the asymmetry distribution for each angle

Errors of the plot are RMS of asymmetry distributions






# The two methods give the same results



## Upstream


ADC spectra after pedestal subtraction



This cut seems not optimal.  
(see Tommaso's presentation about these tails)  
Selected events with  $ADC\_counts > 100$



Asymmetry computed as in the previous case both from ADC spectra distributions and event by events



**Asy comp. evt by evt**  
**Asy comp. From ADC spectra**



# 50 GeV electrons shower with 10mm, 20mm, 35mm of Pb upstream

In these runs observed a problem: pedestal values of physics runs are different from pedestal values of pedestal runs.




The differences are for all runs at level of :

24.5-25.5 ADC counts for ch24 and of 5.5 ADC counts for ch25.


First part of ADC spectrum after pedestal subtraction




# ADC spectra after correct pedestal subtraction





## Asymmetry computed from ADC spectra



## Asymmetry computed event by event





$$\frac{\text{Cerenkov}}{\text{Signal}}(30^\circ) = \frac{2\text{Asym}}{(1 + \text{Asym})}$$

$\frac{\text{Cerenkov}}{\text{Signal}}(30^\circ) = 10.5\%$  (without Pb) from positive angle

$\frac{\text{Cerenkov}}{\text{Signal}}(30^\circ) = 18\%$  (without Pb) from negative angles

$\frac{\text{Cerenkov}}{\text{Signal}}(30^\circ) = 10.5\%$  (without Pb) from previous testbeam with ele of 10 GeV

**The difference must be investigated!**



# To do

---

Understand the difference between negative and positive angles

Look at the data with 8-14-25 mm of Pb in front to the crystal with electrons of 50 GeV

Look data with pions of 70 GeV (Davide Pinci has already analysed them looking at oscilloscope data)