

# Jet substructure: back to basics

Gavin Salam (CERN)


CMS Jet Substructure workshop  
CERN, 15 April 2013

As a field we've devised  $O(10-20)$  powerful methods to tag jet substructure.

Many of the methods have been tried out in searches and work; these kinds of methods will be crucial for searches in the years to come.

# Very active research field



Some of the tools developed for boosted W/Z/H/top reconstruction





apologies for omitted taggers, arguable links, etc.

# Seeing W's and tops in a single jet

## W's in a single jet



## tops in a single jet



As a field we've devised  $O(10-20)$  powerful methods to tag jet substructure.

Many of the methods have been tried out in searches and work; these kinds of methods will be crucial for searches in the years to come.

But from outside, the many methods make the field look pretty confusing.

And from inside, I get the impression we don't always know *why* or *how* the methods work – which is bad if we're looking for robustness.

**Is it time to get back to basics?**

# What was the original motivation?

Normal  $R=0.4/0.5$  jet finding fails to find one jet per prong of a boosted  $[W/Z/H/top/NP]$  hadronic decay.

We need to make sure that this doesn't prevent us from using EW-scale particles in TeV scale searches.

## **Question #1:**

To what extent are the things we do with “normal” jets (and leptons) mirrored in the things we're doing with “fat” jets?

# What have we found out in the meantime?

There's a huge number of things you can do with jet substructure.

Many of the things appear to improve mass resolution, background rejection, etc. [at least in MC simulation]

## **Question #2:**

How should we balance improvements v. “complexity” of method?

# What are we comfortable with?

## Resolved Analysis

Find one jet/prong

Cut on jet  $p_t$ ,  $\Delta y$ , ...



# What are we comfortable with?

## Resolved Analysis

Find one jet/prong

Cut on jet  $p_t$ ,  $\Delta y$ , ...



## Fat-jet Analysis

Find subjets

Cut on subjet  $z$ ,  $\Delta R$ , ...

[MDT/Prune/Trim/Filt/XYZTopTagger/Template ...]

# What are we comfortable with?

## Resolved Analysis

Find one jet/prong

Cut on jet  $p_t$ ,  $\Delta y$ , ...

Isolation cut for  
colourless leptons,  $\gamma$

Cut on radiation in jet  
for q/g discrimination



## Fat-jet Analysis

Find subjets

Cut on subjet  $z$ ,  $\Delta R$ , ...

[MDT/Prune/Trim/Filt/XYZTopTagger/Template ...]

# What are we comfortable with?

## Resolved Analysis

Find one jet/prong

Cut on jet  $p_t$ ,  $\Delta y$ , ...

Isolation cut for  
colourless leptons,  $\gamma$

Cut on radiation in jet  
for  $q/g$  discrimination



## Fat-jet Analysis

Find subjets

Cut on subjet  $z$ ,  $\Delta R$ , ...


[MDT/Prune/Trim/Filt/XYZTopTagger/Template ...]

Cut on radiation for  
colourless  $W, H, \dots$

Cut on radiation in  
subjets

[ $\tau_{mn}$ ,  $Q_{jets}$ , deconstruction...]


# What are we comfortable with?



# [Analytic] understanding

Understanding your taggers means you know what tools you can safely use with them

For robustness, you can then choose taggers whose distributions can be predicted in many ways



Just like  
MET( $Z \rightarrow \nu\nu$ ) in  
multijets is reliably  
estimated from  
 $\gamma$ +jets because  
multiple types of  
calculations of the  
ratio agree

# What do we know currently?

## Boost 2010 proceedings:

The [Monte Carlo] findings discussed above indicate that while [pruning, trimming and filtering] have qualitatively similar effects, there are important differences. For our choice of parameters, pruning acts most aggressively on the signal and background followed by trimming and filtering.


## At the time:

- No clear picture of why the taggers might be similar or different
- No clear picture of how the parameter choices affect the taggers

## Today:


- I'll show a selection of **preliminary** lessons from studies for background jets in progress with Dasgupta, Fregoso and Marzani

# The “right” MC study can already be instructive (testing on background [quark] jets)






# The “right” MC study can already be instructive (testing on background [quark] jets)



Different taggers  
are apparently  
quite similar

# The “right” MC study can already be instructive (testing on background [quark] jets)




But only for a  
limited range  
of masses


# Do we care about such differences?

- Think data-driven backgrounds: kinks can seriously mess with you [especially if you got used to their being absent, e.g. from moderate  $p_t$  tests]
- How do these structures depend on the  $z_{\text{cut}}$ ,  $y_{\text{cut}}$ ,  $R_{\text{trim}}$ , etc. parameters?
- Are these structures telling us something we might want to know about the taggers? E.g. how to improve them?

**This calls for analysis and calculation**



Pruning [7, 8] takes an initial jet, and from its mass deduces a pruning radius  $R_{\text{prune}} = R_{\text{fact}} \cdot \frac{2m}{p_t}$ , where  $R_{\text{fact}}$  is a parameter of the tagger. It then reclusters the jet and for every clustering step, involving objects  $a$  and  $b$ , it checks whether  $\Delta_{ab} > R_{\text{prune}}$  and  $\min(p_{ta}, p_{tb}) < z_{\text{cut}} p_{t,(a+b)}$ , where  $z_{\text{cut}}$  is a second parameter of the tagger. If so, then the softer of the  $a$  and  $b$  is discarded. Otherwise  $a$  and  $b$  are recombined as usual. Clustering then proceeds with the remaining objects, applying the pruning check at each stage.




## What pruning is meant to do:

Choose an  $R_{\text{prune}}$  such that different hard prongs ( $p_1, p_2$ ) end up in different hard subjects.

Discard any softer radiation.

**Sets pruning radius, & hard enough to end up as subject**

Pruning [7, 8] takes an initial jet, and from its mass deduces a pruning radius  $R_{\text{prune}} = R_{\text{fact}} \cdot \frac{2m}{p_t}$ , where  $R_{\text{fact}}$  is a parameter of the tagger. It then reclusters the jet and for every clustering step, involving objects  $a$  and  $b$ , it checks whether  $\Delta_{ab} > R_{\text{prune}}$  and  $\min(p_{ta}, p_{tb}) < z_{\text{cut}} p_{t,(a+b)}$ , where  $z_{\text{cut}}$  is a second parameter of the tagger. If so, then the softer of the  $a$  and  $b$  is discarded. Otherwise  $a$  and  $b$  are recombined as usual. Clustering then proceeds with the remaining objects, applying the pruning check at each stage.




## What pruning sometimes does

Chooses  $R_{\text{prune}}$  based on a soft  $p_3$  (dominates total jet mass), and leads to a single narrow subjet whose mass is also dominated by a soft emission ( $p_2$ , within  $R_{\text{prune}}$  of  $p_1$ , so not pruned away).

**Sets pruning radius,  
but gets pruned away**

# A simple fix: “sane” pruning


Require at least one successful merging with  $\Delta R > R_{\text{prune}}$  and  $z > z_{\text{cut}}$



“sane” pruning is effectively placing an isolation cut on radiation around the tagged (colour-neutral) object

# Mass drop tagger

1. Break the jet  $j$  into two subjects by undoing its last stage of clustering. Label the two subjects  $j_1, j_2$  such that  $m_{j_1} > m_{j_2}$ .
2. If there was a significant mass drop,  $m_{j_1} < \mu m_j$ , and the splitting is not too asymmetric,  $y = \min(p_{tj_1}^2, p_{tj_2}^2) \Delta R_{j_1 j_2}^2 / m_j^2 > y_{\text{cut}}$ , then deem  $j$  to be the tagged jet.
3. Otherwise redefine  $j$  to be equal to  $j_1$  and go back to step 1 (unless  $j$  consists of just a single particle, in which case the original jet is deemed untagged).




## What MDT does wrong:

Can follow a soft branch ( $p_2 + p_3 < y_{\text{cut}} p_{\text{jet}}$ ) with “accidental” small mass, when the “right” answer was that the (massless) hard branch had no substructure

**Subjet is soft, but has more substructure than hard subjet**

# A simple fix for “modified” Mass Drop Tagger:

When recursing, follow branch with larger  $(m^2+p_t^2)$   
(rather than the one with larger  $m$ )






# What about analytic calculations of the taggers?

Simpler


More complex




Dasgupta, Fregoso, Marzani & GPS, forthcoming:  
LL in all cases, plus some subleading logs  
[NB: LL doesn't mean the same thing in all cases!]

# Modified Mass Drop Tagger

Monte Carlo




Analytic




[mMDT is closest we have to a scale-invariant tagger,  
though exact behaviour depends on q/g fractions]


## Monte Carlo




## Analytic



## Monte Carlo




## Analytic



# Why is this useful?


If MC & analytics always agree, why bother with calculations?



Preliminary

# Why is this useful?

Different MC showers don't always agree.  
That turns into a systematic uncertainty.




Preliminary

# Why is this useful?

Different MC showers don't always agree.  
That turns into a systematic uncertainty.

Analytics can tell you which is "right".



Preliminary

# Bottom line on “understanding”

- Taggers may be quite simple to write, but potentially involved to understand – tiny details can lead even courageous theorists to tears.
- Contrast this with  $p_t$  cuts for standard jet analyses – (mostly) simple
- Still, many taggers/groomers are within calculational reach.
- New “modified” Mass Drop Tagger is especially simple; New “sane” pruning is also interesting
- The better you understand a tagger, the better you can detect signals



# Infrared safety

## Infrared safety:

When the addition of one soft particle with momentum  $\epsilon$  changes the outcome of tagging by an amount  $O(1)$ .

It means that perturbative calculations give  $\infty$

It means that the physics of hard objects may be irremediably contaminated by non-perturbative physics – not good for robustness or calculability!

Was long an issue in hadron-collider jet-finding.  
Let's make sure it doesn't come back to haunt us!

# Pruning followed by a mass-drop cut:



- Cut on the mass drop,  $\mu = m_1/m$ 
  - $m_1$  is mass of highest pT subjet
  - subjets defined by un-clustering last step

This is  
infrared  
unsafe

# Pruning followed by a mass-drop cut:

- Cut on the mass drop,  $\mu = m_1/m$ 
  - $m_1$  is mass of highest pT subjet
  - subjets defined by **un-clustering last step**



This is infrared unsafe



# Pruning followed by a mass-drop cut:

- Cut on the mass drop,  $\mu = m_1/m$ 
  - $m_1$  is mass of highest pT subjet
  - subjets defined by **un-clustering last step**



This is infrared unsafe



# Pruning followed by a mass-drop cut:

- Cut on the mass drop,  $\mu = m_1/m$ 
  - $m_1$  is mass of highest pT subjet
  - subjets defined by **un-clustering last step**



This is infrared unsafe



# Pruning followed by a mass-drop cut:

- Cut on the mass drop,  $\mu = m_1/m$ 
  - $m_1$  is mass of highest pT subjet
  - subjets defined by **un-clustering last step**


This is infrared unsafe



If you use “sane” pruning, the IR safety problem gets cured

## N-subjettiness $\tau_3 / \tau_2$ :


$\tau_2$  measures departure from 2-parton energy flow  
 $\tau_3$  measures departure from 3-parton energy flow





## N-subjettiness $\tau_3 / \tau_2$ :

$\tau_2$  measures departure from 2-parton energy flow  
 $\tau_3$  measures departure from 3-parton energy flow



Easily cured with a cut on  $\tau_2 / \tau_1$ , which forces 3<sup>rd</sup> prong not to be soft.

Extra cut has almost no impact on performance

Cacciari et al '12

# Scale invariant searches




arXiv:1303.6636 Gouzevitch,  
Oliveira, Rojo, Rosenfeld & Sanz

Experiments often have two distinct searches:


Resolved (small-R multi-jet)

Boosted (large-R fat-jet)

Can resolved and boosted analyses be consistently performed together?





# Look just at $Y \rightarrow 2Z$ decay



## Key [simple] idea:

Cuts on resolved jets should mirror those on subjects inside fat jets






$$\left| \frac{2(m_{Y1} - m_{Y2})}{m_{Y1} + m_{Y2}} \right| \leq f_m \quad M_Y (1 - f_m) \leq m_{Y1}, m_{Y2} \leq M_Y (1 + f_m)$$

$$\Delta y \equiv |y_{Y1} - y_{Y2}| \leq \Delta y_{\max} \quad \Delta y \equiv |y_{Yi,1} - y_{Yi,2}| \leq \Delta y_{\max}^{\text{res}}$$

$$\Delta p_T \equiv p_T^{(1)} - p_T^{(2)} \geq (1 - y_{\text{cut}}) p_T^{(1)} \quad \max(m_{Y,1}, m_{Y,2}) \leq \mu \cdot m_Y$$

Traditional cuts  
on resolved jets




Boosted cuts  
on resolved jets

Jet Reconstruction					
$R$	$R_{\text{sj}}$	$R_f$	$n_{\text{filt}}$	$\mu$	$y_{\text{cut}}$
0.5	1.3	0.3	3	0.67	0.09

Basic cuts		
$p_T^{\min}$	$ y_{\max} $	$H_T^{\min}$
25 GeV	5.0	100 GeV

Quality requirements			
$M_Y$	$\Delta y_{\max}$	$\Delta y_{\max}^{\text{res}}$	$f_m$
125 GeV	1.3	1.5	0.15


$X > 2Y > 4Z$ , Toy MC, Parton Level



Efficiency  
roughly  
stable as  
 $X \rightarrow 2Y \rightarrow 4Z$   
goes from  
 $4 \rightarrow 3 \rightarrow 2$  jets

Cuts are close to those that  
optimise  $S/\sqrt{B}$  in all regions


$X > 2Y > 4Z$ , Toy MC, Hadron Level, LHC 8 TeV



Efficiency roughly stable as  $X \rightarrow 2Y \rightarrow 4Z$  goes from 4  $\rightarrow$  3  $\rightarrow$  2 jets

Cuts are close to those that optimise  $S/\sqrt{B}$  in all regions

$X > 2Y > 4Z$ , Toy MC, Hadron Level, LHC 8 TeV



Efficiency roughly independent of R used in clustering

**Bottom line:**  
traditional and substructure techniques can be used together  
[an analogous method still needs to be worked out for top]



# Pileup in the boosted regime

## Pronged taggers

Some have pileup-**reduction** built in (MassDrop+Filtering, Pruning, HEPTopTagger, Template), essentially by using small ( $R \sim 0.2-0.3$ ) sub-cones, sometimes dynamically adjusted to the jet  $p_t$

For heavy pileup you will need to supplement them with full pileup **subtraction** (e.g. area-based).

[Technically trivial, but so far studied only for filtering & trimming]

## Shape-based taggers

Until recently, no clear way of subtracting pileup.

[Grooming reduces PU, but also discards info]

# Pileup subtraction for shapes

Cacciari, Dutta, JH Kim, GPS & Soyez '12

$n^{\text{th}}$  **derivative of shape** wrt ghost momenta

Shape as a function of particle momenta in jet

$$V_{\text{jet}}^{[n]} \equiv A_g^n \frac{d^n}{dr_{t,g}^n} V(\{p_i\}_{\text{jet}})$$

Ghost area

# Pileup subtraction for shapes

Cacciari, Dutta, JH Kim, GPS & Soyez '12

$n^{\text{th}}$  **derivative of shape** wrt ghost momenta

Shape as a function of particle momenta in jet

$$V_{\text{jet}}^{[n]} \equiv A_g^n \frac{d^n}{dr_{t,g}^n} V(\{p_i\}_{\text{jet}})$$

Ghost area


$$V_{\text{jet,sub}} = V_{\text{jet}} - \rho V_{\text{jet}}^{[1]} + \frac{1}{2} \rho^2 V_{\text{jet}}^{[2]} + \dots$$

Subtracted shape

pileup density

# Practical test: $\tau_{32}$ and top tagging

## Correcting the $\tau_{32}$ distribution




**Green: no PU**

**Red: with PU**


**Blue/Black: subtracted**

# Practical test: $\tau_{32}$ and top tagging

## Correcting the $\tau_{32}$ distribution



## Tagging efficiency



**Green: no PU**

**Red: with PU**

**Blue/Black: subtracted**

# Closing

# Other things I would have liked to talk about

## **Jet deconstruction**

Matrix-element method extended to all orders.

Soper & Spannowsky, [arXiv:1102.3480](#), [arXiv:1211.3140](#)

## **Q-jets**

Does clustering have to give a unique answer? What if you probe multiple possible clustering histories?

Ellis et al. [arXiv:1201.1914](#), [arXiv:1304.2394](#)

## **Jet substructure by accident**

Rather than looking for 16 jets (e.g. in BSM  $\rightarrow$  4 low- $p_t$  tops), look for  $O(4)$  fat jets, each with substructure. May be easier to reliably predict backgrounds

Cohen, Izaguirre, Lisanti & Lou [arXiv:1212.1456](#)

## **Quark–Gluon discrimination**

What information are we exploiting? Can we exploit it better?

Gallicchio & Schwartz '12  
Larkoski, GPS & Thaler, in preparation



# Other things I would have liked to talk about

## FastJet Contrib

A space for people to contribute their own jet-tool libraries, to provide users with uniform, regularly updated and reliable access to a broad range of jet tools.



The screenshot shows a web browser window with the address bar containing `fastjet.hepforge.org/contrib/`. The page title is "FastJet Contrib". The main text explains that the space is for 3rd party extensions of FastJet. It provides instructions to download the current version, `fjcontrib-1.002`, released on 12 April 2013, and to run `./configure`, `make`, `make check`, and `make install`. It also notes that contributions are accessed by including `fastjet/contrib/SomeContrib.hh` in C++ files.

**FastJet Contrib**

The fastjet-contrib space is intended to provide a common location for access to 3rd party extensions of **FastJet**.

**Download** the current version: `fjcontrib-1.002` (released 12 April 2013), which contains **these contributions**. Changes relative to earlier versions are briefly described in the **NEWS** file.

After downloading and unpacking, enter the `fjcontrib-1.002/` directory and then run

```
./configure [--fastjet-config=FILE] [--prefix=...] [...]  
make  
make check          # optional  
make install
```

By default the package installs to the same directories as the FastJet installation.

A contribution named "SomeContrib" is usually accessed by including "fastjet/contrib/SomeContrib.hh" in your C++ file, and linking with `-ISomeContrib`.

# Summary [and points for discussion]

It's time to make the transition to a deep understanding of our tools, the only way of guaranteeing robustness

- ▶ Analytical control of “pronged” taggers now seems to be possible [though still early days]
- ▶ Taggers can have surprises in store for us – especially when we explore full LHC14  $p_t$  range.  
[They can also be “fixed up”, e.g. sane pruning, modified MDT]
- ▶ When do we want want to use “radiation-based” taggers?  
[Should resolved analyses always exploit q/g discrimination?]
- ▶ Do we need/want continuous resolved–fat-jet analyses?
- ▶ Pileup: it's time to start dealing with it systematically in our taggers [beyond just grooming, even as part of grooming]

EXTRAS

# Different fat-jet tagger types

## Prong based

(e.g. HEPTopTagger,  
Template Tagger)


- Identifies prongs
- Requires prongs be consistent with kinematics of  $t \rightarrow Wb \rightarrow 3$  quarks

## Radiation based

(e.g. N-subjettiness =  $\tau_3/\tau_2$   
+ mass cut)

- Requires top-mass consistency (maybe with some grooming)
- Exploits weaker radiation from top (3 quarks) than background (1q+2g or 3g)

# Top quarks, Tops from quark-jets & Top fat-jets [and their radiation pattern]




# Are top pairs in high- $p_t$ events always back-to-back?


A reminder that top-quarks at LHC are almost “light”

An 8 TeV study with POWHEG, top-pair production, no decay and no parton showering (to keep things simple)

## Flavour Creation



## Flavour Excitation – tops inside your PDFs





## Gluon Splitting

