

Flavour Physics & CP Violation

Lecture 4 of 4

Tim Gershon
University of Warwick

CERN Summer Student Lecture Programme
5th August 2013


Contents

- Part 1
 - What is flavour physics & why is it interesting?
- Part 2
 - What do we know from previous experiments?
- **Part 3**
 - **What do we hope to learn from current experiments?**
- **Part 4**
 - **The future of flavour physics**

Rare Decays

$b \rightarrow s\gamma$ rate and photon energy spectrum

Archetypal FCNC probe for new physics



Belle PRL 103 (2009) 241801


$$B(B \rightarrow X_s \gamma)_{E_\gamma > 1.7 \text{ GeV}} = (3.45 \pm 0.15 \pm 0.40) \times 10^{-4}$$

consistent with the SM prediction

... but SM also predicts that the photon is polarised


$b \rightarrow s\gamma$ photon polarisation measurement

- Search for time-dependent asymmetry
 - Observable effect requires NP: left-handed current & new CP phase



Excellent prospects for LHCb with $B_s \rightarrow \phi\gamma$

NPB 867 (2013) 1



First measurement of photon polarisation in $b \rightarrow s\gamma$ transitions





- New LHCb measurement

LHCb-CONF-2013-009

- uses $B^+ \rightarrow K^+\pi\pi\gamma$ decays
- compare γ direction relative to $K^+\pi\pi$ plane


$$A_{ud} = -0.085 \pm 0.019 \text{ (stat)} \pm 0.003 \text{ (syst)}$$

A_{ud} proportional to polarisation
(unfortunately, constant of proportionality has large uncertainty)



$$B \rightarrow K^* \mu^+ \mu^-$$

- $b \rightarrow s l^+ l^-$ processes also governed by FCNCs
 - rates and asymmetries of many exclusive processes sensitive to NP
- Queen among them is $B_d \rightarrow K^{*0} \mu^+ \mu^-$
 - superb laboratory for NP tests
 - **experimentally clean signature**
 - many kinematic variables ...
 - ... with clean theoretical predictions (at least at low q^2)



Operator Product Expansion

Build an effective theory for b physics

- take the weak part of the SM
- integrate out the heavy fields (W,Z,t)
- (like a modern version of Fermi theory for weak interactions)

$$\mathcal{L}_{(\text{full EW} \times \text{QCD})} \longrightarrow \mathcal{L}_{\text{eff}} = \mathcal{L}_{\text{QED} \times \text{QCD}} \left(\begin{array}{l} \text{quarks } \neq t \\ \& \text{ leptons} \end{array} \right) + \sum_n C_n(\mu) Q_n$$

Q_n - local interaction terms (operators), C_n - coupling constants (Wilson coefficients)

Wilson coefficients

- encode information on the weak scale
- are calculable and known in the SM (at least to leading order)
- are affected by new physics

For $K^* \mu \mu$ we care about C_7 (also affects $b \rightarrow s \gamma$), C_9 and C_{10}

Effective operators

$$\mathcal{H}_W^{\Delta B=1, \Delta C=0, \Delta S=-1} = 4 \frac{G_F}{\sqrt{2}} \left(\lambda_c^s (C_1(\mu) Q_1^c(\mu) + C_2(\mu) Q_2^c(\mu)) \right. \\ \left. + \lambda_u^s (C_1(\mu) Q_1^u(\mu) + C_2(\mu) Q_2^u(\mu)) - \lambda_t^s \sum_{i=3}^{10} C_i(\mu) Q_i(\mu) \right)$$

where the $\lambda_q^s = V_{qb}^* V_{qs}$ and the operator basis is given by

$$\begin{aligned} Q_1^q &= \bar{b}_L^\alpha \gamma^\mu q_L^\alpha \bar{q}_L^\beta \gamma_\mu s_L^\beta & Q_2^q &= \bar{b}_L^\alpha \gamma^\mu q_L^\beta \bar{q}_L^\beta \gamma_\mu s_L^\alpha \\ Q_3 &= \bar{b}_L^\alpha \gamma^\mu s_L^\alpha \sum_q \bar{q}_L^\beta \gamma_\mu q_L^\beta & Q_4 &= \bar{b}_L^\alpha \gamma^\mu s_L^\beta \sum_q \bar{q}_L^\beta \gamma_\mu q_L^\alpha \\ Q_5 &= \bar{b}_L^\alpha \gamma^\mu s_L^\alpha \sum_q \bar{q}_R^\beta \gamma_\mu q_R^\beta & Q_6 &= \bar{b}_L^\alpha \gamma^\mu s_L^\beta \sum_q \bar{q}_R^\beta \gamma_\mu q_R^\alpha \\ Q_7 &= \frac{3}{2} \bar{b}_L^\alpha \gamma^\mu s_L^\alpha \sum_q e_q \bar{q}_R^\beta \gamma_\mu q_R^\beta & Q_8 &= \frac{3}{2} \bar{b}_L^\alpha \gamma^\mu s_L^\beta \sum_q e_q \bar{q}_R^\beta \gamma_\mu q_R^\alpha \\ Q_9 &= \frac{3}{2} \bar{b}_L^\alpha \gamma^\mu s_L^\alpha \sum_q e_q \bar{q}_L^\beta \gamma_\mu q_L^\beta & Q_{10} &= \frac{3}{2} \bar{b}_L^\alpha \gamma^\mu s_L^\beta \sum_q e_q \bar{q}_L^\beta \gamma_\mu q_L^\alpha \end{aligned}$$

Four-fermion operators (except $Q_{7\gamma}$ & Q_{8g}) – dimension 6

$$Q_{7\gamma} = \frac{e}{16\pi^2} m_b \bar{b}_L^\alpha \sigma^{\mu\nu} F_{\mu\nu} s_L^\alpha$$

$$Q_{8g} = \frac{g_s}{16\pi^2} m_b \bar{b}_L^\alpha \sigma^{\mu\nu} G_{\mu\nu}^A T^A s_L^\alpha$$


$$Q_{9V} = \frac{1}{2} \bar{b}_L^\alpha \gamma^\mu s_L^\alpha \bar{l} \gamma_\mu l$$

$$Q_{10A} = \frac{1}{2} \bar{b}_L^\alpha \gamma^\mu s_L^\alpha \bar{l} \gamma_\mu \gamma_5 l$$

Theory of $B \rightarrow K^* \mu^+ \mu^-$

- Given for inclusive $b \rightarrow s \mu^+ \mu^-$ for simplicity
 - physics of exclusive modes \approx same but equations are more complicated (involving form factors, etc.)
- Differential decay distribution

$$\frac{d^2\Gamma}{dq^2 d \cos \theta_l} = \frac{3}{8} \left[(1 + \cos^2 \theta_l) H_T(q^2) + 2 \cos \theta_l H_A(q^2) + 2 (1 - \cos^2 \theta_l) H_L(q^2) \right]$$




$$H_T(q^2) \propto 2q^2 \left[\left(C_9 + 2C_7 \frac{m_b^2}{q^2} \right)^2 + C_{10}^2 \right],$$

$$H_A(q^2) \propto -4q^2 C_{10} \left(C_9 + 2C_7 \frac{m_b^2}{q^2} \right),$$


$$H_L(q^2) \propto \left[(C_9 + 2C_7)^2 + C_{10}^2 \right].$$

This term gives a forward-backward asymmetry

Angular analysis of $B^0 \rightarrow K^{*0} \mu^+ \mu^-$





LHCb arXiv:1304.6325
 See also CDF PRL 108 (2012) 081807
 BaBar PRD 86 (2012) 032012
 ATLAS-CONF-2013-038 & CMS BPH-11-009



Analysis performed in bins of dimuon invariant mass squared (q^2)

Angular analysis of $B^0 \rightarrow K^{*0} \mu^+ \mu^-$



LHCb arXiv:1304.6325
 See also CDF PRL 108 (2012) 081807
 BaBar PRD 86 (2012) 032012
 ATLAS-CONF-2013-038 & CMS BPH-11-009



First measurement of zero-crossing point of A_{FB}


$$q_0^2 = (4.9 \pm 0.9) \text{ GeV}^2/c^4$$

Consistent with SM expectation



New observables in $B^0 \rightarrow K^{*0} \mu^+ \mu^-$


LHCb-PAPER-2013-037




Interesting tension with the SM prediction

Isospin asymmetry in $B \rightarrow K^{(*)} \mu \mu$

JHEP 07 (2012) 133



Deviation from zero integrated over $q^2 \sim 4.4\sigma$
 Consistent with previous measurements
 (BaBar, Belle, CDF)




Consistent with zero & with SM prediction
 Consistent with previous measurements
 (BaBar, Belle, CDF)

$$B_{(s)}^0 \rightarrow \mu^+ \mu^-$$

$$B_{(s)}^0 \rightarrow \mu^+ \mu^-$$


Killer app. for new physics discovery

- Very small in the SM
- Huge NP enhancement
($\tan \beta = \text{ratio of Higgs vevs}$)
- Clean experimental signature



$$BR(B_s \rightarrow \mu^+ \mu^-)^{SM} = (3.3 \pm 0.3) \times 10^{-8} \quad BR(B_s \rightarrow \mu^+ \mu^-)^{MSSM} \propto \tan^6 \beta / M_{A^0}^4$$


$$B_{(s)}^0 \rightarrow \mu^+ \mu^-$$



Searches over 30 years

$B_{(s)}^0 \rightarrow \mu^+ \mu^-$ – analysis ingredients


- Produce a very large sample of B mesons
- Trigger efficiently on dimuon signatures
- Reject background
 - excellent vertex resolution (identify displaced vertex)
 - excellent mass resolution (identify B peak)
 - also essential to resolve B^0 from B_s^0 decays
 - powerful muon identification (reject background from B decays with misidentified pions)
 - typical to combine various discriminating variables into a multivariate classifier
 - e.g. Boosted Decision Trees algorithm



$$B_{(s)}^0 \rightarrow \mu^+ \mu^-$$


latest results from CMS & LHCb

CMS arXiv:1307.5025




Events weighted by S/(S+B)

LHCb arXiv:1307.5024



Only events with BDT > 0.7

$B_{(s)}^0 \rightarrow \mu^+ \mu^-$ – combined results




$$B(B_s^0 \rightarrow \mu^+ \mu^-) = (2.9 \pm 0.7) \times 10^{-9}$$

Future flavour physics projects


LHCb upgrade

- To fully exploit LHC potential for heavy flavour physics will require an upgrade to LHCb
 - full readout & trigger at 40 MHz to enable high L running
 - “high L” = $10^{33}/\text{cm}^2/\text{s}$ (so independent of machine upgrade)
 - planned for 2018 shutdown
- Physics case:
 - “exploration” of 1st phase will become “precision studies”
 - new opportunities for exploration open up (e.g. testing consistency of CP violation in tree vs. loop processes)

LHC upgrade and the all important trigger




higher luminosity
 → need to cut harder at L0 to keep rate at 1 MHz
 → lower efficiency



- readout detector at 40 MHz
- implement trigger fully in software → efficiency gains
- run at L_{inst} up to $2 \times 10^{33} / \text{cm}^2 / \text{s}$

LHCb detector upgrade



LHCb upgrade timeline

- 2011
 - Letter of Intent: [CERN-LHCC-2011-001](#)
- 2012
 - Framework TDR: [CERN-LHCC-2012-007](#)
 - **Endorsed by LHCC and approved by CERN Research Board** ([minutes](#))
 - LHCb upgrade features prominently in draft European Strategy for Particle Physics
 - See also [arXiv:1208.3355](#) for physics discussion
- **2013**
 - Sub-detector TDRs ← **preparation of TDRs already started**
- 2014-17
 - Final R&D, production and construction
- 2018 (LS2)
 - Installation of upgraded LHCb detector (requires 18 months)

Upgrade – expected sensitivities

Type	Observable	Current precision	LHCb 2018	Upgrade (50 fb ⁻¹)	Theory uncertainty
B_s^0 mixing	$2\beta_s (B_s^0 \rightarrow J/\psi \phi)$	0.10 [9]	0.025	0.008	~ 0.003
	$2\beta_s (B_s^0 \rightarrow J/\psi f_0(980))$	0.17 [10]	0.045	0.014	~ 0.01
	$A_{fs}(B_s^0)$	6.4×10^{-3} [18]	0.6×10^{-3}	0.2×10^{-3}	0.03×10^{-3}
Gluonic penguin	$2\beta_s^{\text{eff}}(B_s^0 \rightarrow \phi\phi)$	–	0.17	0.03	0.02
	$2\beta_s^{\text{eff}}(B_s^0 \rightarrow K^{*0}\bar{K}^{*0})$	–	0.13	0.02	< 0.02
	$2\beta_s^{\text{eff}}(B^0 \rightarrow \phi K_S^0)$	0.17 [18]	0.30	0.05	0.02
Right-handed currents	$2\beta_s^{\text{eff}}(B_s^0 \rightarrow \phi\gamma)$	–	0.09	0.02	< 0.01
	$\tau^{\text{eff}}(B_s^0 \rightarrow \phi\gamma)/\tau_{B_s^0}$	–	5 %	1 %	0.2 %
Electroweak penguin	$S_3(B^0 \rightarrow K^{*0}\mu^+\mu^-; 1 < q^2 < 6 \text{ GeV}^2/c^4)$	0.08 [14]	0.025	0.008	0.02
	$s_0 A_{\text{FB}}(B^0 \rightarrow K^{*0}\mu^+\mu^-)$	25 % [14]	6 %	2 %	7 %
	$A_1(K\mu^+\mu^-; 1 < q^2 < 6 \text{ GeV}^2/c^4)$	0.25 [15]	0.08	0.025	~ 0.02
	$\mathcal{B}(B^+ \rightarrow \pi^+\mu^+\mu^-)/\mathcal{B}(B^+ \rightarrow K^+\mu^+\mu^-)$	25 % [16]	8 %	2.5 %	$\sim 10\%$
Higgs penguin	$\mathcal{B}(B_s^0 \rightarrow \mu^+\mu^-)$	1.5×10^{-9} [2]	0.5×10^{-9}	0.15×10^{-9}	0.3×10^{-9}
	$\mathcal{B}(B^0 \rightarrow \mu^+\mu^-)/\mathcal{B}(B_s^0 \rightarrow \mu^+\mu^-)$	–	$\sim 100\%$	$\sim 35\%$	$\sim 5\%$
Unitarity triangle angles	$\gamma (B \rightarrow D^{(*)}K^{(*)})$	$\sim 10\text{--}12^\circ$ [19, 20]	4°	0.9°	negligible
	$\gamma (B_s^0 \rightarrow D_s K)$	–	11°	2.0°	negligible
	$\beta (B^0 \rightarrow J/\psi K_S^0)$	0.8° [18]	0.6°	0.2°	negligible
Charm	A_Γ	2.3×10^{-3} [18]	0.40×10^{-3}	0.07×10^{-3}	–
CP violation	ΔA_{CP}	2.1×10^{-3} [5]	0.65×10^{-3}	0.12×10^{-3}	–

Table 1: Statistical sensitivities of the LHCb upgrade to key observables. For each observable the current sensitivity is compared to that which will be achieved by LHCb before the upgrade, and that which will be achieved with 50 fb⁻¹ by the upgraded experiment. Systematic uncertainties are expected to be non-negligible for the most precisely measured quantities.

- sample sizes in most exclusive B and D final states far larger than those collected elsewhere
- no serious competition in study of B_s decays and CP violation

Other future flavour experiments



- SuperKEKB/Belle2 & SuperB
 - $B \rightarrow \tau \nu$, inclusive measurements, τ physics, ...
- Rare kaon decays
 - $K^+ \rightarrow \pi^+ \nu \nu$ (NA62, CERN); $K^0 \rightarrow \pi^0 \nu \nu$ (KOTO, J-PARC)
- Muon to electron conversion (charged lepton flavour violation)
 - COMET/PRIME (J-PARC); $\mu 2e$ (FNAL)

B → τν and charged Higgs limits

- Pure leptonic decays of charged B mesons very clean
 - clean SM prediction
 - clean effect of charged Higgs (2HDM or SUSY)

$$BR(B^+ \rightarrow l^+ \nu)^{SM} = \frac{G_F m_B}{8\pi} m_l^2 \left(1 - \frac{m_l^2}{m_B^2}\right)^2 f_B^2 |V_{ub}|^2 \tau_B \quad BR(B^+ \rightarrow l^+ \nu)^{NP} = BR(B^+ \rightarrow l^+ \nu)^{SM} \left(1 - \frac{m_B^2}{m_H^2} \tan^2 \beta\right)^2$$


Belle PRD 82 (2010) 071101




KEKB to SuperKEKB



Colliding bunches





New superconducting / permanent final focusing quads near the IP



New beam pipe & bellows

Replace short dipoles with longer ones (LER)




Redesign the lattices of HER & LER to squeeze the emittance

Add / modify RF systems for higher beam current

Low emittance positrons to inject

Damping ring



Positron source


New positron target / capture section

Low emittance gun

Low emittance electrons to inject




TiN-coated beam pipe with antechambers



To obtain x40 higher luminosity

Belle II Detector



EM Calorimeter:
CsI(Tl), waveform sampling (barrel)
Pure CsI + waveform sampling (end-caps)

KL and muon detector:
Resistive Plate Counter (barrel outer layers)
Scintillator + WLSF + MPPC (end-caps, inner 2 barrel layers)

electrons (7GeV)

Particle Identification
Time-of-Propagation counter (barrel)
Prox. focusing Aerogel RICH (fwd)


Beryllium beam pipe
2cm diameter

Vertex Detector
2 layers DEPFET + 4 layers DSSD

positrons (4GeV)

Central Drift Chamber
He(50%):C₂H₆(50%), small cells,
long lever arm, fast electronics

Schedule



The holy grail of kaon physics: $K \rightarrow \pi \nu \nu$

Highest CKM suppression
of the $s \rightarrow d$ coupling:

$$A \sim (m_t/m_W)^2 |V_{ts}^* V_{td}| \sim \lambda^5$$

SM branching ratios

(Brod, Gorbahn, Stamou; PRD83 (2011) 034030)


Mode	$BR_{SM} \times 10^{11}$
$K^+ \rightarrow \pi^+ \nu \bar{\nu} (\gamma)$	$7.81 \pm 0.75 \pm 0.29$
$K_L \rightarrow \pi^0 \nu \bar{\nu}$	$2.43 \pm 0.39 \pm 0.06$



CKM parametric
(mainly $|V_{ts}|$)

Intrinsic


SM: box and penguin diagrams



Next generation experiments should
measure these decays for the 1st time


- $K^+ \rightarrow \pi^+ \nu \nu$ (NA62, CERN)
- $K^0 \rightarrow \pi^0 \nu \nu$ (KOTO, J-PARC)
- Proposals also at FNAL

NA62 Technique: Decay in Flight




$$m_{miss}^2 = (\tilde{p}_K - \tilde{p}_\pi)^2$$

Kinematically Constraint Decays




Unconstraint Decays






NA62: $K^+ \rightarrow \pi^+ \nu \nu$ in-flight @ CERN-SPS




- SPS primary p: 400 GeV/c
Unseparated beam:
- 75 GeV/c
 - 750 MHz
 - $\pi/K/p$ (~6% K^+)



KOTO at JPARC



Future projects



The need for more precision

- “Imagine if Fitch and Cronin had stopped at the 1% level, how much physics would have been missed”
– A.Soni
- “A special search at Dubna was carried out by Okonov and his group. They did not find a single $K_L^0 \rightarrow \pi^+\pi^-$ event among **600 decays** into charged particles (Anikira et al., JETP 1962). At that stage the search was terminated by the administration of the lab. **The group was unlucky.**”
– L.Okun

(remember: $B(K_L^0 \rightarrow \pi^+\pi^-) \sim 2 \cdot 10^{-3}$)

Summary

- We still don't know:
 - why there are so many fermions in the SM
 - what causes the baryon asymmetry of the Universe
 - where exactly the new physics is ...
 - ... and what its flavour structure is
- Prospects are good for progress in the next few years
- We need a continuing programme of flavour physics into the 2020s
 - complementary to the high- p_T programme of the LHC

References and background reading

- Reviews by the Particle Data Group
 - <http://pdg.lbl.gov/>
- Heavy Flavour Averaging Group (HFAG)
 - <http://www.slac.stanford.edu/xorg/hfag/>
- CKMfitter & UTfit
 - <http://ckmfitter.in2p3.fr/> & <http://www.utfit.org/>
- Review journals (e.g. Ann. Rev. Nucl. Part. Phys.)
 - <http://nucl.annualreviews.org>
- Proceedings of CKM workshops
 - Phys.Rept. 494 (2010) 197, eConf C100906
- Books
 - CP violation, I.I.Bigi and A.I.Sanda (CUP)
 - CP violation, G.C.Branco, L.Lavoura & J.P.Silva (OUP)