

Valence quarks and k_T factorisation

Michal Deak, Hannes Jung, **Krzysztof Kutak**

DESY, Hamburg


Motivation

- HERA \rightarrow hints that at rapidities $y > 3$ there can be new kind of dynamics \rightarrow BFKL, CCFM, BK. However, at HERA we cannot go to the more forward region.
- LHC will allow for this study. The interesting region can be studied if we consider QCD Compton scattering:



Jets at LHC - collinear approach


Jets are initiated by hard subprocesses (a), (b). QCD Compton (a) is the relevant hard process if we want to study low x_g effects where x_g is a longitudinal momentum fraction carried by gluon.



To go to low x_g safely we need to consider off-shell gluon...

QCD Compton in k_T factorisation

Incoming gluon is not collinear to the proton it is off-shell. Valence quark is collinear to the proton. The upper quark line \rightarrow replaced by gluon distribution after matrix element is calculated



All five diagrams are required by gauge invariance.

Details of kinematics

The Sudakov decomposition is:

$$k = x_g p_A + z_g p_B + k_T$$

$$q = x_q$$

$$k' = x_{g'} p_A + z_{g'} p_B + k'_T$$

$$q' = z_{q'} p_A + x_{q'} p_B + q'_T$$

Mandelstam variables are:

$$s = (p_1 + p_2)^2$$


$$\hat{s} = (k + q)^2$$

$$\hat{u} = (k - q')^2$$

$$\hat{t} = (k - k')^2$$

We are interested in configuration where:

$$\hat{s}, \hat{t}, \hat{u} \ll s$$





Hard matrix element

After squaring amplitude we obtain:

$$|M|^2 = -(4\pi)^2 \alpha_s^2 \frac{x_g^2 s^2 (x_q^2 + x_{q'}^2)}{18 \hat{s} \hat{u} \hat{t}} \left(\frac{\hat{s}(8x_q + x_{q'}) - \hat{u}(8x_{q'} + x_q)}{x_q - x_{q'}} + k^2 \right)$$




- In collinear limit $k^2 \rightarrow 0$ one obtains QCD Compton
- Symmetry $\hat{s} \rightarrow -\hat{u}$, $x_{q'} \rightarrow x_q$

Some properties of the off-shell ME



- off-shell ME allows for larger q'_q

Some properties of the off-shell ME



- at 0 angle there is a singularity at $k'_q = k_g$

Cross-section for 2 jets

Cross-section for 2-jet production:



$$\sigma \sim f_g(x_g, k_T^2, \mu) \otimes |M|^2 \otimes f_q(x_q, k_{T2}^2, \mu)$$

- $f_g(x_g, k_T^2, \mu)$ unintegrated gluon density ← CCFM
- $f_q(x_q, k_T^2, \mu)$ unintegrated valence quark density (needed for technical reasons) ← CCFM-like. Initial valence quark distribution is provided by CTEQ 6.1
- $\alpha_s \rightarrow \alpha_s(k_T^2)$
- cut on momenta of outgoing jets $\rightarrow p_T > 2.5 \text{ GeV}$

The result for the total x-section is roughly: 10 mb



For comparison total pp x-section at LHC energies is roughly 80 mb

Results - x distribution before collision




- Incoming off-shell gluon carrying low momentum fraction
- Incoming on-shell quark carrying large momentum fraction



Results - pseudo-rapidity distribution of produced jets



- valence quark ($m_q = 0$) is slightly deflected
- produced jets are well separated in rapidity




Results - p_T spectra of produced jets



Conclusions and outlook

- We obtained matrix element in k_T factorisation which allow for studies of low x_g effects at the LHC
 - CCFM-like equation for unintegrated quark distribution has been incorporated in Monte Carlo framework → CASCADE
 - p_T and rapidity spectra of produced jets have been calculated
 - Consistency check with collinear approach has been done
-
- Step towards including multiple interactions for MC generator in k_T factorisation framework
 - Important for testing different models
 - Since gluon is probed at low momentum fraction we are going to include nonlinear evolution equation to parametrize unintegrated gluon distribution

Back up slides



- Rapidity difference between produced jets