

$B_q \rightarrow \mu^+ \mu^-$
The Standard Model Prediction

Emmanuel Stamou

emmanuel.stamou@weizmann.ac.il

Weizmann Institute of Science



CKM 2014

Vienna, Austria

September 10, 2014

In collaboration with:


C. Bobeth, M. Gorbahn

C. Bobeth, M. Gorbahn, T. Hermann, M. Misiak, M. Steinhauser

arXiv:1311.1348

arXiv:1311.0903

$B_s \rightarrow \mu^+ \mu^-$ within and beyond the SM




Within the SM

- FCNC → loop-induced
- helicity suppressed → $\text{BR} \propto \frac{m_\mu^2}{M_{B_s}^2}$
- CKM suppressed → $\text{BR} \propto |V_{tb}^* V_{td}|^2 \propto \sin^4 \theta_c \sim 0.0016$

rare decay $\text{BR} \sim 10^{-9}$

- single dim-6 vectorial operator $Q_{10} = [\bar{s}_L \gamma_\nu b_L][\bar{\mu} \gamma^\nu \gamma_5 \mu]$
(B_s pseudo-scalar, no γ -penguin, scalar operators suppressed)
- single hadronic quantity, decay constant f_{B_s} at the 2% level
theoretically clean decay [follow-up talk by M. Wingate]

$B_s \rightarrow \mu^+ \mu^-$ within and beyond the SM




Beyond the SM

[talk by W. Altmannshofer]


- **sensitive** to scalar and pseudo-scalar operators in models with **extended Higgs sector** (2HDM, MSSM)
i.e. in MSSM $\text{BR} \propto \tan^6 \beta$.
- constraints on effective Z couplings to quarks **comparable** with **electroweak precision** tests
i.e. MFV, models with partial compositeness

[Haisch, Weiler '07, Guadagnoli, Isidori '13]

$B_s \rightarrow \mu^+ \mu^-$ within and beyond the SM




- sensitive to anomalous ttZ couplings [talk by A. Greljo]
- first time possible @ LHC (NLO analysis [Röntsch, Schulze '14])



[Brod, Greljo, ES, Uttayarat '14]

$B_s \rightarrow \mu^+ \mu^-$ within and beyond the SM



Experimental status

[talk by F. Archilli]

- LHCb and CMS measure the time-integrated BR

$$\text{BR}(B_s \rightarrow \mu^+ \mu^-) = (2.8^{+0.7}_{-0.6}) \times 10^{-9}$$


$$\text{BR}(B_d \rightarrow \mu^+ \mu^-) = (3.9^{+1.6}_{-1.4}) \times 10^{-10}$$

[new average, see talk by F. Archilli]

- in SM related to instantaneous BR $^{[t=0]}$:

$$\text{BR} = \frac{1}{1 - \tau_{B_s} \Delta \Gamma_s / 2} \text{BR}^{[t=0]}$$


[De Bruyn, Fleischer, Knegjens, Koppenburg, Merk '12,
arXiv:1204.1737]



Outline


- Recent developments in the SM prediction of $B_s \rightarrow \mu^+ \mu^-$
 - NNLO QCD corrections (1.8% → 0.2%)
[Hermann, Misiak, Steinhauser '13, arXiv:1311.1347]
 - NLO electroweak corrections (8% → 0.6%)
[Bobeth, Gorbahn, ES '13, arXiv:1311.1348]
- New BR prediction in the SM
 - error budget
[Bobeth, Gorbahn, Hermann, Misiak, ES, Steinhauser '13, arXiv:1311.0903]

The multi-scales of B Decays



\mathcal{L}_{SM}

Matching



$$\mathcal{L}_{\text{eff}} = \mathcal{L}_{\text{QCD}} + \mathcal{L}_{\text{QED}} + \sum_i \mathbf{C}_i(\mu) \underbrace{Q_i}_{\text{non-renorm.}}$$

RGE

- simple for QCD, no mixing
(Q_{10} conserved current)
- operator mixing under QED
(mixing of Q_9 , Q_2 & QCD penguins)

$$\langle 0 | \bar{s} \gamma_\nu \gamma_5 b | B_s(p) \rangle = i f_{B_s} p_\nu \text{ computed on the lattice}$$



formally μ_0, μ_b independent
→ estimate of theo. uncertainty

$$\text{BR} \propto f_{B_s}^2 |\mathbf{C}_{10}(\mu_b)|^2$$

NNLO QCD

$B_s \rightarrow \mu^+ \mu^-$ at LO, NLO & NNLO QCD

$$C_{10} = G_F \frac{\alpha_{em}}{4\pi s_w^2} \left(c_{10}^{\text{LO}} + \dots \right)$$





[Inami, Lim '81]

- neglect EW scale/scheme dependence for now

- LO has large μ_0 dependence from $m_t(\mu_0)$
- $m_t(\mu_0)$ $\overline{\text{MS}}$ mass w.r.t. QCD and on-shell w.r.t. EW interactions
($m_t(50\text{GeV}) = 180.8 \text{ GeV}$, $m_t(163.5) = 163.5 \text{ GeV}$, $m_t(300\text{GeV}) = 156.2 \text{ GeV}$)

$B_s \rightarrow \mu^+ \mu^-$ at LO, NLO & NNLO QCD

$$C_{10} = G_F \frac{\alpha_{em}}{4\pi s_w^2} \left(c_{10}^{\text{LO}} + \frac{\alpha_s}{4\pi} c_{10}^{\text{NLO QCD}} + \dots \right)$$



[Buchalla, Buras '93; Misiak, Urban '99]



- neglect EW scale/scheme dependence for now

- @NLO strong reduction of scale uncertainty on BR (1.8%)
- @NLO good convergence for $\mu \simeq m_t$, $\sim +2.2\%$ on BR
- 1.8%NLO \rightarrow 0.2%NNLO scale uncertainty on BR**

[Hermann, Misiak, Steinhauser '13, arXiv:1311.1347]

$B_s \rightarrow \mu^+ \mu^-$ at LO, NLO & NNLO QCD

$$C_{10} = G_F \frac{\alpha_{em}}{4\pi s_w^2} \left(C_{10}^{\text{LO}} + \frac{\alpha_s}{4\pi} C_{10}^{\text{NLO QCD}} + \frac{\alpha_s^2}{(4\pi)^2} C_{10}^{\text{NNLO QCD}} + \dots \right)$$



[Hermann, Misiak, Steinhauser '13,
arXiv:1311.1347]

- neglect EW scale/scheme dependence for now

- @NLO strong reduction of scale uncertainty on BR (1.8%)
- @NLO good convergence for $\mu \simeq m_t$, $\sim +2.2\%$ on BR
- 1.8%NLO \rightarrow 0.2%NNLO scale uncertainty on BR**

[Hermann, Misiak, Steinhauser '13, arXiv:1311.1347]

NLO EW

Turning on Electroweak Interactions

- so far we treated EW parameters as numbers

Questions

- which numbers?
EW parameters G_F , M_Z^{pole} , M_W^{pole} , M_H^{pole} , M_t^{pole} , $\alpha_{em}^{\overline{\text{MS}}}$, $s_w^{\overline{\text{MS}}}$ are not all independent
- how does the BR depend on this choice (EW renormalisation scheme)?
i.e. $s_w^2 \overline{\text{MS}} = 0.231$ VS $s_w^2 \text{on-shell} = 0.223$
- what about scale dependence?

→ 8% uncertainty on the BR

- uncertainty was not included in SM prediction so far
- ambiguities removed at NLO in EW interactions

Electroweak Corrections I/II

Step 1: effective Lagrangian normalisation

$$\mathcal{L} = \frac{4G_F}{\sqrt{2}} \frac{\alpha_{em}}{4\pi s_w^2} \left(c_{10}^{\text{LO}} \left(\frac{m_t^2}{M_W^2} \right) + \frac{\alpha_{em}}{4\pi} c_{10}^{\text{EW}}(m_t, M_Z, s_w, \dots) + \dots \right)$$
$$\tilde{\mathcal{L}} = \frac{G_F^2 M_W^2}{\pi^2} \left(\tilde{c}_{10}^{\text{LO}} \left(\frac{m_t^2}{M_W^2} \right) + \frac{\alpha_{em}}{4\pi} \tilde{c}_{10}^{\text{EW}}(m_t, M_Z, s_w, \dots) + \dots \right)$$

- $\tilde{\mathcal{L}} \rightarrow$ no EW ambiguity at LO [Misiak '11, arXiv:1112.5978]
- only the product invariant under EW scheme

Step 2: choice of numerical input

$$G_F, \quad \alpha_{em}^{\overline{\text{MS}}}, \quad M_Z^{\text{pole}}, \quad M_H^{\text{pole}}, \quad M_t^{\text{pole}}$$

Step 3: EW renormalisation schemes (α_{em} always $\overline{\text{MS}}$)


OS masses, $s_w^2 \equiv 1 - M_W^2/M_Z^2$

$\overline{\text{MS}}$ everything, “unbroken” parameters $v, g_1, g_2, y_t, \lambda$

HY masses on-shell, $s_w^2 \overline{\text{MS}}$


Electroweak Corrections II/II

- 2-loop matching calculation in $R_\xi=1$ gauge, divergences cancel, ...



- here, switch off QCD and neglect mixing


[Bobeth, Gorbahn, ES '13, arXiv:1311.1348]



- @LO large shifts $\rightarrow \pm 8\%$ on BR
- @LO considerable μ_0 -dependence in single G_F normalisation


Electroweak Corrections II/II

- 2-loop matching calculation in $R_\xi=1$ gauge, divergences cancel, ...



- here, switch off QCD and neglect mixing

[Bobeth, Gorbahn, ES '13, arXiv:1311.1348]




- @NLO prediction aligns in diff. schemes
- large shift in OS-1, better convergence in HY

- **EW scheme uncert.**
 $\pm 8\% \rightarrow \pm 0.8\%$

Combining NLO EW & NNLO QCD + RGE to μ_b

LO + NLO QCD + NNLO QCD + Log QED + NLO EW

- Choose OS-2 as default scheme
- RGE evolution
 $C_{10}(\mu_b) = \sum_i U(\mu_b, \mu_0) C_i(\mu_0)$
- Log-enhanced QED corrections known
[Bobeth, Gambino, Gorbahn, Haisch '03 hep-ph/0312090, Huber, Lunghi, Misiak, Wyler '05, hep-ph/0512066]
- **NLO EW reduce BR by 4% w.r.t. NNLO QCD**



Estimate of higher-order uncertainties


- μ_0 variation between $[m_t/2, 2m_t] \rightarrow \pm 0.2\% \text{ (QCD)} \pm 0.2\% \text{ (EW)}$
- residual scheme dependence from OS-2 & HY $\rightarrow \pm 0.2\%$

Residual μ_b Dependence

LO + NLO QCD + NNLO QCD + Log QED + NLO EW

- the Log enhanced QED corrections further reduce the BR
- the residual μ_b dependence cancels by yet **unknown** virtual QED corrections at μ_b .
- variation of μ_b gives a measure of uncertainty

$\pm 0.3\%$ residual μ_b dependence




Soft and Hard Photons

Soft photons from muons

Theoretical prediction is fully inclusive in bremsstrahlung. Otherwise sizeable corrections from phase-space cut.


[Buras, Girrbach, Guadagnoli, Isidori '12,
arXiv:1208.0934]




Direct emission

Is phase-space suppressed for invariant mass $m_{\mu\mu}$ close to M_{B_s} .

[Aditya, Healey, Petrov '12, arXiv:1212.4166]



Soft and Hard Photons



Suppose collaborations were using simple signal windows:

(they do not, analysis more involved)

- direct emission treated as background, tiny in signal window
- simulate signal fully inclusive in bremsstrahlung (PHOTOS)

$$\mathbf{BR}(B_q \rightarrow \mu^+ \mu^-)$$

The new SM Prediction & Error Budget

$$\text{BR}(B_s \rightarrow \mu^+ \mu^-)^{\text{exp}} = (2.8^{+0.7}_{-0.6}) \times 10^{-9}$$

[new average, see talk by F. Archilli]

$$\text{BR}(B_s \rightarrow \mu^+ \mu^-)^{\text{SM}} = (3.65 \pm 0.23) \times 10^{-9}$$

[Bobeth, Gorbahn, Hermann, Misiak, ES, Steinhauser '13, arXiv:1311.0903]

Error Budget

f_{B_s}	CKM	τ_H^s	M_t	α_s	other param.	non-param.	\sum
4.0%	4.3%	1.3%	1.6%	0.1%	< 0.1%	1.5%	6.4%

param.

- $f_{B_s} = 227.4(4.5)$ MeV
[FLAG '13, arXiv:1310.8555]
- V_{cb} from recent inclusive fit
[Gambino, Schwanda '13, arXiv:1307.4551]
[Update! See talk by P. Gambino]

non param.

- 0.3% from $\mu_b \in [m_b/2, 2m_b]$
- $2 \times 0.3\%$ from $O(\alpha_s^3, \alpha_{em}^2, \alpha_s \alpha_{em})$ for $\mu_0 \in [m_t/2, 2m_t]$
- 0.3% from top-mass conversion
- 0.5% additional uncertainties ($O(m_b^2/M_W^2) + \dots$)

The new SM Prediction & Error Budget

$$\text{BR}(B_d \rightarrow \mu^+ \mu^-)^{\text{exp}} = (3.9^{+1.6}_{-1.4}) \times 10^{-10}$$

[new average, see talk by F. Archilli]

$$\text{BR}(B_d \rightarrow \mu^+ \mu^-)^{\text{SM}} = (1.06 \pm 0.09) \times 10^{-10}$$

[Bobeth, Gorbahn, Hermann, Misiak, ES, Steinhauser '13, arXiv:1311.0903]

Error Budget

f_{B_d}	CKM	τ_H^s	M_t	α_s	other param.	non-param.	\sum
4.5%	6.9%	0.5%	1.6%	0.1%	< 0.1%	1.5%	8.5%

param.

- $f_{B_d} = 190.5(4.2)$ MeV

[FLAG '13, arXiv:1310.8555]

non param.

- 0.3% from $\mu_b \in [m_b/2, 2m_b]$
- $2 \times 0.3\%$ from $O(\alpha_s^3, \alpha_{em}^2, \alpha_s \alpha_{em})$ for $\mu_0 \in [m_t/2, 2m_t]$
- 0.3% from top-mass conversion
- 0.5% additional uncertainties ($O(m_b^2/M_W^2) + \dots$)

Conclusions

We revisited and improved the SM prediction of the rare decay $B_q \rightarrow \mu^+ \mu^-$

(actually of all $B_q \rightarrow \ell^+ \ell^-$ decays)

- NNLO QCD corrections reduce μ_0 dependence from $m_t(\mu_0)$
from 1.8%@NLO QCD → 0.2%@NNLO QCD
- NLO EW corrections reduce EW scheme dependence
from 8%@LO → 0.6%@NLO EW
- the size of the NLO EW corrections is
(3 – 5)% depending on μ_0 and scheme
- the theory uncertainty of the BR is
≤ 7% mainly from f_{B_s} (4%), V_{cb} (4.3%), non-param.(1.5%)

The SM prediction is

$$\text{BR}(B_s \rightarrow \mu^+ \mu^-)^{\text{SM}} = (3.65 \pm 0.23) \times 10^{-9}$$

in good agreement with the LHCb and CMS measurements.

Backup

The Branching Ratio of $B_s \rightarrow \mu^+ \mu^-$

- LHCb, CMS report average time-integrated BR
- within the SM it is related to instantaneous BR $^{[t=0]}$:

$$\text{BR} = \frac{1}{1 - \tau_{B_s} \Delta \Gamma_s / 2} \text{BR}^{[t=0]}$$

[De Bruyn, Fleischer, Kneijens, Koppenburg, Merk '12, arXiv:1204.1737]

$$\text{BR}(B_s \rightarrow \mu^+ \mu^-) = \frac{M_{B_s}^3 f_{B_s}^2}{32\pi \Gamma_H^s} \beta r^2 |V_{tb}^* V_{ts}|^2 \left| \frac{G_F^2 M_W^2}{\pi^2} C_{10}(\mu_b) \right|^2 + O(\alpha_{em})$$

with $r = 2m_\mu/M_{B_s}$ and $\beta = \sqrt{1 - r^2}$

r^2 : helicity suppression

$1/\Gamma_H$: effect of $B_s^0 - \bar{B}_s^0$ mixing

f_{B_s} : decay constant from lattice input

$|...|^2$: perturbative Wilson coefficient