



Contribution ID: 68

Type: not specified

## **$B \rightarrow D^{**} l \nu$ - puzzle 1/2 vs 3/2**

*Wednesday 10 September 2014 10:25 (20 minutes)*

Understanding the composition of final states in  $B \rightarrow Xc l \nu$  could help to get a feedback on the persisting disagreement between exclusive and inclusive determinations of  $V_{cb}$ . In particular the series of orbital excitations  $D$  and radial excitations ( $D'$ ,  $D^{**}$ ) has received a lot of attention; a misinterpretation as a scalar state of the ( $D' \rightarrow D \pi$ ) spectrum tail could have induced an experimental overestimate of the broad states contribution to the total  $B \rightarrow Xc l \nu$  width with respect to theoretical expectations, all of them made however in the infinite mass limit: it is the so-called 1/2 vs 3/2 puzzle. We will describe first attempts to measure on the lattice form factors of  $B \rightarrow D l \nu$  at realistic quark masses. Cleaner processes, like hadronic decays  $B \rightarrow D \pi$  and semileptonic decays  $B_s \rightarrow D_s l \nu$  in the strange sector have recently been examined by phenomenologists, putting new interesting ideas on those issues with, again, the need of lattice inputs.

**Presenter:** Dr BLOSSIER, Benoit (CNRS/Laboratoire de Physique Théorique d'Orsay)

**Session Classification:** WG2