

Next-to-eikonal corrections in the CGC

Tolga Altinoluk

Universidade de Santiago de Compostela

DIS - 2014 Warsaw, Poland

April 30, 2014

[T. A. , N. Armesto, G. Beuf, M. Martínez, C. Salgado, arXiv:1404.2219]

Eikonal approximation for high-energy (dense-dilute) scattering


- dense target \equiv classical background field $A_a^\mu(x)$
- dilute projectile \equiv color charge $J_a^\mu(x)$

Coupling between the projectile and the target $\rightarrow \int d^4x J_a^\mu(x) A_a^\mu(x)$

In the high energy limit (within eikonal approximation):

- boosting the projectile:
 - $J_a^+ \gg J_a^i \gg J_a^-$
 - $J_a^\mu(x) = \delta^{\mu+} J^+(x) \propto \delta(x^-) \Rightarrow$ projectile localized at $x^- = 0$
- boosting the target:
 - $A_a^- \gg A_a^i \gg A_a^+$
 - $A^\mu(x) \equiv \delta^{\mu-} A_a^-(x^+, \mathbf{x}) \propto \delta(x^+) \Rightarrow$ target localized at $x^+ = 0$
& independent of x^-

Finite width target : relaxing the eikonal approximation



Consider a finite width target, but still assuming that

- $J_a^\mu(x) \propto \delta^{\mu+} \delta(x^-)$
- $A^\mu(x) \equiv \delta^{\mu-} A_a^-(x^+, x_\perp)$

The projectile $\rightarrow j_a^\mu(x) = \delta^{\mu+} \delta(x^-) \mathcal{U}^{ab}(x^+, 0, \mathbf{x}) \rho^b(\mathbf{x} - \mathbf{B})$

$$\mathcal{P}_+ \exp \left\{ ig \int_0^{x^+} dz^+ T \cdot \mathcal{A}^-(z^+, \mathbf{x}) \right\}^{ab}$$

The single inclusive gluon spectrum for pA:


$$(2\pi)^3 (2k^+) \frac{dN}{dk^+ d^2\mathbf{k}}(\mathbf{B}) = \sum_{\lambda \text{ phys.}} \left\langle \left\langle |\mathcal{M}_\lambda^a(\underline{k}, \mathbf{B})|^2 \right\rangle_\rho \right\rangle_A$$

gluon production amplitude

The single inclusive gluon cross section :

$$\frac{d\sigma}{dk^+ d^2\mathbf{k}} = \int d^2\mathbf{B} \frac{dN}{dk^+ d^2\mathbf{k}}(\mathbf{B})$$

$$\mathcal{M}_\lambda^a(\underline{k}, \mathbf{B}) =$$



at LO in g , LSZ reduction formula $\Rightarrow \mathcal{M}_\lambda^a(\underline{k}, \mathbf{B}) = \varepsilon_\lambda^{\mu*} \int d^4x e^{ik \cdot x} \square_x A_\mu^a(x)$

power counting : dense target $\Rightarrow \mathcal{A}_a^-(x^+, \mathbf{x}) = \mathcal{O}(1/g)$

dilute projectile $\Rightarrow j_a^+(\underline{x}) = \mathcal{O}(g)$

perturbative expansion of the classical field : $A_a^\mu(x) = \mathcal{A}_a^\mu(x) + a_a^\mu(x) + \mathcal{O}(g^3)$

$$\downarrow$$

$$-i \int d^4y G_R^{\mu-}(x, y)_{ab} j_b^+(y)$$

In the LC gauge : $A_a^+ = 0$ & $\varepsilon_\lambda^{+*} = 0 \Rightarrow \varepsilon_\lambda^{\mu*} \mathcal{A}_\mu^a(x) = -\varepsilon_\lambda^{i*} a_a^i$

$$\mathcal{M}_\lambda^a(\underline{k}, \mathbf{B}) = \varepsilon_\lambda^{i*} (2k^+) \lim_{x^+ \rightarrow +\infty} \int d^2\mathbf{x} \int dx^- e^{ik \cdot x} \int d^4y G_R^{i-}(x, y)_{ab} j_b^+(y)$$

$G_R^{\mu\nu}(x, y)_{ab}$ is the background retarded gluon propagator

Properties of the background retarded gluon propagator

$G_R^{\mu\nu}(x, y)_{ab}$ is the retarded solution of the linearized Yang-Mills equations around the classical field $\mathcal{A}_a^-(x^+, \mathbf{x})$:

$$\left[g_{\mu\nu} \left(\delta_{ab} \square_x - 2ig \left(\mathcal{A}^-(x^+, \mathbf{x}) \cdot T \right)_{ab} \partial_{x^-} \right) - \delta_{ab} \partial_{x^\mu} \partial_{x^\nu} \right] G_R^{\nu\rho}(x, y)_{bc} = i g_{\mu}{}^\rho \delta_{ac} \delta^{(4)}(x-y)$$

$\mathcal{A}_a^-(x^+, \mathbf{x})$ is independent of the x^- , so it is convenient to introduce the 1-d Fourier transform of $G_R^{\mu\nu}(x, y)_{ab}$

$$G_R^{\mu\nu}(x, y)_{ab} = \int \frac{dk^+}{2\pi} e^{-ik^+(x^- - y^-)} \frac{1}{2(k^+ + i\epsilon)} \mathcal{G}_{k^+}^{\mu\nu}(\underline{x}; \underline{y})_{ab}$$

The ($i-$) component of the $\mathcal{G}_{k^+}^{\mu\nu}(\underline{x}; \underline{y})_{ab}$ can be written as

$$\mathcal{G}_{k^+}^{i-}(\underline{x}; \underline{y})^{ab} = \frac{i}{k^+ + i\epsilon} \partial_{\mathbf{y}^i} \mathcal{G}_{k^+}^{ab}(\underline{x}; \underline{y})$$

$\mathcal{G}_{k^+}^{ab}(\underline{x}; \underline{y})$ is the background scalar propagator.

The background scalar propagator satisfies the scalar Green's equation :

$$\left[\delta^{ab} \left(i\partial_{x^+} + \frac{\partial_{\underline{x}}^2}{2(k^+ + i\epsilon)} \right) + g \left(\mathcal{A}^-(\underline{x}) \cdot T \right)^{ab} \right] \mathcal{G}_{k^+}^{bc}(\underline{x}; \underline{y}) = i \delta^{ac} \delta^{(3)}(\underline{x} - \underline{y})$$

Schrödinger equation in 2 + 1 dimensions in a space-time dependent matrix potential $-g \left(\mathcal{A}^-(\underline{x}) \cdot T \right)^{ab}$.

The solution can be written formally as a path integral

$$\mathcal{G}_{k^+}^{ab}(\underline{x}; \underline{y}) = \theta(x^+ - y^+) \int_{\mathbf{z}(y^+) = \underline{y}}^{\mathbf{z}(x^+) = \underline{x}} \mathcal{D}\mathbf{z}(z^+) \exp \left[\frac{ik^+}{2} \int_{y^+}^{x^+} dz^+ \dot{\mathbf{z}}^2(z^+) \right] \\ \times \mathcal{U}^{ab}(x^+, y^+, [\mathbf{z}(z^+)])$$

with the Wilson line

$$\mathcal{U}^{ab}(x^+, y^+, [\mathbf{z}(z^+)]) = \mathcal{P}_+ \exp \left\{ ig \int_{y^+}^{x^+} dz^+ T \cdot \mathcal{A}^-(z^+, \mathbf{z}(z^+)) \right\}^{ab}$$

following the Brownian trajectory $\mathbf{z}(z^+)$.

Discretization of the background propagator

Discretized form of the background propagator :

$$\mathcal{G}_{k^+}^{ab}(\underline{x}; \underline{y}) = \lim_{N \rightarrow +\infty} \theta(x^+ - y^+) \int \left(\prod_{n=1}^{N-1} d^2 \mathbf{z}_n \right) \left(\frac{-i(k^+ + i\epsilon)N}{2\pi(x^+ - y^+)} \right)^N \\ \times \exp \left[\frac{i(k^+ + i\epsilon)N}{2(x^+ - y^+)} \sum_{n=0}^{N-1} (\mathbf{z}_{n+1} - \mathbf{z}_n)^2 \right] \mathcal{U}^{ab}(x^+, y^+, \{\mathbf{z}_n\})$$


with the discretized Wilson line

$$\mathcal{U}^{ab}(x^+, y^+, \{\mathbf{z}_n\}) = \mathcal{P}_+ \left\{ \prod_{n=0}^{N-1} \exp \left[ig \frac{(x^+ - y^+)}{N} (\mathcal{A}^-(z_n^+, \mathbf{z}_n) \cdot T) \right] \right\}^{ab}$$

with $z_n^+ = y^+ + \frac{n}{N}(x^+ - y^+)$

Expansion of the background propagator

Perturbative expansion around free classical path:



eikonal limit : $\frac{k^+}{(x^+ - y^+)} \gg Q_1^2$ in the problem

\mapsto large k^+ limit (classical free path!)

\Rightarrow perturbative expansion around the free classical path:

$$\mathbf{z}_n = \mathbf{z}_n^{\text{cl}} + \mathbf{u}_n \text{ with } \mathbf{z}_n^{\text{cl}} = \mathbf{y} + \frac{n}{N}(\mathbf{x} - \mathbf{y})$$

Then


$$\mathcal{G}_{k^+}^{ab}(\underline{x}; \underline{y}) = 2\pi i \mathcal{G}_{0, k^+}(\underline{x}; \underline{y}) \frac{(x^+ - y^+)}{k^+} \lim_{N \rightarrow +\infty} \int \left(\prod_{n=1}^{N-1} d^2 \mathbf{u}_n \right) \times \mathcal{P}_+ \prod_{n=0}^{N-1} \left\{ \mathcal{G}_{0, k^+}(\mathbf{z}_{n+1}^+, \mathbf{u}_{n+1}; \mathbf{z}_n^+, \mathbf{u}_n) \exp \left[\frac{(x^+ - y^+)}{N} igT \cdot \mathcal{A}^-(\mathbf{z}_n^+, \mathbf{z}_n^{\text{cl}} + \mathbf{u}_n) \right] \right\}$$

- Taylor expand the gauge link around $\mathbf{u}_n = 0$ at each discretization step.
- Take the product of the Taylor expansion of all links and collect terms according to power of \mathbf{u}_n .

Expansion around the initial transverse position :

The first expansion is performed for fixed initial and final positions.

In the large k^+ limit , the result has to be re-expanded since $\mathbf{z}^{\text{cl}}(z^+) - \mathbf{y}$ is small at each step.



After all :

$$\int d^2\mathbf{x} e^{-ik\cdot\mathbf{x}} \mathcal{G}_{k^+}^{ab}(\underline{x}; \underline{y}) = \theta(x^+ - y^+) e^{-ik\cdot\mathbf{y}} e^{-ik^-(x^+ - y^+)} \left\{ \mathcal{U}(x^+, y^+, \mathbf{y}) \right. \\ \left. + \frac{(x^+ - y^+)}{k^+} \mathbf{k}^i \mathcal{U}_{(1)}^i(x^+, y^+, \mathbf{y}) + i \frac{(x^+ - y^+)}{2k^+} \mathcal{U}_{(2)}(x^+, y^+, \mathbf{y}) \right\}^{ab}$$

strict eikonal term
first corrections to strict eikonal limit

The explicit form of the corrections :

$$\mathcal{U}_{(1)}^{i,ab}(x^+, y^+, \mathbf{y}) = \int_{y^+}^{x^+} dz^+ \left(\frac{z^+ - y^+}{x^+ - y^+} \right) \left\{ \mathcal{U}(x^+, z^+, \mathbf{y}) \right. \\ \left. \times [igT \cdot \partial_{\mathbf{y}^i} A^-(z^+, \mathbf{y})] \mathcal{U}(z^+, y^+, \mathbf{y}) \right\}^{ab}$$

and

$$\mathcal{U}_{(2)}^{ab}(x^+, y^+, \mathbf{y}) = \int_{y^+}^{x^+} dz^+ \left(\frac{z^+ - y^+}{x^+ - y^+} \right) \left\{ \mathcal{U}(x^+, z^+, \mathbf{y}) [igT \cdot \partial_{\mathbf{y}^i}^2 A^-(z^+, \mathbf{y})] \mathcal{U}(z^+, y^+, \mathbf{y}) \right\}^{ab} \\ + 2 \int_{y^+}^{x^+} dz^+ \int_{y^+}^{z^+} dw^+ \left(\frac{w^+ - y^+}{x^+ - y^+} \right) \left\{ \mathcal{U}(x^+, z^+, \mathbf{y}) [igT \cdot \partial_{\mathbf{y}^i} A^-(z^+, \mathbf{y})] \right. \\ \left. \times \mathcal{U}(z^+, w^+, \mathbf{y}) [igT \cdot \partial_{\mathbf{y}^i} A^-(w^+, \mathbf{y})] \mathcal{U}(w^+, y^+, \mathbf{y}) \right\}^{ab}$$

Reduced amplitude and the cross section

Defining Fourier transform of the color charge density

$$\rho^a(\mathbf{y} - \mathbf{B}) = \int \frac{d^2\mathbf{q}}{(2\pi)^2} e^{i\mathbf{q}\cdot(\mathbf{y}-\mathbf{B})} \tilde{\rho}^a(\mathbf{q})$$

and gluon-nucleus reduced amplitude

$$\mathcal{M}_\lambda^a(\underline{k}, \mathbf{B}) = \int \frac{d^2\mathbf{q}}{(2\pi)^2} e^{-i\mathbf{q}\cdot\mathbf{B}} \overline{\mathcal{M}}_\lambda^{ab}(\underline{k}, \mathbf{q}) \tilde{\rho}^b(\mathbf{q})$$

The cross section is written as

$$k^+ \frac{d\sigma}{dk^+ d^2\mathbf{k}} = \int \frac{d^2\mathbf{q}}{(2\pi)^2} \varphi_p(\mathbf{q}) \frac{\mathbf{q}^2}{4} \frac{1}{N_c^2 - 1} \sum_{\lambda \text{ phys.}} \left\langle \overline{\mathcal{M}}_\lambda^{ab}(\underline{k}, \mathbf{q})^\dagger \overline{\mathcal{M}}_\lambda^{ab}(\underline{k}, \mathbf{q}) \right\rangle_A.$$

Expanded reduced amplitude reads

$$\begin{aligned} \overline{\mathcal{M}}_\lambda^{ab}(\underline{k}, \mathbf{q}) &= i \varepsilon_\lambda^{i*} \int d^2\mathbf{y} e^{-i(\mathbf{k}-\mathbf{q})\cdot\mathbf{y}} \left\{ 2 \left[\frac{\mathbf{k}^i}{\mathbf{k}^2} - \frac{\mathbf{q}^i}{\mathbf{q}^2} \right] \mathcal{U}(L^+, 0; \mathbf{y}) \right. \\ &\left. + \frac{L^+}{k^+} \left[\delta^{ij} - 2 \frac{\mathbf{q}^i \mathbf{k}^j}{\mathbf{q}^2} \right] \mathcal{U}_{(1)}^j(L^+, 0; \mathbf{y}) - i \frac{L^+}{k^+} \frac{\mathbf{q}^i}{\mathbf{q}^2} \mathcal{U}_{(2)}^i(L^+, 0; \mathbf{y}) + \mathcal{O} \left(\left(\frac{L^+}{k^+} \partial_\perp^2 \right)^2 \right) \right\}^{ab} \end{aligned}$$

By defining the new operators as

$$S_A(\mathbf{r}, \mathbf{b}) = \frac{1}{N_{\epsilon}^2 - 1} \left\langle \text{tr} \left[\mathcal{U}^\dagger(L^+, 0; \mathbf{b} - \frac{\mathbf{r}}{2}) \mathcal{U}(L^+, 0; \mathbf{b} + \frac{\mathbf{r}}{2}) \right] \right\rangle_A$$

$$\mathcal{O}_{(1)}^j(\mathbf{r}, \mathbf{b}) = \frac{1}{N_{\epsilon}^2 - 1} \left\langle \text{tr} \left[\mathcal{U}^\dagger(L^+, 0; \mathbf{b} - \frac{\mathbf{r}}{2}) \mathcal{U}_{(1)}^j(L^+, 0; \mathbf{b} + \frac{\mathbf{r}}{2}) \right] \right\rangle_A$$

$$\mathcal{O}_{(2)}(\mathbf{r}, \mathbf{b}) = \frac{1}{N_{\epsilon}^2 - 1} \left\langle \text{tr} \left[\mathcal{U}^\dagger(L^+, 0; \mathbf{b} - \frac{\mathbf{r}}{2}) \mathcal{U}_{(2)}(L^+, 0; \mathbf{b} + \frac{\mathbf{r}}{2}) \right] \right\rangle_A$$

Square of the reduced amplitude then reads

$$\frac{1}{N_{\epsilon}^2 - 1} \sum_{\lambda \text{ phys.}} \left\langle \overline{\mathcal{M}}_{\lambda}^{ab}(\underline{k}, \mathbf{q})^\dagger \overline{\mathcal{M}}_{\lambda}^{ab}(\underline{k}, \mathbf{q}) \right\rangle_A = \frac{1}{\mathbf{k}^2 \mathbf{q}^2} \int_{\mathbf{b}, \mathbf{r}} e^{-i(\mathbf{k}-\mathbf{q})\cdot\mathbf{r}} \left\{ 4(\mathbf{k}-\mathbf{q})^2 S_A(\mathbf{r}, \mathbf{b}) \right.$$

$$\left. + \frac{L^+}{k^+} \left[f(\mathbf{k}^i, \mathbf{q}^j) \underbrace{[\mathcal{O}_{(1)}^j(\mathbf{r}, \mathbf{b}) + \mathcal{O}_{(1)}^j(-\mathbf{r}, \mathbf{b})]} + g(\mathbf{k}, \mathbf{q}) \underbrace{[\mathcal{O}_{(2)}(\mathbf{r}, \mathbf{b}) - \mathcal{O}_{(2)}(-\mathbf{r}, \mathbf{b})]} \right] \right\}$$

Vanish upon integration over \mathbf{b} due to rotational symmetry

$$k^+ \frac{d\sigma}{dk^+ d^2\mathbf{k}} = \frac{1}{\mathbf{k}^2} \int \frac{d^2\mathbf{q}}{(2\pi)^2} \varphi_p(\mathbf{q}) (\mathbf{k}-\mathbf{q})^2 \int_{\mathbf{b}, \mathbf{r}} e^{-i(\mathbf{k}-\mathbf{q})\cdot\mathbf{r}} S_A(\mathbf{r}, \mathbf{b}) + \mathcal{O}\left(\left(\frac{L^+}{k^+} \partial_{\perp}^2\right)^2\right)$$

Well known k_{\perp} factorization formula!!

Single transverse spin asymmetry : polarized target

Consider the process $p + A^\uparrow \rightarrow g + X$. The SSA is defined as

$$A_N = \frac{k^+ \frac{d\sigma^\uparrow}{dk^+ d^2\mathbf{k}} - k^+ \frac{d\sigma^\downarrow}{dk^+ d^2\mathbf{k}}}{k^+ \frac{d\sigma^\uparrow}{dk^+ d^2\mathbf{k}} + k^+ \frac{d\sigma^\downarrow}{dk^+ d^2\mathbf{k}}} = \frac{k^+ \frac{d\sigma^\uparrow}{dk^+ d^2\mathbf{k}} - k^+ \frac{d\sigma^\downarrow}{dk^+ d^2\mathbf{k}}}{2k^+ \frac{d\sigma}{dk^+ d^2\mathbf{k}}}$$

Transversely polarized target \Rightarrow \mathbf{s} dependent S_A , $\mathcal{O}_{(1)}^j$ and $\mathcal{O}_{(2)}$

Due to transverse rotational symmetry around the center of the target:

$$S_A(-\mathbf{r}, -\mathbf{b}, -\mathbf{s}) = S_A(\mathbf{r}, \mathbf{b}, \mathbf{s})$$

$$\mathcal{O}_{(1)}^j(-\mathbf{r}, -\mathbf{b}, -\mathbf{s}) = -\mathcal{O}_{(1)}^j(\mathbf{r}, \mathbf{b}, \mathbf{s})$$

$$\mathcal{O}_{(2)}(-\mathbf{r}, -\mathbf{b}, -\mathbf{s}) = \mathcal{O}_{(2)}(\mathbf{r}, \mathbf{b}, \mathbf{s}) \text{ Then:}$$

$$k^+ \left(\frac{d\sigma^\uparrow}{dk^+ d^2\mathbf{k}} - \frac{d\sigma^\downarrow}{dk^+ d^2\mathbf{k}} \right) = \frac{L^+}{k^+} \int_{\mathbf{q}\mathbf{r}\mathbf{b}} \varphi_p(\mathbf{q}) \left\{ f(\mathbf{k}^j, \mathbf{q}^j, \mathbf{r}) \mathcal{O}_{(1)}^j(\mathbf{r}, \mathbf{b}, \mathbf{s}) + g(\mathbf{k}, \mathbf{q}, \mathbf{r}) \mathcal{O}_{(2)}(\mathbf{r}, \mathbf{b}, \mathbf{s}) \right\}$$

The strict eikonal term vanishes!!

Next-to-eikonal terms in the expansion become leading terms!!

Analogy between twist-3 contributions to hard processes and next-to-eikonal contributions to high-energy processes??

Polarized gluon production in unpolarized pA

Consider a polarized gluon produced in unpolarized pA which then which then fragments in to polarized hadron preserving this asymmetry.

This asymmetry is given by difference between $\lambda = \pm 1$ helicity contributions of the produced gluon:

$$k^+ \frac{d\sigma^+}{dk^+ d^2\mathbf{k}} - k^+ \frac{d\sigma^-}{dk^+ d^2\mathbf{k}} = \int \frac{d^2\mathbf{q}}{(2\pi)^2} \varphi_P(\mathbf{q}) \frac{\mathbf{q}^2}{4} \frac{1}{N_c^2 - 1} \sum_{\lambda \text{ phys.}} \lambda \left\langle \overline{\mathcal{M}}_\lambda^{ab}(\underline{k}, \mathbf{q})^\dagger \overline{\mathcal{M}}_\lambda^{ab}(\underline{k}, \mathbf{q}) \right\rangle_A$$

By using the identity

$$\sum_{\lambda \text{ phys.}} \lambda \varepsilon_\lambda^i \varepsilon_\lambda^{*j} = -i \epsilon^{ij}$$

one gets

$$k^+ \frac{d\sigma^+}{dk^+ d^2\mathbf{k}} - k^+ \frac{d\sigma^-}{dk^+ d^2\mathbf{k}} = \frac{L^+}{k^+} \int \frac{d^2\mathbf{q}}{(2\pi)^2} \varphi_P(\mathbf{q}) \mathbf{q}^2 \int_{\mathbf{b}, \mathbf{r}} e^{-i(\mathbf{k}-\mathbf{q})\cdot\mathbf{r}} \\ \times \left\{ -i \left[\left(\frac{\mathbf{k}^i}{k^2} - \frac{\mathbf{q}^i}{q^2} \right) \epsilon^{ij} - 2 \frac{(\epsilon^{il} \mathbf{k}^l \mathbf{q}^j)}{k^2 q^2} \mathbf{k}^j \right] \mathcal{O}_{(1)}^j(\mathbf{r}, \mathbf{b}) - \frac{(\epsilon^{ij} \mathbf{k}^i \mathbf{q}^j)}{k^2 q^2} \mathcal{O}_{(2)}(\mathbf{r}, \mathbf{b}) \right\}$$

Strict eikonal terms vanish leaving the next-to-eikonal terms as leading contribution!!!

Conclusions and Outlook

- We develop a systematic eikonal expansion of the retarded gluon propagator in a background field.
- We apply this method to single inclusive gluon cross section in order to study the corrections to the CGC beyond eikonal limit. The strict eikonal term provides the usual k_{\perp} -factorization formula, whereas the first next-to-eikonal corrections vanish for this particular observable.
- For SSA and polarized gluon production the strict eikonal terms vanish leaving the next-to-eikonal corrections as the leading contribution.
- The first non-vanishing contribution to strict eikonal limit of single inclusive gluon cross section arises at order $(L^+ Q_s^2/k^+)^2$ which can be obtained by continuing the expansion to higher orders. [In preparation]
- The same expansion can be applied to energy loss calculations in jet quenching physics. [In preparation]