



Contribution ID: 88

Type: Oral presentation

Search for QCD Instantons using the H1 experiment at HERA

Tuesday 29 April 2014 10:05 (20 minutes)

Signals of QCD instanton-induced processes are searched for in deep-inelastic scattering (DIS) at the electron-proton collider HERA in the kinematic region defined by the Bjorken-scaling variable $x > 10^{-3}$, the inelasticity $0.2 < y < 0.7$ and the photon virtuality $150 < Q^2 < 15000 \text{ GeV}^2$. The search is performed using H1 data corresponding to an integrated luminosity of 350 pb^{-1} . Several observables of the hadronic final state of the events are exploited to identify a potentially instanton-enriched domain. Two Monte Carlo models, RAPGAP and ARLADNE, are used to estimate the background from the standard DIS processes, and the instanton-induced scattering processes are modeled by the program QCDINS. In order to extract the expected signal a multivariate data analysis technique is used.

Primary authors: DAUM, Karin (University of Wuppertal/DESY); SCHMITT, Stefan (Deutsches Elektronen-Synchrotron (DE))

Presenter: SOPICKI, Pawel (IFJ PAN)

Session Classification: WG4: QCD and Hadronic Final States

Track Classification: WG4: QCD and Hadronic Final States