



# Exclusive Photoproduction of $\rho^0$ Meson with Forward Neutron at HERA



Sergey Levonian

*On behalf of H1 Collaboration*




# Introduction

- First observation of exclusive photoproduction on (virtual) pion

- ▷ Unique for HERA (before that  $\gamma, \pi$  beams did exist, but no target)
- ▷ Extends further (very powerful) VM field at HERA

- Key observables:

- ▷  $x_L = E_n/E_p$  (or  $x_\pi = 1 - x_L$ ) distribution:  $\sim f_{\pi/p}(x_L)$
- ▷  $W$  dependence:  $\sim W^\delta$  – nature of exchange object(s)
- ▷  $t$ -slope of  $\rho^0$  ( $b \propto R^2$  in geometric picture)



- Main experimental difficulty:

- ▷ Trigger (tagged  $\gamma p$  – too large  $W$  to observe VM; untagged  $\gamma p$  – too high rates/prescales)
- ▷ Limited acceptance for forward  $\pi$  and  $N$  ( $\eta_{\text{lab}} \geq 6$ )

# Introduction

- First observation of exclusive photoproduction on (virtual) pion

- ▷ Unique for HERA (before that  $\gamma, \pi$  beams did exist, but no target)
- ▷ Extends further (very powerful) VM field at HERA

- Key observables:


- ▷  $x_L = E_n/E_p$  (or  $x_\pi = 1 - x_L$ ) distribution:  $\sim f_{\pi/p}(x_L)$
- ▷  $W$  dependence:  $\sim W^\delta$  – nature of exchange object(s)
- ▷  $t$ -slope of  $\rho^0$  ( $b \propto R^2$  in geometric picture)

- Main experimental difficulty:

- ▷ Trigger (tagged  $\gamma p$  – too large  $W$  to observe VM; untagged  $\gamma p$  – too high rates/prescales)
- ▷ Limited acceptance for forward  $\pi$  and  $N$  ( $\eta_{\text{lab}} \geq 6$ )

- Advantages of H1@HERA2

- ▷ Improved FNC (distinguish and measure  $n$  and  $\gamma/\pi^0$ )



# Introduction

- First observation of exclusive photoproduction on (virtual) pion

- ▷ Unique for HERA (before that  $\gamma, \pi$  beams did exist, but no target)
- ▷ Extends further (very powerful) VM field at HERA

- Key observables:


- ▷  $x_L = E_n/E_p$  (or  $x_\pi = 1 - x_L$ ) distribution:  $\sim f_{\pi/p}(x_L)$
- ▷  $W$  dependence:  $\sim W^\delta$  – nature of exchange object(s)
- ▷  $t$ -slope of  $\rho^0$  ( $b \propto R^2$  in geometric picture)

- Main experimental difficulty:

- ▷ Trigger (tagged  $\gamma p$  – too large  $W$  to observe VM; untagged  $\gamma p$  – too high rates/prescales)
- ▷ Limited acceptance for forward  $\pi$  and  $N$  ( $\eta_{\text{lab}} \geq 6$ )



- Advantages of H1@HERA2

- ▷ Improved FNC (distinguish and measure  $n$  and  $\gamma/\pi^0$ )
- ▷ Powerful fast track trigger (allows untagged soft  $\gamma p$  to be collected)





# Contributing processes and their modelling



$$W_{\gamma p} \simeq \sqrt{2(E - p_z)_\rho E_p}$$

$$W_{\gamma\pi} \simeq W_{\gamma p} \sqrt{1 - x_L}$$

- DPP expectations:  $f_{\pi/p}(x_L, t) \Rightarrow x_L$  shape,  $p_{t,\rho}^2$  slope,  $b = b_{\text{eff}}(M_{\pi N})$

- Diffractive bgr is well known (but has an irreducible part:  $M_Y = N^* \rightarrow n\pi^+$ )

# Analysis Summary

## • Data sample

- ▷ 2006 – 2007  $e^+$  runs,  $\sqrt{s} = 319$  GeV,  $\mathcal{L} = 1.16$  pb $^{-1}$   $\simeq$  6600 events in final sample
- ▷ Trigger:  $\langle \epsilon_{L1} \rangle \simeq 0.8$ ,  $\langle \epsilon_{L2} \rangle \simeq 1.0$

## • Tracking

- ▷ 2 tracks with  $p_t^{\text{tr}} > 0.2$  GeV,  $20^\circ < \theta^{\text{tr}} < 160^\circ$  fitted to event vertex  $|z_{vx}| < 30$  cm, net charge = 0
- ▷ Effective mass range:  $0.6 < M_{\pi\pi} < 1.1$  GeV (analysis);  $\Rightarrow \sigma(\rho^0)$  for  $0.28 < M_{\pi\pi} < 1.5$  GeV

## • FNC

- ▷ High energy neutron,  $E_n > 120$  GeV, within good acceptance region:  $\theta_n < 0.75$  mrad
- ▷ Background fraction determined from  $x_L$  shape:  $F_{\text{bg}} = 0.36 \pm 0.06$  (subtracted from the data)

## • Exclusivity


- ▷ Nothing above noise level in the detector except two tracks from  $\rho^0$  decay and the leading neutron

## • Cross section measurement phase space and precision

- ▷ Photoproduction:  $Q^2 < 2$  GeV $^2$ ,  $20 < W_{\gamma p} < 100$  GeV
- ▷ Leading neutron:  $0.35 < x_L < 0.95$ ,  $p_{t,n} < x_L \cdot 0.69$  GeV
- ▷  $\rho^0$  meson:  $0.28 < M_{\pi\pi} < 1.5$  GeV,  $p_{t,\rho} < 1$  GeV

$$\delta_{\text{stat}} = 2.1\% \oplus \delta_{\text{sys}} = 15.5\% \oplus \delta_{\text{norm}} = 5.9\% \Rightarrow \delta_{\text{tot}} = 16.6\%$$

# $\rho$ -meson shape




$$\frac{dN(M_{\pi\pi})}{dM_{\pi\pi}} \propto BW_{\rho}(M_{\pi\pi}) \left(\frac{M_{\rho}}{M_{\pi\pi}}\right)^{n_{RS}}$$

$$M = 764 \pm 3 \text{ MeV}$$

$$\Gamma = 154 \pm 5 \text{ MeV}$$


$$n_{RS} = 4.17 \pm 0.27$$



Analysis region:  $0.6 < M_{\pi^+\pi^-} < 1.1$  GeV extrapolated using BW to the full range:  $0.28 < M_{\rho^0} < 1.5$  GeV

# Control Plots for basic kinematics

Exclusive photoproduction of  $\rho^0$  with Forward Neutrons



Data points are shown with stat. errors only; green band represents estimated bgr fraction uncertainty


# OPE and pion fluxes

$$\frac{d^2\sigma_{\gamma p}(W^2, x_L, t)}{dx_L dt} = f_{\pi/p}(x_L, t)\sigma_{\gamma\pi}((1-x_L)W^2)$$

$$\frac{d\sigma_{\gamma p}}{dx_L} = \int_{t_0(x_L)}^{t_{min}(x_L)} f_{\pi/p}(x_L, t) dt \cdot \sigma_{\gamma\pi}(W_{\gamma\pi})$$

$$\text{where } t = -\frac{p_{t,n}^2}{x_L} - \frac{(1-x_L)(m_n^2 - m_p^2 x_L)}{x_L}$$

$$\sigma_{\gamma\pi}(W_{\gamma\pi}) = \frac{1}{\Gamma_\pi(x_L)} \frac{d\sigma_{\gamma p}}{dx_L} \quad \text{and} \quad \overline{\sigma_{\gamma\pi}(\langle W_{\gamma\pi} \rangle)} = \frac{\sigma_{\gamma p}}{\int \Gamma_\pi}$$




# OPE and pion fluxes

$$\frac{d^2\sigma_{\gamma p}(W^2, x_L, t)}{dx_L dt} = f_{\pi/p}(x_L, t)\sigma_{\gamma\pi}((1-x_L)W^2)$$

$$\frac{d\sigma_{\gamma p}}{dx_L} = \int_{t_0(x_L)}^{t_{min}(x_L)} f_{\pi/p}(x_L, t) dt \cdot \sigma_{\gamma\pi}(W_{\gamma\pi})$$

$$\text{where } t = -\frac{p_{t,n}^2}{x_L} - \frac{(1-x_L)(m_n^2 - m_p^2 x_L)}{x_L}$$

$$\sigma_{\gamma\pi}(W_{\gamma\pi}) = \frac{1}{\Gamma_\pi(x_L)} \frac{d\sigma_{\gamma p}}{dx_L} \quad \text{and} \quad \overline{\sigma_{\gamma\pi}(\langle W_{\gamma\pi} \rangle)} = \frac{\sigma_{\gamma p}}{\int \Gamma_\pi}$$



Typical examples:


$$f_{\pi^+/p}(x_L, t) = \frac{1}{2\pi} \frac{g_{p\pi N}^2}{4\pi} (1-x_L) \frac{-t}{(m_\pi^2 - t)^2} \exp[-R_{\pi n}^2 \frac{m_\pi^2 - t}{1-x_L}]$$

— H. Holtmann et al., *Nucl. Phys.* **A596** (1996) 631.

$$f_{\pi^+/p}(x_L, t) = \frac{1}{2\pi} \frac{g_{p\pi N}^2}{4\pi} (1-x_L)^{1-2\alpha'_\pi t} \frac{-t}{(m_\pi^2 - t)^2} \exp[-R_\pi^2 (m_\pi^2 - t)]$$


— B. Kopeliovich et al., *Z. Phys.* **C73** (1996) 125.

Problem: too many different fluxes on the market




# Pion fluxes confronted with H1 data

Make restricted selection of  $\pi$ -fluxes on the basis of shape comparison only



Example of fluxes **excluded by the data**  
(too soft pions 'in the proton')



Fluxes **compatible with H1 data**  
( $\chi^2 = 2.1$  to 5.5 for 6 points)

# Total cross sections



$$\sigma_{\gamma p} = \frac{\sigma_{ep}}{\int f_{\gamma/e}(y, Q^2) dy dQ^2} = \frac{N_{\text{data}} - N_{\text{bgr}}}{\mathcal{L}(A \cdot \epsilon) \mathcal{F}} \cdot C_{\rho}$$

Where

$N_{\text{bgr}}$  – diffractive dissociation bgr from MC

$\mathcal{L}$  – integrated luminosity


$A \cdot \epsilon$  – correction for detector acceptance and efficiency

$\mathcal{F}$  – photon flux integrated over kinematic domain  $20 < W < 100$  GeV,  $Q^2 < 2$  GeV<sup>2</sup>

$C_{\rho}$  – numerical factor accounting for extrapolation to full  $\rho^0$  mass range

For OPE dominated range,  $0.35 < x_L < 0.95$ , and  $20 < W_{\gamma p} < 100$  GeV,  $\theta_n < 0.75$  mrad

$$\sigma(\gamma p \rightarrow \rho^0 n(\pi^+)) = (280 \pm 6_{\text{stat}} \pm 46_{\text{sys}}) \text{ nb}$$



$$\sigma_{\gamma\pi}(\langle W_{\gamma\pi} \rangle) = \frac{\sigma_{\gamma p}}{\int f_{\pi^+/p}(x_L, t) dx_L dt},$$

and for  $\langle W_{\gamma\pi} \rangle = 22$  GeV


$$\sigma_{\text{el}}(\gamma\pi^+ \rightarrow \rho^0\pi^+) = (2.03 \pm 0.34_{\text{exp}} \pm 0.51_{\text{model}}) \mu\text{b}$$

Taking interpolated value of  $\sigma(\gamma p \rightarrow \rho^0 p) = 9.5 \pm 0.5 \mu\text{b}$  at corresponding energy, we obtain


$$r_{\text{el}} = \sigma_{\gamma\pi}^{\text{el}} / \sigma_{\gamma p}^{\text{el}} = 0.21 \pm 0.06 \quad (\text{cf. } r_{\text{tot}} = \sigma_{\gamma\pi}^{\text{tot}} / \sigma_{\gamma p}^{\text{tot}} = 0.32 \pm 0.03 \text{ [ZEUS, 2002]})$$

# Total $\gamma p$ and $\gamma\pi$ cross sections

Inner error bars – statistical uncertainty  
 outer error bars –  $\sqrt{\text{stat}^2 + \text{sys}^2}$



Inner error bars – total experimental uncertainty  
 outer error bars –  $\sqrt{\text{exp}^2 + \text{model}^2}$




Regge motivated power law fit  $W^\delta$  yields  $\delta < 0$

(in qualitative agreement with DPP and in contrast to MC,  
 $\delta_{MC} = 0.08 \pm 0.02$ , which is expected from purely  $\mathbb{P}$  exchange)

Holtmann flux is used for the central values.


Conservative model uncertainty  $\sim 25\%$

# Differential cross sections in $\eta$





Different energy dependence in data and MC is also reflected in  $\eta$  shape

# Differential cross section in $p_t^2$






# Differential cross section in $p_t^2$




Geometric interpretation:  $\langle r^2 \rangle = 2b_1 \cdot (\hbar c)^2 \simeq 2 \text{ fm}^2 \Rightarrow (1.6 R_p)^2 \Rightarrow$  ultra-peripheral process

# Differential cross section in $p_t^2$



DPP 2 → 3



Geometric interpretation:  $\langle r^2 \rangle = 2b_1 \cdot (\hbar c)^2 \simeq 2 \text{ fm}^2 \Rightarrow (1.6 R_p)^2 \Rightarrow$  ultra-peripheral process

DPP explanation: low mass  $\pi^+ n$  state  $\rightarrow$  large slope, high masses  $\rightarrow$  less steep slope

# Summary

---

- Photoproduction cross section for exclusive  $\rho^0$  production associated with leading neutron is measured for the first time at HERA.
- Differential cross sections for the reaction  $\gamma p \rightarrow \rho^0 n \pi^+$  exhibit features typical for exclusive double peripheral process.
- The elastic photon-pion cross section,  $\sigma(\gamma \pi^+ \rightarrow \rho^0 \pi^+)$ , is extracted in the OPE approximation.