DIS 2014 - XXII. International Workshop on Deep-Inelastic Scattering and Related Subjects



Contribution ID: 85

Type: Oral presentation

Exclusive Photoproduction of Rho Meson with Leading Neutron at HERA

Tuesday 29 April 2014 11:40 (30 minutes)

The first measurement of exclusive photoproduction of rho meson associated with the leading neutron at HERA is presented. The data are taken with the H1 detector in the years 2006-2007 and correspond to an integrated luminosity of 1.16 pb-1. The rho mesons are reconstructed from decay pions in the central tracking chamber, while the leading neutrons, carrying a large fraction of the incoming proton momentum x_L>0.4, are detected in the Forward Neutron Calorimeter. The phase space of the measurement is restricted to by photon virtualities $Q^2 < 2 \text{ GeV}^2$, total energies of the gamma-p system 20 < W_gp < 100 GeV and polar angles of the leading neutrons, theta_n < 0.75 mrad. The cross section of the reaction gamma+p -> rho+n+Y, where Y is a small mass system escaping undetected in the proton direction, is measured as a function of the neutron energy, the invariant mass of the pi+pi- system and the transverse momentum and the pseudorapidity of the rho meson. The data are interpreted in terms of a double peripheral process, involving a pion exchange at the proton vertex followed by elastic photoproducton of a rho meson on a virtual pion.

Primary authors: DAUM, Karin (University of Wuppertal/DESY); SCHMITT, Stefan (Deutsches Elektronen-Synchrotron (DE))

Presenter: LEVONIAN, Sergey (Deutsches Elektronen-Synchrotron (DE))

Session Classification: WG2: Small-x, Diffraction and Vector Mesons

Track Classification: WG2: Small-x, Diffraction and Vector Mesons