

# Non-planar Feynman diagrams and Mellin-Barnes representations with AMBRE 3.0

I. Dubovyk<sup>1</sup>, J. Gluza<sup>2</sup>, Tord Riemann<sup>1</sup>

<sup>1</sup> Deutsches Elektronen-Synchrotron, DESY, Platanenallee 6, 15738 Zeuthen, Germany

<sup>2</sup> Institute of Physics, University of Silesia, Uniwersytecka 4, PL-40007 Katowice, Poland

E-mail: e.a.dubovyk@gmail.com, janusz.gluza@us.edu.pl, Tord.Riemann@desy.de

**Abstract.** We introduce the Mellin-Barnes representation of general Feynman integrals and discuss their evaluation. The Mathematica package **AMBRE** has been recently extended in order to cover consistently non-planar Feynman integrals with two loops. Prospects for the near future are outlined. This write-up is an introductory to new results which have also been presented elsewhere.

## 1. Introduction

The evaluation of Feynman integrals is a central numerical problem of perturbative quantum field theory and is not solved in generality beyond the one-loop case. One has to consider arbitrary  $L$ -loop integrals  $G(X)$  with loop momenta  $k_l$ , with  $E$  external legs with momenta  $p_e$ , and with  $N$  internal lines with masses  $m_i$  and propagators  $1/D_i$ ,

$$G(X) = \frac{1}{(i\pi^{d/2})^L} \int \frac{d^d k_1 \dots d^d k_L X(k_1, \dots, k_L)}{D_1^{n_1} \dots D_i^{n_i} \dots D_N^{n_N}}, \quad (1)$$

with  $d = 4 - 2\epsilon$ ,  $D_i = q_i^2 - m_i^2 = [\sum_{l=1}^L c_i^l k_l + \sum_{e=1}^E d_i^e p_e] - m_i^2$ , and some tensors  $X(k_1, \dots, k_L)$  in the loop momenta. For several reasons, one is interested in analytical solutions either:

- Analytical results are well suited for the exact cancellation of certain intermediate terms, arising either from the renormalization procedure, from regularization, or just from the organization of the calculation.
- Analytical results may stabilize the numerics.
- With analytical results, analytical continuations of presentations of Feynman integrals into regions of physical interest may be performed. Typically, it is easier to determine a Feynman integral in the Euclidean region, but often it is needed in the Minkowskian.

There are quite different approaches to tackle multi-loop Feynman integrals. Without aiming at completeness, we like to mention:

- Direct evaluation of a Feynman parameter representation (see e.g. the seminal articles [1, 2], and for later improvements [3, 4, 5] and references therein)
- Solving systems of differential or difference equations (see e.g. [6, 7, 8] and refs. therein)

- Expansion by regions (see e.g. [9] and refs. therein)
- Sector decomposition (see e.g. [10, 11, 12, 13, 14, 15, 16] and refs. therein)
- Mellin-Barnes (MB-) representations (see e.g. [17, 18, 19, 20, 21, 22, 23] and refs. therein)

For an overview of most recent developments we recommend [24]. We are concentrating here on the Mellin-Barnes approach. Quite some work has been devoted to an automation of it, see [25, 26, 27, 28, 29, 30, 31] and software at the webpage <http://projects.hepforge.org/mbtools>. Nevertheless, the formalism is much less developed than e.g. that with differential equations, where algorithms have been worked out for solving them in terms of certain classes of functions. An analogue idea would be here using Cauchy's theorem in order to derive from a Mellin-Barnes representation multiple sums of residues, and then to sum up these sums analytically [32]. How this might work with the software of the RISC (Linz) group [33, 34, 35, 36, 37, 38, 39, 40, 41] around the packages `SIGMA` [42, 43], `EvaluateMultiSums`, and `SumProduction` [44, 45, 46] is indicated in [47]. For not too involved classes of functions, one may try to apply the packages `SUMMER` (<http://www.nikhef.nl/~t68/>) or `XSUMMER` [48]. For automation, several problems have to be solved:

- Determine if the topology is planar or non-planar – Mathematica package `PlanarityTest` [31], see <http://prac.us.edu.pl/~gluza/ambre/planarity/>.
- Construct the appropriate Mellin-Barnes representations – Mathematica package `AMBRE` [26, 27, 28, 47], see <http://prac.us.edu.pl/~gluza/ambre/>. Here, also the Mathematica packages `MB` [25] and `MBresolve` [29] are integrated – for the analytic continuation of Mellin-Barnes integrals in  $\epsilon$ ; as well as `MBasymptotics` (2005, <http://projects.hepforge.org/mbtools/>) – for the parametric expansions of Mellin-Barnes integrals and `barnesroutines` by D. Kosower (2008, <http://projects.hepforge.org/mbtools/>) – for the automatic application of the first and second Barnes lemmas.
- Change the MB-integrals into multiple nested sums – Mathematica package `MBsums`, under development by M. Ochman (DESY) et al., unpubl., see [47].
- Finally, try to perform the multiple sums analytically, see [47, 49].
- Alternatively, a purely numerical evaluation of the multi-dimensional Mellin-Barnes integral may be envisaged. Here it is wishful to cover not only Euclidean cases, but also the Minkowskian kinematics; this has not been studied so far.

Of course, a solution of the general case is not to be expected. The limitations of the method have several reasons; we mention here:

- The number of loops;
- The number of different scales due to internal masses and the kinematics;
- The number of external legs;
- A planar or non-planar topology.

A Feynman parameter integral with  $N$  internal legs has, essentially, a dimensionality  $N - 1$ . For the corresponding MB-integral, the dimensionality will be different, and for complex problems it often will come out much higher.

## 2. Mellin-Barnes representations

For the Feynman integral (1) one may derive the following Feynman parameter integral:

$$G(X) = \frac{(-1)^{N_\nu} \Gamma(N_\nu - \frac{d}{2}L)}{\prod_{i=1}^N \Gamma(n_i)} \int \prod_{j=1}^N dx_j x_j^{n_j-1} \delta(1 - \sum_{i=1}^N x_i) \frac{U(x)^{N_\nu - d(L+1)/2}}{F(x)^{N_\nu - dL/2}} \quad (2)$$

The functions  $U$  and  $F$  are called graph or Symanzik polynomials.

The repeated application of the Mellin-Barnes representation [17]

$$\frac{1}{(A+B)^\lambda} = \frac{1}{\Gamma(\lambda)} \frac{1}{2\pi i} \int_{-i\infty}^{+i\infty} dz \Gamma(\lambda+z) \Gamma(-z) \frac{B^z}{A^{\lambda+z}} \quad (3)$$

to the Symanzik polynomials  $U$  and  $F$  allows to replace the sums of monomials in their definitions (here:  $A(x)$  and  $B(x)$ ) by products of these monomials and  $\Gamma$ -functions and, subsequently, to perform the  $x$ -integrations. The integration path separates poles of  $\Gamma[\lambda+z]$  and  $\Gamma[-z]$ . **AMBRE 1.1**, **1.2**, **2.0** perform so-called loop-by-loop integrations with  $U=1$  and are efficient enough for many applications (figure 1, left). For the massless, on-shell non-planar diagram of figure 1, right, which was first solved in [21], one would have to treat:

$$U(x) = +x[1]x[2] + x[1]x[4] + x[2]x[4] + x[1]x[5] + x[2]x[5] + x[2]x[6] \\ + x[4]x[6] + x[5]x[6] + x[1]x[7] + x[4]x[7] + x[5]x[7] + x[6]x[7], \quad (4)$$

$$F(x) = -s x[1]x[2]x[5] - s x[1]x[3]x[5] - s x[2]x[3]x[5] - u x[2]x[4]x[6] \\ - s x[3]x[5]x[6] - t x[1]x[4]x[7] - s x[3]x[5]x[7] - s x[3]x[6]x[7]. \quad (5)$$


Here it becomes operational to apply the Cheng-Wu theorem [50, 51] which states that (2) holds also with a modified delta function  $\delta(1 - \sum_{i \in \Omega} x_i)$  where  $\Omega$  is an arbitrary subset of the lines  $1, \dots, N$ , when the integration over the rest of the variables, i.e. for  $i \notin \Omega$ , is extended to the integration from 0 to  $\infty$ . With **AMBRE 3.0**, non-planar Feynman integrals with two loops may be efficiently represented by a direct approach with use of the Cheng-Wu theorem. Planar integrals are treated by the loop-by-loop ansatz with the earlier **AMBRE** versions. For the example of the *massless non-planar double box* mentioned, **AMBRE 3.0** derives without manual interaction the 4-dimensional Mellin-Barnes representation of [21]. It might look more promising to apply here instead the loop-by-loop approach, but experience shows that the dimensionality becomes, without further interventions, higher. At the other hand, for the *massive non-planar double box*, the loop-by-loop approach gives the MB-presentation derived first in [22]. We mention here shortly that 3-loop integrals may be treated in a hybrid way by a loop-by-loop approach, subsequently using **AMBRE 1.2** and **AMBRE 3.0** [47]; see the example files `MB_hybrid_3loopNP_massless.nb`, `MB_hybrid_3loopNP_massless.m`, `out_MB_hybrid_3loopNP_massless` at the webpage <http://prac.us.edu.pl/~gluza/ambre/>.

Finally we would like to mention that the mathematica package **AMBRE 3.0** is released for public use at the webpage <http://prac.us.edu.pl/~gluza/ambre/>. There is no source file made publicly available so far. We consider this to be appropriate; for closer information we refer to the webpages <http://fh.desy.de/projekte/gfitter01/Gfitter01.htm> (March 2013) and <http://zfitter-gfitter.desy.de/> (April 2014). For a collection of experts' views on proper distribution of scientific software in basic research, we would like to refer to [52, 53], a Summary of a round table discussion at ACAT 2014.

### 3. Summary

We sketched essential features of the Mellin-Barnes approach to Feynman integrals and its implementation in **AMBRE**. In the new version **AMBRE 3.0** the Cheng-Wu theorem is implemented and the replacement of the Symanzik polynomials by MB-integrals is performed globally. The alternative loop-by-loop approach is implemented in **AMBRE 2.0** or older versions. Which of the approaches is more appropriate for a given problem has to be investigated.

A treatment of non-planar three-loop integrals deserves the combined application of **AMBRE 1.2** and of **AMBRE 3.0** in a hybrid approach. This will be improved in the nearest future. The *analytical* summation of series of MB-residues is under study. Concerning the alternative to analytical summation, namely a *numerical* evaluation of the MB-integrals, we are restricted with **AMBRE** to the Euclidean kinematics so far. The Minkowskian numerics is on our to-do list.



**Figure 1.** Left: The planar double box with loop momenta as used for the loop-by-loop presentation. Right: The non-planar double box.

### Acknowledgements

Work is supported in part by Sonderforschungsbereich/Transregio SFB/TRR 9 “Computergestützte Theoretische Teilchenphysik” of Deutsche Forschungsgemeinschaft (DFG), by the Polish National Center of Science (NCN) under the Grant Agreement number DEC-2013/11/B/ST2/04023, and by the European Initial Training Network LHCPHENOnet PITN-GA-2010-264564.

### References

- [1] G. 't Hooft, M. Veltman, Scalar One Loop Integrals, Nucl. Phys. B153 (1979) 365–401. doi:10.1016/0550-3213(79)90605-9.
- [2] G. Passarino, M. Veltman, One Loop Corrections for  $e^+e^-$  Annihilation Into  $\mu^+\mu^-$  in the Weinberg Model, Nucl. Phys. B160 (1979) 151. doi:10.1016/0550-3213(79)90234-7.
- [3] J. Fleischer, T. Riemann, Complete algebraic reduction of one-loop tensor Feynman integrals, Phys. Rev. D83 (2011) 073004. arXiv:1009.4436, doi:10.1103/PhysRevD.83.073004.
- [4] J. Fleischer, T. Riemann, Calculating contracted tensor Feynman integrals, Phys. Lett. B701 (2011) 646–653. arXiv:1104.4067, doi:10.1016/j.physletb.2011.06.033.
- [5] J. Fleischer, T. Riemann, A solution for tensor reduction of one-loop N-point functions with  $N \geq 6$ , Phys. Lett. B707 (2012) 375–380. arXiv:1111.5821, doi:10.1016/j.physletb.2011.12.060.
- [6] A. Kotikov, Differential equations method: New technique for massive Feynman diagrams calculation, Phys. Lett. B254 (1991) 158–164. doi:10.1016/0370-2693(91)90413-K.
- [7] E. Remiddi, Differential equations for Feynman graph amplitudes, Nuovo Cim. A110 (1997) 1435–1452. arXiv:hep-th/9711188.
- [8] J. M. Henn, Multiloop integrals in dimensional regularization made simple, Phys. Rev. Lett. 110 (25) (2013) 251601. arXiv:1304.1806, doi:10.1103/PhysRevLett.110.251601.
- [9] M. Beneke, V. A. Smirnov, Asymptotic expansion of Feynman integrals near threshold, Nucl. Phys. B522 (1998) 321–344. arXiv:hep-ph/9711391, doi:10.1016/S0550-3213(98)00138-2.
- [10] K. Hepp, Proof of the Bogolyubov-Parasiuk theorem on renormalization, Commun. Math. Phys. 2 (1966) 301–326, <http://www.projecteuclid.org/euclid.cmp/1103815087>.
- [11] M. Roth, A. Denner, High-energy approximation of one loop Feynman integrals, Nucl. Phys. B479 (1996) 495–514. arXiv:hep-ph/9605420, doi:10.1016/0550-3213(96)00435-X.
- [12] T. Binoth, G. Heinrich, Numerical evaluation of multiloop integrals by sector decomposition, Nucl. Phys. B680 (2004) 375–388. arXiv:hep-ph/0305234, doi:10.1016/j.nuclphysb.2003.12.023.
- [13] A. Denner, S. Pozzorini, An Algorithm for the high-energy expansion of multi-loop diagrams to next-to-leading logarithmic accuracy, Nucl. Phys. B717 (2005) 48–85. arXiv:hep-ph/0408068, doi:10.1016/j.nuclphysb.2005.03.036.
- [14] G. Heinrich, Sector Decomposition, Int. J. Mod. Phys. A23 (2008) 1457–1486. arXiv:0803.4177, doi:10.1142/S0217751X08040263.
- [15] A. Smirnov, M. Tentyukov, Feynman Integral Evaluation by a Sector decomposition Approach (FIESTA), Comput. Phys. Commun. 180 (2009) 735–746. arXiv:0807.4129, doi:10.1016/j.cpc.2008.11.006.
- [16] A. Smirnov, V. Smirnov, Hepp and Speer Sectors within Modern Strategies of Sector Decomposition, JHEP 0905 (2009) 004. arXiv:0812.4700, doi:10.1088/1126-6708/2009/05/004.
- [17] E. W. Barnes, A new development of the theory of the hypergeometric functions, Proc. Lond. Math. Soc. (2) 6 (1908) 141–177. doi:10.1112/plms/s2-6.1.141.

- [18] N. Usyukina, On a Representation for Three Point Function, *Teor.Mat.Fiz.* 22 (1975) 300–306. doi:10.1007/BF01037795.
- [19] E. Boos, A. I. Davydychev, A Method of evaluating massive Feynman integrals, *Theor. Math. Phys.* 89 (1991) 1052–1063. doi:10.1007/BF01016805.
- [20] V. A. Smirnov, Analytical result for dimensionally regularized massless on shell double box, *Phys. Lett. B* 460 (1999) 397–404. arXiv:hep-ph/9905323, doi:10.1016/S0370-2693(99)00777-7.
- [21] J. Tausk, Nonplanar massless two loop Feynman diagrams with four on-shell legs, *Phys. Lett. B* 469 (1999) 225–234. arXiv:hep-ph/9909506, doi:10.1016/S0370-2693(99)01277-0.
- [22] G. Heinrich, V. Smirnov, Analytical evaluation of dimensionally regularized massive on-shell double boxes, *Phys. Lett. B* 598 (2004) 55–66. arXiv:hep-ph/0406053.
- [23] M. Czakon, J. Gluza, T. Riemann, The Planar four-point master integrals for massive two-loop Bhabha scattering, *Nucl. Phys. B* 751 (2006) 1–17. arXiv:hep-ph/0604101, doi:10.1016/j.nuclphysb.2006.05.033.
- [24] J. Blümlein, P. Marquard, Tord Riemann (eds.), *Loops and Legs in Quantum Field Theory*, PoS(LL2014) Proceedings of the 12th DESY Workshop on Elementary Particle Theory, April 27 – May, 2, 2014, Weimar, Germany. <http://pos.sissa.it/cgi-bin/reader/conf.cgi?confid=211>.
- [25] M. Czakon, Automatized analytic continuation of Mellin-Barnes integrals, *Comput. Phys. Commun.* 175 (2006) 559–571. arXiv:hep-ph/0511200, doi:10.1016/j.cpc.2006.07.002.
- [26] J. Gluza, K. Kajda, T. Riemann, AMBRE - a Mathematica package for the construction of Mellin-Barnes representations for Feynman integrals, *Comput. Phys. Commun.* 177 (2007) 879–893. arXiv:0704.2423, doi:10.1016/j.cpc.2007.07.001.
- [27] J. Gluza, K. Kajda, T. Riemann, V. Yundin, News on Ambre and CSectors, *Nucl. Phys. Proc. Suppl.* 205-206 (2010) 147–151. arXiv:1006.4728, doi:10.1016/j.nuclphysbps.2010.08.034.
- [28] J. Gluza, K. Kajda, T. Riemann, V. Yundin, Numerical Evaluation of Tensor Feynman Integrals in Euclidean Kinematics, *Eur. Phys. J. C* 71 (2011) 1516. arXiv:1010.1667, doi:10.1140/epjc/s10052-010-1516-y.
- [29] A. Smirnov, V. Smirnov, On the Resolution of Singularities of Multiple Mellin-Barnes Integrals, *Eur. Phys. J. C* 62 (2009) 445–449. arXiv:0901.0386, doi:10.1140/epjc/s10052-009-1039-6.
- [30] B. Jantzen, New proofs for the two Barnes lemmas and an additional lemma, *J. Math. Phys.* 54 (2013) 012304. arXiv:1211.2637, doi:10.1063/1.4775770.
- [31] K. Bielas, I. Dubovyk, J. Gluza, T. Riemann, Some Remarks on Non-planar Feynman Diagrams, *Acta Phys. Polon. B* 44 (11) (2013) 2249–2255. arXiv:1312.5603, doi:10.5506/APhysPolB.44.2249.
- [32] J. Gluza and T. Riemann, *Simple Feynman diagrams and simple sums*, Talk held at RISC - DESY *Workshop on Advanced Summation Techniques and their Applications in Quantum Field Theory* on the occasion of the 5th year jubilee of the RISC-DESY cooperation, May 7-8, 2012, RISC Institute, Castle of Hagenberg, Linz, Austria. Conference link: <http://www.risc.jku.at/conferences/RISCDESY12/>, talk link: <http://www-zeuthen.desy.de/~riemann/Talks/riemann-Linz-2012-05.pdf>.
- [33] M. Karr, Summation in finite terms, *J. ACM* 28 (1981) 305–350. doi:10.1145/322248.322255.
- [34] C. Schneider, Symbolic summation in difference fields, Tech. Rep. 01-17, RISC-Linz, J. Kepler University, PhD Thesis, [http://www.risc.jku.at/publications/download/risc\\_3017/SymbSumTHESIS.pdf](http://www.risc.jku.at/publications/download/risc_3017/SymbSumTHESIS.pdf) (November 2001).
- [35] C. Schneider, Solving parameterized linear difference equations in terms of indefinite nested sums and products, *J. Differ. Equations Appl.* 11 (2005) 799–821. doi:10.1080/10236190500138262.
- [36] C. Schneider, Simplifying Sums in  $\Pi\Sigma$ -Extensions, *J. Algebra Appl.* 6 (2007) 415–441. doi:10.1142/S0219498807002302.
- [37] C. Schneider, A refined difference field theory for symbolic summation, *J. Symbolic Comput.* 43 (2008) 611–644. arXiv:0808.2543, doi:10.1016/j.jsc.2008.01.001.
- [38] C. Schneider, Structural Theorems for Symbolic Summation, *Appl. Algebra Engrg. Comm. Comput.* 21 (2010) 1–32. doi:10.1007/s00200-009-0115-3.
- [39] C. Schneider, A Symbolic Summation Approach to Find Optimal Nested Sum Representations, in: A. Carey, D. Ellwood, S. Paycha, S. Rosenberg (Eds.), *Motives, Quantum Field Theory, and Pseudodifferential Operators*, Vol. 12 of Clay Mathematics Proceedings, Amer. Math. Soc, 2010, pp. 285–308, [http://www.risc.jku.at/publications/download/risc\\_3806/DepthSequ.pdf](http://www.risc.jku.at/publications/download/risc_3806/DepthSequ.pdf).
- [40] C. Schneider, Parameterized Telescoping Proves Algebraic Independence of Sums, *Ann. Comb.* 14 (2010) 533–552. arXiv:0808.2596.
- [41] C. Schneider, Fast Algorithms for Refined Parameterized Telescoping in Difference Fields, in: M. W. J. Guitierrez, J. Schicho (Ed.), *Computer Algebra and Polynomials*, Lecture Notes in Computer Science (LNCS), to appear, Springer, 2014. arXiv:1307.7887.
- [42] C. Schneider, Symbolic summation assists combinatorics, *Sém. Lothar. Combin.* 56 (2007) 1–36, Article B56b,

- <http://www.emis.de/journals/SLC/wpapers/s56.html>.
- [43] C. Schneider, Simplifying multiple sums in difference fields, in: C. Schneider, J. Blümlein (Eds.), *Computer Algebra in Quantum Field Theory: Integration, Summation and Special Functions*, Texts and Monographs in Symbolic Computation, Springer, 2013, pp. 325–360. [arXiv:1304.4134](#).
  - [44] J. Ablinger, J. Blümlein, S. Klein, C. Schneider, Modern Summation Methods and the Computation of 2- and 3-loop Feynman Diagrams, *Nucl. Phys. Proc. Suppl.* 205-206 (2010) 110–115. [arXiv:1006.4797](#), [doi:10.1016/j.nuclphysbps.2010.08.028](#).
  - [45] J. Blümlein, A. Hasselhuhn, C. Schneider, Evaluation of Multi-Sums for Large Scale Problems, *PoS RADCOR2011* (2011) 032. [arXiv:1202.4303](#).
  - [46] C. Schneider, Modern Summation Methods for Loop Integrals in Quantum Field Theory: The Packages Sigma, EvaluateMultiSums and SumProduction, *J.Phys.Conf.Ser.* 523 (2014) 012037. [arXiv:1310.0160](#), [doi:10.1088/1742-6596/523/1/012037](#).
  - [47] J. Blümlein, I. Dubovyk, J. Gluza, M. Ochman, C. G. Raab, T. Riemann, C. Schneider, Non-planar Feynman integrals, Mellin-Barnes representations, multiple sums, *PoS(LL2014)052*. [http://pos.sissa.it/archive/conferences/211/052/LL2014\\_052.pdf](http://pos.sissa.it/archive/conferences/211/052/LL2014_052.pdf). [arXiv:1407.7832](#).
  - [48] S. Moch, P. Uwer, XSummer: Transcendental functions and symbolic summation in form, *Comput. Phys. Commun.* 174 (2006) 759–770. [arXiv:math-ph/0508008](#), [doi:10.1016/j.cpc.2005.12.014](#).
  - [49] J. Ablinger, J. Blümlein, C. G. Raab, C. Schneider, Nested (inverse) binomial sums and new iterated integrals for massive Feynman diagrams, *PoS(LL2014)*. [arXiv:1407.4721](#).
  - [50] N. Nakanishi, Parametric integral formulas and analytic properties in perturbation theory, *Prog. Theor. Phys. Supplement* 18 (1961) 1.  
URL <http://ptps.oxfordjournals.org/content/18/1.full.pdf>
  - [51] N. Nakanishi, *Graph Theory and Feynman Integrals*, Gordon and Breach, 1971.
  - [52] F. Carminati, D. Perret-Gallix, T. Riemann, Summary of the ACAT [2013] Round Table Discussion: Open-source, knowledge sharing and scientific collaboration, *J. Phys. Conf. Ser.* 523 (2014) 012066. [doi:10.1088/1742-6596/523/1/012066](#).
  - [53] F. Carminati, D. Perret-Gallix, T. Riemann, Summary of the ACAT [2013] Round Table Discussion: Open-source, knowledge sharing and scientific collaboration. Red report DESY 14-079. [arXiv:1407.0540](#).