

STRUCTURE OF MATTER

Discoveries and Mysteries

Rolf Landua
CERN

(Presenter and marginal contributions : Giacinto de Cataldo INFN Bari, Italy)

PREFACE

**This is a lecture about 100 years of particle physics.
It covers about 100 years of ideas, theories and experiments.**

More than 50 Nobel prize winners on particle physics
This is a broad overview about the main discoveries.

**In the early 1900s, most physicists believed that physics was complete,
described by classical mechanics, thermodynamics, and the Maxwell
theory.**




Lord Kelvin

“There is nothing new to be discovered in physics now. All that remains is more and more precise measurement.” (Lord Kelvin, 1900)

DARK CLOUDS:

- 1) Blackbody radiation - Quantum Physics
- 2) Michelson-Morley experiment - Special Relativity



Higgs boson

MATTER IS MADE OF PARTICLES




J.J. Thomson

1897: ELECTRON - the first 'discrete' building block of matter



A. Einstein

1905: ATOMS ARE REAL - Explanation of Brownian Motion (Perrin)




$$\langle x^2 \rangle = \frac{2kTt}{\alpha} = \frac{kTt}{3\pi\eta a}$$

ENERGY COMES IN QUANTA



M. Planck

1900: ELECTROMAGNETIC RADIATION IS EMITTED IN QUANTA



$$\epsilon = h \nu$$


$$I(n) \sim n^2 \frac{hn}{e^{\frac{hn}{kT}} - 1}$$



P. von Lenard

1902: PHOTOELECTRIC EFFECT

“The electron energy does not show the slightest dependence on the light intensity”



A. Einstein

1905: LIGHT IS EMITTED AND ABSORBED IN QUANTA

$$E_{\max} = h\nu - W$$


“My only revolutionary contribution to physics”

SPECIAL RELATIVITY



A. Einstein

1905: SPEED OF LIGHT IS ALWAYS CONSTANT




$$c^2 t^2 = v^2 t^2 + w^2$$

$$t = \frac{w/c}{\sqrt{1 - \frac{v^2}{c^2}}} = g \times t$$

- 1) Time dilation, space contraction
- 2) Modification of Newton's laws, relativistic mass increase.

$$E = mc^2$$

THE BEGINNING OF ATOMIC PHYSICS



Rutherford

1909: NUCLEI: very small + heavy within (almost) empty atom



Hydrogen

1913: BOHR MODEL- (empirical) explanation of discrete spectral lines

(using Planck's constant h) to quantize angular momentum




L. de Broglie

1923: DE BROGLIE

Particles are (also) waves
(particle-wave duality)


$$\lambda = \frac{h}{p}$$



Which role of the classical mechanics in the atomic world ?

(gdc)

In classical mechanics the measurement of observables (E, p, x) describes the state of the system .



QUANTUM MECHANICS




Heisenberg

1923: UNCERTAINTY RELATION

$$\Delta x \Delta p \geq \frac{\hbar}{2}$$

$$\Delta E \Delta t \geq \hbar$$



Schrödinger

1926: SCHRÖDINGER EQUATION


$$H\psi(\mathbf{r}, t) = (T + V)\psi(\mathbf{r}, t) = \left[-\frac{\hbar^2}{2m} \nabla^2 + V(\mathbf{r}) \right] \psi(\mathbf{r}, t) = i\hbar \frac{\partial \psi}{\partial t}(\mathbf{r}, t)$$

(electrons in atoms form 'standing waves')

Interpretation (Born, 1927):

ψ = probability amplitude

$|\psi|^2$ = probability



RELATIVISTIC QUANTUM MECHANICS



Paul A.M. Dirac
(1928)

$$E^2 = p^2 + m^2 \rightarrow$$
$$E = \pm(\alpha \cdot p) + \beta m$$

$$(i\gamma^\mu \partial_\mu - m)\psi = 0$$

$$\Psi = \begin{pmatrix} e^- \uparrow \\ e^- \downarrow \\ e^+ \uparrow \\ e^+ \downarrow \end{pmatrix}$$

Spin

Antimatter

CONSEQUENCES:

ELECTRON **SPIN** EXPLAINED
ANTIPARTICLES MUST EXIST !
ELECTRONS OBEY 'PAULI PRINCIPLE' (1940) - **FERMIONS**

ANTIPARTICLES




Anderson

1932: POSITRON DISCOVERY



EVERY PARTICLE HAS AN ANTIPARTICLE

$$E=mc^2$$



WHEN ENERGY CONVERTS TO MASS,
PARTICLES AND ANTIPARTICLES ARE PRODUCED

QUANTUM FIELD THEORY (1927 - 1948)



S.I. Tomonaga

It was known that the electromagnetic field consists of photons



J. Schwinger

How could the interaction between electrons and photons be correctly described, respecting quantum mechanics and special relativity?




F. Dyson

Many people worked on this problem ...

EMPTY SPACE HAD BECOME COMPLICATED !

Quantum physics says that 'oscillators' (e.g. field quanta) cannot be at absolute rest (uncertainty relation)


The lowest energy states of e.g. electromagnetic fields can produce (virtual) electron-positron pairs: VACUUM FLUCTUATIONS



Quantum Electrodynamics (QED)





R.P. Feynman



QED: Charged particles interact by exchanging photons

- 1) **Massless virtual photons are continuously emitted by electric charges**
- 2) The **$1/r^2$ law** comes from the probability to hit another particle at distance r
(directly connected with the 3 dimensions of space)




$1/r^2$ law

QED became a model for other interactions

RENORMALIZATION : HOW TO DEAL WITH INFINITIES

The 'naked' electron + vacuum fluctuations = measured electron


(“infinite” - “infinite” = “finite”)



vacuum fluctuations modify its charge and mass
('Debye shielding')


Vacuum fluctuations have observable effects

... and Quantum Electrodynamics allows to calculate them precisely




Lamb Shift

(shift of atomic energy levels)



Electron (anomalous) magnetic moment

$$\frac{1}{2}(g - 2) = \frac{1}{2} \frac{\alpha}{\pi} - 0.32848 \left(\frac{\alpha}{\pi} \right)^2 + (1.183 \pm 0.011) \left(\frac{\alpha}{\pi} \right)^3 .$$



Casimir effect

(force on two uncharged metal plates)


SPOOKY !

THE BEGINNING OF NUCLEAR PHYSICS



M. Curie

1895-1900: RADIOACTIVITY - strange radiation phenomena



E. Rutherford


1903: Alpha-, Beta-, Gamma-Radiation known
(different penetration; Alpha = He-Nucleus)



J. Chadwick

1911: Nucleus positive, small - surrounded by electrons

1932: DISCOVERY OF THE NEUTRON



Fields

'Strong' interaction

The "Strong Interaction" - Nuclear forces

What keeps the protons and neutrons together in the nucleus?


- 1) This force must be stronger than the electromagnetic repulsion
- 2) It must be of short range ($\sim 1-2$ fm) to explain the size of nuclei

Yukawa's idea:


a massive particle ("pion") is exchanged between two nucleons



Yukawa (1934)



Electromagnetic vs Nuclear



$$V(r) = -e^2 \frac{1}{r}$$

Coulomb law




emits massive pions

$$\Delta E \Delta t \geq \hbar \quad (\Delta E \sim m)$$

$$r = c \Delta t = \frac{\hbar c}{m} \sim \frac{200 \text{ MeV fm}}{m}$$

$$V(r) = -g^2 \frac{e^{-mr}}{r}$$


Yukawa potential ~ Modified "Coulomb" law




Fields

'Strong' interaction

Metaphors for 'particle exchange'



Allowed by uncertainty relation: $1.4 \text{ fm} \sim 140 \text{ MeV}$



1913: Cosmic Rays were discovered


Physicists went on mountain tops for experiments!

1937: Negative particle with $M \sim 200 m_e$

Very long range in matter !? Not Yukawa's "pion" !

Muon = 'heavy electron'

Who ordered that ?




1948: Discovery of the 'pion'

PARTICLE SPECTRUM


1948

In 1948, the particle spectrum started to look ugly:





Rolf Wideroe, 1928




Linear accelerator

Accelerate particles between electrode gaps
Tune RF frequency to match particle motion

Accelerators

"Man-made cosmic rays"

Ernest Lawrence, 1931



Cyclotron

Use magnetic field to bend particles into circular orbit
Particles pass through same accelerating gap many times and reach higher energies

1931: 80 keV


1932: 1000 keV

1939: 19 MeV*

1946: 195 MeV ("synchrocyclotron")

* first limitations by relativistic mass increase

Accelerators (2)



Synchrotron

Similar to cyclotron, but change magnetic field to keep particles on the same orbit
(also overcomes relativistic mass increase)

1947: US constructs two 'synchrotrons'

Brookhaven (1952) - 3 GeV

Berkeley (1954) - 6.2 GeV ('antiproton')

1954: Europe competes with US

CERN (1959) - 24 GeV

Brookhaven (1960) - 30 GeV

Detectors

Geiger counters
Cloud chambers
Emulsions
Bubble chambers

Cerenkov counters
Photomultipliers
Spark chambers

After 1967:

Wire chambers
Drift chambers
Calorimeters

PARTICLE SPECTRUM

1950- 1968

Particle
zoo

With new accelerators and detectors,
the "particle zoo" grew to more than ~ 200 'elementary particles'

π^+ π^- π^0

Pions

K^+ K^- K^0

η'

Kaons

Eta-Prime

η

Eta

ϕ

ρ^+ ρ^- ρ^0

Rho

Phi

Mesons

Δ^{++} , Δ^+ , Δ^0 , Δ^-

Delta

Λ^0

Σ^+ , Σ^0 , Σ^-

Lambda (strange!)

Sigma (strange!)

Ξ^0 , Ξ^-

Sigma(very strange!)

BARYONS

What was the underlying structure ?




SU(3) - Classification scheme based on 'quarks'



Fig. 6.35 Murray Gell-Mann (b.1929).


Gell-Mann, 1963
(G. Zweig, 1963, CERN)

- 1) 3 types of "quarks" : up, down, strange
- 2) Carry electric charges: $+2/3$, $-1/3$, $-1/3$
- 3) Appear in combinations:
 Meson = quark+antiquark
 Baryon = quark(1) + quark(2) + quark(3)




PARTICLE SPECTRUM


Some mesons (quark+antiquark):




π^0




π^+



π^-




K^+



PARTICLE SPECTRUM

1967 Discovery of quarks

Electron-Proton scattering (1956)




1956 Hofstadter: measured finite proton radius



Stanford Linear Accelerator Centre

1967 Friedmann, Kendall, Taylor (SLAC):


'hard scattering' of electron on three 'point-like particles'



Measured cross-sections perfectly compatible with presence of 2 up- and 1 down-quark in proton

The concept of “Colour” charge

PROBLEM: three fermions are not allowed to be in identical states (Pauli exclusion principle)



Since the three up-quarks must have parallel spin - there are in a symmetric state

The three quarks must be different in one quantum number: “colour”

(Bardeen, Fritsch, Gell-Mann)

PARTICLE SPECTRUM


Quantum Chromo Dynamics

this has nothing to do with our visible colours, just an analogy

Theory constructed in analogy to QED

QCD: 3 different charges (“colour charge”) [red, green, blue]*

‘Strong force’ between quarks is transmitted by (8) gluons (massless)



Dogma of QCD: Only colour-neutral bound states are allowed, explains:


MESONS = Quark-Antiquark

BARYONS = 3-Quark states

GLUONS CARRY COLOUR CHARGE : SELF-INTERACTION !



At low energies, QCD long-range forces increase with distance. These forces keep the quarks as 'prisoners' .



At high energies, for small distances, the force decreases: asymptotic freedom of quarks (phenomenon: the quark-gluon plasma)