

# The Structure and Signals of Neutron Stars, from Birth to Death



Contribution ID: 36

Type: **not specified**

## Energy transformations in the birth of neutron stars

*Monday 24 March 2014 18:00 (15 minutes)*

Millisecond pulsars, radio pulsars, and magnetars are distinct classes of neutron stars that currently exhibit different periods, magnetic fields and space velocities. In my talk, I speculate with the idea that these current properties of neutron stars could have had similar values during the birth of these stars. I develop a relatively simple model based on the assumption that neutron stars experienced three abrupt physical changes at the end of their birth, which could have been originated in birth magneto-rotational instabilities: an increase in period from the initial value to the current value, implying a change of rotational energy; an exponential decay of its magnetic field from the initial value to the current surface value, implying a change of radiative energy; and finally, an increase of space velocity from the initial value to the current value, implying a change of kinetic energy. These birth energy changes are assumed to be connected by an energy conversion in which a radiation loss and an increase of kinetic energy occur at the expense of a rotational energy loss. According to this model, if the assumed energy conversion occurs in times of order  $10^{-4}$  s then neutron stars are born with magnetic fields in the range of  $10^{15}$ - $10^{16}$  G and initial periods in range 1-20 ms. This means that neutron stars are born with magnetic fields typical of magnetars and periods typical of millisecond pulsars.

**Primary author:** Dr HERAS, Ricardo (University College London)

**Presenter:** Dr HERAS, Ricardo (University College London)

**Session Classification:** Evening session - Parallel B