

PONT Avignon – April 15, 2014



# Dark Matter in the Black Hills

## The LUX Experiment at SURF

First Results from October 2013  
and a few more things that have happened since then



[www.luxdarkmatter.org](http://www.luxdarkmatter.org)

# The LUX Collaboration



## Brown

Richard Gaijskell	PI, Professor
Simon Fiorucci	Research Associate
Monica Panglinan	Postdoc
Jeremy Chapman	Graduate Student
David Malling	Graduate Student
James Verbus	Graduate Student
Samuel Chung Chan	Graduate Student
Dongqing Huang	Graduate Student



## Case Western

Thomas Shutt	PI, Professor
Dan Akerib	PI, Professor
Karen Gibson	Postdoc
Tomasz Biesiadzinski	Postdoc
Wing H To	Postdoc
Adam Bradley	Graduate Student
Patrick Phelps	Graduate Student
Chang Lee	Graduate Student
Kati Pech	Graduate Student



## Imperial College London

Henrique Araujo	PI, Reader
Tim Sumner	Professor
Alastair Currie	Postdoc
Adam Bailey	Graduate Student



## Lawrence Berkeley + UC Berkeley

Bob Jacobsen	PI, Professor
Murdock Gilchriese	Senior Scientist
Kevin Lesko	Senior Scientist
Carlos Hernandez Faham	Postdoc
Victor Gehman	Scientist
Mia Ihm	Graduate Student



## Lawrence Livermore

Adam Bernstein	PI, Leader of Adv. Detectors Group
Dennis Carr	Mechanical Technician
Kareem Kazkaz	Staff Physicist
Peter Sorensen	Staff Physicist
John Bower	Engineer



## LIP Coimbra

Isabel Lopes	PI, Professor
Jose Pinto da Cunha	Assistant Professor
Vladimir Solovov	Senior Researcher
Luiz de Viveiros	Postdoc
Alexander Lindote	Postdoc
Francisco Neves	Postdoc
Claudio Silva	Postdoc



## SD School of Mines

Xinhua Bai	PI, Professor
Tyler Liebsch	Graduate Student
Doug Tiedt	Graduate Student



## SDSTA

David Taylor	Project Engineer
Mark Hanhardt	Support Scientist



## Texas A&M

James White †	PI, Professor
Robert Webb	PI, Professor
Rachel Mannino	Graduate Student
Clement Sofka	Graduate Student



## UC Davis

Mani Tripathi	PI, Professor
Bob Svoboda	Professor
Richard Lander	Professor
Britt Holbrook	Senior Engineer
John Thomson	Senior Machinist
Ray Gerhard	Electronics Engineer
Aaron Manalaysay	Postdoc
Matthew Szydagis	Postdoc
Richard Ott	Postdoc
Jeremy Mock	Graduate Student
James Morad	Graduate Student
Nick Walsh	Graduate Student
Michael Woods	Graduate Student
Sergey Uvarov	Graduate Student
Brian Lenardo	Graduate Student



## UC Santa Barbara

Harry Nelson	PI, Professor
Mike Witherell	Professor
Dean White	Engineer
Susanne Kyrre	Engineer
Carmen Carmona	Postdoc
Curt Nehrhorn	Graduate Student
Scott Haselschwardt	Graduate Student



## University College London

Chamkaur Ghag	PI, Lecturer
Lea Reichhart	Postdoc



Collaboration Meeting, Sanford Lab, April 2013



## University of Edinburgh

Alex Murphy	PI, Reader
Paolo Beltrame	Research Fellow
James Dobson	Postdoc



## University of Maryland

Carter Hall	PI, Professor
Attila Dobi	Graduate Student
Richard Knoche	Graduate Student
Jon Balajthy	Graduate Student



## University of Rochester

Frank Wolfs	PI, Professor
Wojtek Skutski	Senior Scientist
Eryk Druszkiewicz	Graduate Student
Mongkol Moongweluwan	Graduate Student



## University of South Dakota

Dongming Mei	PI, Professor
Chao Zhang	Postdoc
Angela Chiller	Graduate Student
Chris Chiller	Graduate Student
Dana Byram	*Now at SDSTA



## Yale


Daniel McKinsey	PI, Professor
Peter Parker	Professor
Sidney Cahn	Lecturer/Research Scientist
Ethan Bernard	Postdoc
Markus Horn	Postdoc
Blair Edwards	Postdoc
Scott Hertel	Postdoc
Kevin O'Sullivan	Postdoc
Nicole Larsen	Graduate Student
Evan Pease	Graduate Student
Brian Tennyson	Graduate Student
Ariana Hackenburg	Graduate Student
Elizabeth Boulton	Graduate Student

# LUX Design – Dual Phase Xenon TPC



Instrument paper:

NIM. A 704,  
111-126 (2013)  
[arXiv:1211.3788](https://arxiv.org/abs/1211.3788)

- Can measure single electrons and photons
- Charge yield reduced for nuclear recoils
- Excellent 3D imaging
  - Reject multiple scatters
  - Eliminate edge events to take advantage of Xe self shielding




The LUX Detector Grid Configuration



# Typical Event in LUX


~ 1.5 keV gamma






# The LUX Program

LUX0.1 - CWRU



2007 – 2009

LUX - Surface



2010 – 2011


LUX - Underground



2012+




# The Sanford Laboratory at Homestake



[www.sanfordlab.org](http://www.sanfordlab.org)

# Sanford Lab – Davis Laboratory






# Davis Campus – September 2012





# LUX Run 3: Some Statistics




- Detector cool-down January 2013, Xe condensed mid-February 2013
- 95% Data taking efficiency during WIMP search period (minus storms)
- Waited until after WS data before precision  $\text{CH}_3\text{T}$  calibration


# Light and Charge Yields

- Modeled Using Noble Element Simulation Technique (NEST) - [arxiv:1106.1613](https://arxiv.org/abs/1106.1613)
- NEST based on canon of existing experimental data.
- **Artificial cutoff** in light and charge yields assumed **below 3 keV<sub>nr</sub>**, to be conservative.
- Includes predicted electric field quenching of light signal, to 77-82% of the zero field light yield

S1 Light yield  $L_{eff}$





S2 yield  $Q_y$








# ER and NR Calibration Data




CH<sub>3</sub>T injection system



<sup>83m</sup>Kr injections



# LUX WIMP Search, 85.3 live-days, 118 kg



## The Economist

*"Absence of evidence, or evidence of absence?"*


## New York Times

*"Dark Matter Experiment Has Detected Nothing, Researchers Say Proudly"*




# LUX Run 3 – Spin Independent Sensitivity

- Deviation of  $1\sigma$  in detection efficiency shifts the limit by 5%



# Projected LUX 300 day WIMP Search Run


- **Wrap up post-Run3 items until Summer 2014**
  - Increased calibration stats
  - Measure NR response
  - Looking at improving E-fields
- **More results from Run 3 data**
  - Exploring low-energy regime
  - “Exotic” searches
- **We intend to run LUX for a new run of 300 days in 2014/15**
  - Extending sensitivity by another factor ~5
  - Discovery still possible
- **LZ 20x increase in target mass**
  - If approved, plans to be deployed in Davis Lab in 2016+







# DD Neutron Generator – Result (preliminary!)

- 105 live hours of beam time accumulated
- Remove many instrumental systematics
- Analysis still ongoing
  - Improve MC stats and accuracy
  - Push energy threshold down
  - Refine understanding of systematics
- Dedicated publication will be forthcoming
- **Direct implications for low mass sensitivity!**



Compare to slide 26



---

Result paper published in PRL 112.091303

<http://arxiv.org/abs/1310.8214>

Instrument paper NIM A 704 111-126 (2013)

<http://arxiv.org/abs/1211.3788>

Backgrounds paper submitted to Astropart. Phys.

<http://arxiv.org/abs/1403.1299>

Several LUX theses now available on luxdarkmatter.org



**Run 4 to start first half of 2014. Expect more exciting results in the future!**

**Thank you!**







Because this was not long enough already...

**Additional Slides**

# LUX Design – Water Tank



- Water Tank:  $d = 8\text{ m}$ ,  $h = 6\text{ m}$ 
  - 300 tonnes, 3.5 m thickness on the sides
  - Inverted steel pyramid (20 tonnes) under tank to increase shielding top/bottom
- Cherenkov muon veto
- Ultra-low background facility
  - Gamma event rate reduction:  $\sim 10^{-9}$
  - High-E neutrons ( $>10\text{ MeV}$ ):  $\sim 10^{-3}$



# LUX Krypton Removal System

arXiv:1103.2714


- $^{85}\text{Kr}$  - beta decay – intrinsic background in liquid Xe
  - Research grade Xenon:  $\sim 100$  ppb Kr  $\Rightarrow 10^4 - 10^5$  reduction needed
- August 2012 - January 2013: Kr removal at dedicated facility
  - Chromatographic separation system
- Kr concentration reduced from 130 ppb to  $3.5 \pm 1$  ppt, (factor of 35000)
  - 1 ppt is achievable, working on sub-ppt (useful for next-generation detectors)






# Run 3 Parameters Overview

- **Xenon Purity: electron drift length 87 – 135 cm during Run 3**
  - Circulation at 250 kg / day
  - Monitored weekly using  $^{83m}\text{Kr}$  data
- **Light collection efficiency: 14%**
  - Incl. geometry and PMT QE
  - $^{83m}\text{Kr}$  data provides 3D corrections
- **Drift field: 181 V/cm**
  - Drift speed  $1.51 \pm 0.01$  mm /  $\mu\text{s}$
  - ER discrimination 99.6%
- **Electron extraction efficiency: 65%**
- **Fiducial mass:  $118.3 \pm 6.5$  kg**
  - Defined by edges  $\alpha$  background
  - Measured with homogeneous ER calibration data...  $^{83m}\text{Kr}$ , again!



# LUX Calibrations – $^{83\text{m}}\text{Kr}$

- $^{83}\text{Rb}$  produces  $^{83\text{m}}\text{Kr}$  when it decays; this krypton gas can then be flushed into the LUX gas system to calibrate the detector as a function of position.
- Provides reliable, efficient, homogeneous calibration of both S1 and S2 signals, which then decays away in a few hours, restoring low-background operation.





- Bonus: tomography of Xe flow

$^{83\text{m}}\text{Kr}$  source ( $^{83}\text{Rb}$  infused into zeolite, within xenon gas plumbing)



# LUX Calibrations – CH<sub>3</sub>T


- LUX uses tritiated methane, doped into the detector, to accurately calibrate the efficiency of background rejection.
- This beta source (endpoint energy 18 keV) allows electron recoil S2/S1 band calibration with unprecedented accuracy
- The tritiated methane is then fully removed by circulating the xenon through the getter
- This was tested first with natural methane injection, and monitored with our in-line xenon sampling system
- Dedicated paper coming





# ER Discrimination

Average discrimination from 2-30 S1 photoelectrons measured to be 99.6% (with 50% nuclear recoil acceptance)




Black circles show leakage from counting events from the dataset

Red circles show projections of Gaussian fits below the nuclear recoil band mean

# LUX Run 3 – Background Levels

- Full gamma spectrum, excluding region  $\pm 2$  cm from top/bottom grids




# LUX Run 3 – Background Levels

Background Component	Source	$10^{-3} \times \text{evts/keVee/kg/day}$
Gamma-rays	Internal Components including PMTS (80%), Cryostat, Teflon	$1.8 \pm 0.2_{\text{stat}} \pm 0.3_{\text{sys}}$
$^{127}\text{Xe}$ (36.4 day half-life)	Cosmogenic 0.87 $\rightarrow$ 0.28 during run	$0.5 \pm 0.02_{\text{stat}} \pm 0.1_{\text{sys}}$
$^{214}\text{Pb}$	$^{222}\text{Rn}$	0.11-0.22(90% CL)
$^{85}\text{Kr}$	Reduced from 130 ppb to $3.5 \pm 1$ ppt	$0.17 \pm 0.1_{\text{sys}}$
Predicted	Total	$2.6 \pm 0.2_{\text{stat}} \pm 0.4_{\text{sys}}$
Observed	Total	$3.1 \pm 0.2_{\text{stat}}$


- Neutron background negligible (expect 0.06 evt)
- Dedicated publication is now available:

[arXiv:1403.1299](https://arxiv.org/abs/1403.1299)

ER < 5 keVee in 118 kg



Last 44 days






# LUX Run 3 – Data Selection

Cut	Explanation	Events Remaining
All Triggers	S2 Trigger >99% for $S2_{\text{raw}} > 200$ phe	83,673,413
Detector Stability	Cut periods of excursion for Xe Gas Pressure, Xe Liquid Level, Grid Voltages	82,918,901
Single Scatter Events	Identification of S1 and S2. Single Scatter cut.	6,585,686
S1 energy	Accept 2-30 phe (energy $\sim 0.9$ -5.3 keVee, $\sim 3$ -18 keVnr)	26,824
S2 energy	Accept 200-3300 phe (>8 extracted electrons) Removes single electron / small S2 edge events	20,989
S2 Single Electron Quiet Cut	Cut if >100 phe outside S1+S2 identified +/-0.5 ms around trigger (0.8% drop in livetime)	19,796
Drift Time Cut away from grids	Cutting away from cathode and gate regions, $60 < \text{drift time} < 324$ us	8731
Fiducial Volume (R,Z)t cut	Radius < 18 cm, $38 < \text{drift time} < 305$ us, 118 kg fiducial	160

- Simple, obvious cuts
- No “tuning” beyond selecting a threshold, higher energy cutoff, fiducial volume
- PLR analysis not so sensitive to how many events are “in the box”

# LUX Run 3 – NR Detection Efficiency



True Recoil Energy equivalence based on LUX 2013 Neutron Calibration/NEST Model

# Profile Likelihood Ratio (PLR) Analysis

$$\mathcal{L}_{WS} = \frac{e^{-N_s - N_{Compt} - N_{Xe-127} - N_{Rn222}}}{\mathcal{N}!} \prod_{i=1}^{\mathcal{N}} \left( N_s P_s(\mathbf{x}; \sigma, \theta_s) + N_{Compt} P_{ER}(\mathbf{x}; \theta_{Compt}) + N_{Xe-127} P_{ER}(\mathbf{x}; \theta_{Xe-127}) + N_{Rn} P_{ER}(\mathbf{x}; \theta_{Rn}) \right)$$


**Observables:**  $\mathbf{x} = (S1, \log_{10}(S2/S1), r, z)$

**Parameter of interest:**  $N_s$


**Nuisance parameters:**  $N_{Compt}, N_{Xe-127}, N_{Rn, Kr-85}$

**SIGNAL MODEL:** simulated 2D PDFs including resolution/efficiencies; uniform in  $(r^2, z)$


LUX Simulation



LUX Simulation



LUX Simulation






# Precision NR Response Calibration for LXe



- Use a mono-energetic source of neutrons at 2.45 MeV
- Collimate them with a 3 m tube of air through water
- Identify double scatters along beam line inside LUX. Angle gives deposited energy.
  - → Absolute calibration of ionization response
- Apply ionization scale found above to single scatters
  - → Absolute calibration of scintillation response

$$E_r = E_n \frac{4m_n m_{Xe}}{(m_n + m_{Xe})^2} \frac{1 - \cos \theta}{2}$$




# Further Exploring very-low Energies

- Simply applying new  $L_{\text{eff}}$  and  $Q_y$  to existing analysis already provides significant improvement below 10 GeV
  - As much as possible we want to explore all options for a “new” Run3 result together instead of piecemeal
- With DD neutron data in hand, we can be more confident about exploring data below current analysis thresholds
  - Results from simulated data shown at APS:
- Also looking at S2-only analysis, similar to what was done in Xenon10
  - Challenges: fiducialization, single-e background
  - Progress on algorithms also presented at APS






# Grids HV Conditioning

- Fields achieved during Run 3 were sufficient, but we can do better
  - Drift field – 181 V/cm. Discrimination ~99.6%. Higher field may get even better (?)
  - Gas extraction field – 6 kV/cm. Efficiency 65%, can definitely get better.
  - Both limited by electroluminescence
- Conditioning = continuously apply high voltage, watch current on grids, glow patterns, electron emission...
- In this mode since January 2014. Campaign reaching its end.
  - Some hints of progress
  - Will know for sure when cool down and condense again
  - A few more details presented at APS meeting



# Contribution from Intrinsic Sources: $^{127}\text{Xe}$

- Isotope of interest for WIMP search =  $^{127}\text{Xe}$ 
  - EC decay with gammas 203 or 375 keV, possibility to escape the active volume.
  - X-ray / Auger emission corresponding to  $^{127}\text{I}$  levels: 33.2 keV<sub>ee</sub> (K), 5.3 (L), 1.1 (M), 0.19 (N)
  - Depth-dependent background profile; data follows prediction
  - Contribution modeled as a nuisance in the PLR analysis
  - Accounts for **0.5 mDRU<sub>ee</sub> (avg) in WIMP ROI** over Run 3
  - It will have disappeared for Run 4





# Uranium and Thorium Chains

## HIGH-A DECAY CHAINS

