



Higgs Physics at CMS

Albert De Roeck
CERN, Geneva, Switzerland
Antwerp University Belgium
UC-Davis California USA




7th October 2013
Tehran, Iran




The 2nd IPM meeting on
LHC PHYSICS
7 - 12 October 2013
Tehran, Iran



Sponsors:





Outline

- Short introduction
- Standard Model Higgs channel studies overview
- Studies of Higgs properties

Note: not all slides will be discussed in detail




OR: The year after!!
CERN/Melbourne
4th of July




The Origin of Particle Masses

- At 'low' energy the Weak force is much weaker than the Electromagnetic force: **Electroweak Symmetry Breaking: ESB**
- The W and Z bosons are very massive (~ 100 proton masses) while the photon is massless.
- The proposed mechanism^(*) in 1964 gives mass to W and Z bosons and predicts the existence of a new elementary 'Higgs' particle,. Extend the mechanism to give mass to the Fermions via Yukawa couplings.



(*) Higgs, Brout Englert, Kibble, Hagen and Guralnik, and...




The Higgs (H) particle is the quantum of the new postulated field and has been searched for since decades at other particle colliders such as **LEP** and the **Tevatron**, and now at the **large hadron collider @ CERN**

ESB Heroics

The year is 1964


Electroweak Symmetry Breaking



François Englert



Robert Brout



Peter Higgs



Gerald Guralnik



Carl Hagen



Tom Kibble

BROKEN SYMMETRY AND THE MASS OF GAUGE VECTOR

F. Englert and R. Brout

Faculté des Sciences, Université Libre de Bruxelles, Bruxelles
(Received 26 June 1964)

BROKEN SYMMETRIES, PART I

Nobel Institute of Mathematics

VOLUME 13, NUMBER 16 PHYS

BROKEN SYMMETRY



CONSERVATION LAWS AND MASSLESS PARTICLES*

S. Guralnik,† C. R. Hagen,‡ and T. W. B. Kibble
Department of Physics, Imperial College, London, England
(Received 12 October 1964)

LDS

Tomorrow the Nobel Prize Committee will announce the 2013 winners... Will they reward any of these gentlemen for their very important work done almost 50 years ago?? We will know on December 8th...

The Hunt for the Higgs



Where do the masses of elementary particles come from?

The key question (pre-2012):
Does the Higgs particle exist?
If so, where is the Higgs?

Massless particles move at the speed of light \rightarrow no atom formation!!

We do not know the mass of the Higgs Boson


$$\mathcal{L}_{\text{Higgs}} = (\partial_\mu \phi)^\dagger (\partial^\mu \phi) - V(\phi)$$
$$V(\phi) = \mu^2 \phi^\dagger \phi + \lambda (\phi^\dagger \phi)^2$$



Scalar field with at least one scalar particle


It could be anywhere from 114 to ~ 700 GeV

Higgs Production & Decay



- ## Processes
- Gluon fusion
 - Vector Boson Fusion
 - W/Z associated prod.
 - Top associated prod.

Decays



Numbers taken from the LHC Higgs Cross Section WG

- YR1: Inclusive cross sections
arXiv:1101.0593
- YR2: Differential cross sections
arXiv:1201.3084
- YR3: Properties
arXiv:1307.1347

Higgs Hunting in CMS

Processes/decays studied:

Results released
 In progress

	untagged	VBF	VH	ttH
H-> gamgam				
H-> ZZ				
H-> WW				
H-> bb				
H-> tau tau				
H-> Zgamma				
H-> mumu				
H-> invisible				

Main decay channel characteristics:

+ more exotic channels

Channel	m_H range (GeV/ c^2)	Data used 7+8 TeV (fb^{-1})	m_H resolution
H -> $\gamma\gamma$	110-150	5.1+19.6	1-2%
H -> tautau	110-145	4.9+19.6	15%
H -> bb	110-135	5.0+19.0	10%
H -> WW -> lnu lnu	110-600	4.9+19.5	20%
H -> ZZ -> 4l	110-1000	5.1+19.6	1-2%

Higgs Hunters

Higgs Hunting Basics

Needle-in-the-hay-stack problem


- need high energy:

$$E = mc^2$$

- need lots of data

non-deterministic and very rare

order 1 in 10^9












* for us finding the Higgs it was
48 years = 1,513,728,000 sec


Higgs Boson Searches (simulation)

Low $M_H < 140 \text{ GeV}/c^2$


Medium $130 < M_H < 500 \text{ GeV}/c^2$ High $M_H > \sim 500 \text{ GeV}/c^2$

simulation





100 meter underground




Luminosity = # events/cross section/time

LHC operation is now stopped for 2 years, and the machine is being prepared for running at 13-14 TeV from 2015 onwards

Note: the LHC is a Higgs Factory: 1 Million Higgses already produced
15 Higgses/minute with present luminosity

Higgs Decay into Bosons

The Decay $H \rightarrow \gamma\gamma$





Analysis

- Two high momentum photons
- Low mass Higgs is narrow
- Two photon resolution is excellent
- Looking for a narrow peak
- Large irreducible background from direct two photons
- Smaller fake photon background

Key analysis features

- Energy resolution (calibration)
- Fake photon rejection
- Optimize use of kinematics

A Collision with two Photons



A Higgs or
a 'background'
process without
a Higgs?

CMS: The Decay $H \rightarrow \gamma\gamma$



- Two inclusive analyses:

PRIMARY

- MVA:** photons selected with an MVA. Variable in the MVA: photon kinematics, photon ID MVA score (shower shape, isolation), di-photon mass resolution. 4 MVA categories with different S/B

CROSS-CHECK


- Cut-based:** photons selected with cuts. 4 categories based on: γ in Barrel/Endcap, (un)converted γ . Each category has different mass resolution and S/B




Event categories weighted by $S/(S+B)$ for visualization

CMS: The Decay $H \rightarrow \gamma\gamma$

MVA mass-factorized





Cut-based



	MVA analysis (at $m_H=125 \text{ GeV}$)	cut-based analysis (at $m_H=124.5 \text{ GeV}$)
7 TeV	$1.69^{+0.65}_{-0.59}$	$2.27^{+0.80}_{-0.74}$
8 TeV	$0.55^{+0.29}_{-0.27}$	$0.93^{+0.34}_{-0.32}$
7 + 8 TeV	$0.78^{+0.28}_{-0.26}$	$1.11^{+0.32}_{-0.30}$



The Decay $H \rightarrow \gamma\gamma$

MVA mass-factorized



7+8 TeV: σ/σ_{SM} for a mass of 125.0 GeV = $0.78^{+0.28}_{-0.26}$

The Decay $H \rightarrow Z\gamma$




- Z decays into 2 charged leptons. The BR ($H \rightarrow Z\gamma$) is comparable to BR($H \rightarrow \gamma\gamma$), but BR ($Z \rightarrow ll$) reduces sensitivity (factor 15)
- Search for a narrow $ll\gamma$ peak on top of a falling background, as for $H \rightarrow \gamma\gamma$
- No significant excess seen over the entire search region

In certain models this channel could be largely enhanced

The Decay $H \rightarrow ZZ \rightarrow 4l$

Analysis

- 4 isolated high p_T leptons consistent with Z decays from same vertex
- Use a di-jet tagged and untagged category, and kinematics
- Clear mass peak
- Little background, main comes from non-resonant ZZ production, also Zbb and top ($2l2\nu2b$), fakes



CMS-PAS-HIG-13-002

Analysis procedure rather stable since ICHEP2012

Searches for the Higgs Particle



A Higgs particle will decay immediately, eg in two heavy quarks or two heavy (W,Z) bosons

Example: Higgs(?) decays into ZZ and each Z boson decays into $\mu\mu$



So we look for 4 muons in the detector

But two Z bosons can also be produced in LHC collisions, without involving a Higgs!

We cannot say for one event by event (we can reconstruct the total mass with the 4 muons)



CMS: The Decay $H \rightarrow ZZ \rightarrow 4l$



Use event kinematics (MELA)

Significance Expected: 7.1σ
 Observed: 6.7σ


σ/σ_{SM} at 125.7 GeV = 0.92 ± 0.28

Significance is well over 6 standard deviations in this channel

The Decay $H \rightarrow ZZ \rightarrow 4l$




Matrix **E**lement **L**ikelihood **A**nalysis:
uses kinematic inputs to build a kinematic discriminant (K_D) for signal to background discrimination using $\{m_1, m_2, \theta_1, \theta_2, \theta^*, \Phi, \Phi_1\}$

$$\text{MELA} = \left[1 + \frac{\mathcal{P}_{\text{bkg}}(m_1, m_2, \theta_1, \theta_2, \Phi, \theta^*, \Phi_1 | m_{4l})}{\mathcal{P}_{\text{sig}}(m_1, m_2, \theta_1, \theta_2, \Phi, \theta^*, \Phi_1 | m_{4l})} \right]^{-1}$$




SIGNAL

BACKGROUND




The Decay $H \rightarrow ZZ \rightarrow 4l$

Mass Measurements



$$M_H = 125.8 \pm 0.5(\text{stat.}) \pm 0.2(\text{syst.}) \text{ GeV.}$$

Coupling scale factors to vector bosons and fermions




$$\mu_V (qqH, ZH, WH) = 1.0^{+2.4}_{-2.3}$$

$$\mu_F (gg \rightarrow H, t\bar{t}H) = 0.9^{+0.5}_{-0.4}$$

The Decay $H \rightarrow WW \rightarrow 2l 2\nu$

CMS-PAS-HIG-13-003

μ : 32 GeV




Analysis

- Two opposite charged leptons (leptons only e, μ)
- Two neutrinos == missing transverse energy (MET)
- No Higgs mass peak
- Counting & 2D shape analyses
- Enhance sensitivity by subdividing into + (0,1,2) jets categories

Analysis challenges



- Understand backgrounds WW, W+jets, top, Drell-Yan
- Determined from control regions



Scalar decay and V-A structure of W decay lead to a small opening angle between leptons

The Decay $H \rightarrow WW \rightarrow 2l 2\nu$


Events with 0 jets and different flavour leptons (7+8 TeV Data)




A significant excess is observed...

The Decay $H \rightarrow WW \rightarrow 2l 2\nu$

Standard Analysis





Using $m_H = 125 \text{ GeV}$ as a "background"



- Exclusion at 95% in the mass range 128-600 GeV
- Large excess in the low mass region
- When including $M_H=125 \text{ GeV}$ as part of the background, no significant excess is seen over the entire mass range

The Decay $H \rightarrow WW \rightarrow 2l 2\nu$




A 4.0σ (5.1σ) observed (expected) significance at $m_H \sim 125$ GeV

$\sigma/\sigma_{\text{SM}}$ signal strength: 0.76 ± 0.21


VBF: $H \rightarrow WW \rightarrow 2l 2\nu$

2-jet category in CMS
(VBF dominated)



Mild broad excess in the low mass

CMS-PAS-HIG-13-022




analysis	limits		significance		best μ
	expected / observed	expected / observed	expected / observed	expected / observed	observed
VBF, $e\mu$ and $ee/\mu\mu$ final states combined					
7 + 8 TeV (cut-based)	1.1 / 0.9	2.0 / 0.0	-0.35 ^{+0.43} _{-0.45}		
7 + 8 TeV (fit to m_{ee})	1.1 / 1.7	2.1 / 1.3	0.62 ^{+0.58} _{-0.47}		

Associated Production $VH H \rightarrow WW$


$WH \rightarrow WWW \rightarrow 3l 3\nu$

$VH \rightarrow VWW \rightarrow 2l 2\nu + V \rightarrow jj$

CMS-PAS-HIG-13-009



CMS-PAS-HIG-12-017



- Three high p_T leptons with moderate missing transverse momentum

- WW analysis cuts plus two central jets


Limited Standard Model Higgs sensitivity ($\sim 3.5\text{-}4 \cdot \text{SM}$ at 125 GeV)

Higgs Decay into Fermions


The Decay $H \rightarrow \tau\tau$

Topologies studied


CMS-PAS-HIG-13-004



Inclusive
0-jets




Boosted
1-jet



VBF
2-jets



Analysis

- Tau decays to $e, \mu, \tau_{\text{had}}$ used to reconstruct a tau
- Reconstruct corrected $\tau\tau$ invariant mass
- Use many categories to increase the sensitivity





CMS Experiment at LHC, CERN
Data recorded: Sun Nov 25 00:15:46 2012 CEST
Run/Event: 207898 / 97057018

VBF candidate event for
 $H \rightarrow \tau\tau \rightarrow \mu\tau_h$



The Decay $H \rightarrow \tau\tau$





...plus all other tau decay modes: $e\tau_h$, $e\mu$, $\mu\mu$, $\tau_h\tau_h$

Combine the sensitive categories of all channels with a S/B weight

Associated Production $VH \rightarrow V\tau\tau$




- Study topologies of 3 and 4 lepton final states
- Use tau decay channels into electrons muons and hadronic final states



Upper limits of 2.9 to 4.6 times the predicted Standard Model value for $\sigma \cdot \text{BR}$ at 95% CL.

The Decay $H \rightarrow \tau\tau$

Results include also the VH channels



Significance:
 2.93σ for $m_H = 120$ GeV
 2.85σ for $m_H = 125$ GeV

Signal strength
 $\mu = 1.1 \pm 0.4$

Mass: all $\tau\tau$ channels combined
 $m_H = 120^{+9}_{-7}$ (stat+syst) GeV



Excess building up in the region 120-130 GeV

The Decay $H \rightarrow bb$

CMS-PAS-HIG-13-012



Analysis

- By far largest number of Higgs decays
- But lots of QCD background (jets)
- Trigger based on leptons and missing E_T
- b-jets identified through displaced tracks
- Go to high p_T where Higgs is enhanced
- Main background W/Z+jets and top






The Decay $H \rightarrow b\bar{b}$

$M_{b\bar{b}}$ for all categories and 7+8 TeV



The Decay $H \rightarrow bb$





For 125 GeV:

- Significance = 2.1σ (2.1σ expected)
- Signal strength $\mu = 1.0 \pm 0.5$

Mild excess observed in data.


VBF Process with $H \rightarrow bb$



bb event + ≥ 2 non-b jets at large $\Delta\eta$




CMS-PAS-HIG-13-011

At 125 GeV the upper limit
on $\sigma \cdot \text{BR} = 3.6 \cdot \text{SM}$ (3.0 exp.)



New: Higgs Decay into two muons

We do not expect to see any signal yet in this channel if it is really a Higgs particle that we have found





No significant excess observed

- Limits at 125 GeV 7.4 (5.1) observed (expected)
- Excess at 125 GeV is 1.1σ
- Excess at high mass -148 GeV- is 2.3σ (0.8 with LLE)

Higgs Decay into two Electrons

This is even less expected to be observed: about 4000 times lower cross section for $H \rightarrow ee$ than $H \rightarrow \mu\mu$!!






No significant excess observed

- Limits No signal at 125 GeV, as expected
- Sensitivity in ee and $\mu\mu$ channel very similar


Higgs Associated with Top Production

Direct measurement of top-Higgs coupling
 Channels studied $H \rightarrow \text{CC}$ and $H \rightarrow \text{bb} + \dots$



Select top events with additional Higgs

- A recent result in $ttH(\text{CC})$ channel
 - $\mu < 5.4$ (5.3 exp.) @ 95% CL, $m_H = 125$ GeV
- New $ttH(\text{bb} + \dots)$ results:
 - $\mu < 5.2$ (4.1 exp.) @ 95% CL, $m_H = 125$ GeV





Higgs Associated with Top Production

- **New analysis** exploring various top and Higgs decays resulting in like-sign di-lepton, tri-lepton, and quadri-lepton final states ($H \rightarrow WW$ etc...)
- An excess ($\sim 2.5\sigma$) seen in like-sign dimuons has been extensively scrutinized and shown **to have all the features of a statistical fluctuation**
- Overall consistency with the SM: 3%


NEW

CMS-HIG-13-020



All Channel ttH Combination

- All channels combined
- Impressive expected sensitivity $\mu < 2!$
Excess is driven by the di-muon excess in the multi-lepton analysis



Sensitivity is between 1 and 2 times the SM Expectation!

Channel Combination & Higgs Properties

April 2013 Combination



Since fall 2012 we have been especially concentrating on measurements of properties of the new particle

Summary of the Five Main Channels

For a mass of $m_H = 125.7$ GeV

CMS-PAS-HIG-13-005

Decay	Expected	Observed	
<i>ZZ</i>	7.1 σ	6.7 σ	
<i>$\gamma\gamma$</i>	3.9 σ	3.2 σ	
<i>WW</i>	5.3 σ	3.9 σ	
<i>bb</i>	2.2 σ	2.1 σ	} 3.4 σ combined!
<i>$\tau\tau$</i>	2.6 σ	2.8 σ	



bb: includes VH and VBF

WW: includes ggF, VH, VBF

Mass of the New Particle


$H \rightarrow ZZ \rightarrow 4l$: $m_H = 125.8 \pm 0.5$ (stat.) ± 0.2 (syst.) GeV

$H \rightarrow \gamma\gamma$: $m_H = 125.4 \pm 0.5$ (stat.) ± 0.6 (syst.) GeV




$m_H = 125.7 \pm 0.3^{(stat)} \pm 0.3^{(syst)}$ GeV
 $= 125.7 \pm 0.4$ GeV

Consistency with SM Hypothesis



p-value = 0.65 w.r.t. $\mu=1$



p-value = 0.52 w.r.t. $\mu=1$


Combined signal strength: $\mu = 0.80 \pm 0.14$

Here and further: bb results based on 12 fb^{-1} at 8 TeV and 5 fb^{-1} at 7 TeV

Consistency with SM Hypothesis

2-dimensional view: test production modes in the various decay modes

Vector Boson
Couplings




Fermion
Couplings


Signal Strength for Different Modes

Likelihood scans versus the different μ values, using all decay modes


Gluon Gluon Fusion



Vector Boson Fusion



Associated production




Data in good agreement with the expectation

Approximately a 2σ significance for the VBF channel

Couplings to Fermions and Bosons



Couplings scaled by κ_X :

Hff : κ_f HVV : κ_V
 HWW : κ_W $\lambda_{WZ} = \kappa_W / \kappa_Z$
 HZZ : κ_Z In SM, $\kappa_X = 1$





For $m_H = 125.7$ GeV
 $\Gamma(H \rightarrow \gamma\gamma) \sim |\alpha \kappa_V + \beta \kappa_f|^2$
 $\alpha/\beta = -0.2, \Gamma_{BSM} = 0$

Contributions from all decay channels



Results within 1σ of the Standard Model Prediction

Summary of the Couplings Test





For the fermions, the values of the fitted **yukawa couplings** are shown, while for vector bosons the square-root of the coupling for the **hVV vertex** divided by twice the vacuum expectation value of the Higgs boson field. _

Custodial Symmetry Test

Modify the SM Higgs boson couplings to the W and Z bosons introducing two scaling factors κ_W and κ_Z and perform combinations to assess if

$$\lambda_{WZ} = \kappa_W / \kappa_Z = 1$$

for $m_H = 125.7$ GeV




95% CL interval for λ_{WZ} : [0.62, 1.19]

Summary of the Couplings Test


Summary of the fits for deviations in the couplings

for a LHC XS WG benchmark model
parametrisation (arXiv:1209.0040)



for a generic five parameter model
(no eff. loop couplings)

$$\Gamma_{BSM} = 0$$



The best fit values of the most interesting parameters are shown, with the corresponding 68% and 95% CL intervals, and the overall p -value p_{SM} of the SM Higgs hypothesis is given.


Higgs Properties from $H \rightarrow \gamma\gamma$

CMS-PAS-HIG-13-016

Upper limit on the Higgs width


- Dominated by experimental resolution
- Breit-Wigner + Gaussian fit
- Observed (exp) upper limit = **6.9 (5.9) GeV 95% CL**

Use interference? arXiv:1305.3854 & more





Additional Higgs-like states:

- Take SM 125 GeV as part of the background
- Search for additional Higgses
- Largest excess: **136.5 GeV with 2.9σ** ($<2\sigma$ after LEE)



Search for near mass degenerate states




- Two signals with relative strength \times mass difference Δm
- Perform a 2D scan
- No signal at 95% CL for **$\Delta m > 4$ GeV**



Spin/Parity Hypothesis Tests

Spin/parity hypothesis tests: $H \rightarrow ZZ \rightarrow 4l$ channel

Kinematic discriminant built to describe the kinematics of production and decay of different J^P state of a "Higgs"



More J^P hypotheses have been tested in a similar way \rightarrow

J^P	CL_s
0^-	0.16%
0_h^+	8.1%
$2_{m\bar{g}g}^+$	1.5%
$2_{mq\bar{q}}^+$	<0.1%
1^-	<0.1%
1^+	<0.1%

Summary

- The mid-2012 discovery has been confirmed with more added collisions. Moving on to measuring properties.
- Rare processes now studied: $H \rightarrow Z\gamma$, ttH , $(H \rightarrow \mu\mu)$...
- The spin/parity is compatible with a 0^+ state and not with (simple) 0^- or spin 2 states
- The mass value by CMS is 125.7 ± 0.4 GeV
- Signs of decays into fermion decay channels. The significance of the combined $\tau+b$ channels is $\sim 3.4\sigma$
- The couplings to bosons and fermions are consistent with SM predictions, but these are tested so far up to $\sim 20\text{-}30\%$ precision only; Surprises still possible!!
- Hunt for rare decays & processes is going on...
- ATLAS is having similar results (see backup slides)

Tomorrow ???



Fingers crossed....

Search for Other Higgses

- High mass search for SM-like Higgses
- Invisible Higgs
- MSSM Neutral Higgs ($\mu\mu$, bb , $\tau\tau$)
- MSSM Charged Higgs
- Double Charged Higgs
- Light pseudoscalar a_1 production
- Fermiophobic/SM4 studies
- ...

See talk by A. Nikitenko







Backup: Comparison with ATLAS

Summer 2012: Results

Higgs \rightarrow 2 photons!!

Higgs \rightarrow 2Z \rightarrow 4 leptons!!

Higgs \rightarrow 2W \rightarrow 2l2v!!





Summer 2012: Results

Both experiments see an excess ~ 125 GeV in the $\gamma\gamma$, ZZ and WW channel

→ Adding up all the channels gives the following combination



Shown is the compatibility with a 'background only hypothesis'

5 fb⁻¹/2011 and 5 fb⁻¹/2012





CMS and ATLAS observe a **new boson** with a significance of **about 5 sigma** (1 chance in 3 million to be wrong!!!)

Update with the Full 2012 Data Sample




Increased data sample with a factor of ~ 3

The particle is clearly still with us, now with a significance of $>10\sigma$!!




We now enter the phase of measuring the properties of the new particle


The Birth of a Particle




“History” of the data accumulation during the last two years



ATLAS




CMS




The Mass of the Particle

Determine the mass from ZZ and 2-photon channels which show a peak!



$$\hat{m}_H = 125.5 \pm 0.2(\text{stat})_{-0.6}^{+0.5}(\text{syst}) \text{ GeV}$$





$$\hat{m}_H = 125.7 \pm 0.3(\text{stat}) \pm 0.3(\text{syst}) \text{ GeV}$$

ATLAS and CMS observe the same particle!! 😊

Signal Strength

- Signal strength μ is the observed over Standard Model expected cross section
- For $\mu=1$ the production rate is compatible with Standard Model expectation




ATLAS a bit above and CMS a bit below $\mu=1$...

The Spin of the New Particle


A Higgs particle should be a spin 0^+ state

- Study angular correlations in the decays of the particle; build likelihoods and test spin- and parity hypotheses
- Use the ZZ, 2-photon and WW final states


=> Particle is consistent with a 0^+ state!!



ATLAS




CMS




Couplings to the New Particle


- Use information of all production and decay channels
- κ_f and κ_V are scale factors w.r.t. the Standard Model values for **fermions** and **vector bosons**



ATLAS



CMS



⇒ Couplings compatible Standard Model values, but large uncertainties
 ...Future data will decide...