

# **Study of soft gluon resummation as a function of particle mass & center-of-mass energy in high-energy p-p collisions**

**The 2nd IPM Meeting on LHC Physics  
October 7-12, 2013 Tehran, IRAN**

**David d'Enterria**


CERN

**Soodeh Zarepour**

Shiraz University


# Typical proton-proton collision

1. Incoming hadron
  - ➔ Parton distribution function (fitted to data)
2. Initial-state radiation (ISR)
  - ➔ DGLAP parton evolution
3. Hard scattering:
  - ➔ Matrix element calculation at LO, NLO, ... level
4. Final state radiation
  - ➔ DGLAP parton evolution
5. Underlying event
  - ➔ Multiple softer parton interactions
6. Hadronization
  - ➔ Parton fragmentation functions (fitted to data)




# Hard scattering example: top pair production (NLO)

The (differential) cross section can be expressed as:  $\frac{d\sigma}{dQ_T^2} \sim \alpha_s + \alpha_s^2 + \alpha_s^3 + \dots$



# Soft gluon radiation

Soft gluons are very easy to radiate and this affects the  $p_T$  distribution of particle X:



$$\frac{1}{(q+g)^2 - m_q^2} = \frac{1}{2E_g E_q (1 - \beta_q \cos \theta_{qg})}$$

soft divergence if  $E_g \rightarrow 0$   
collinear if  $\theta_{qg} \rightarrow 0$  (only if  $m_q = 0$ )

In some kinematics regions (e.g. at low  $Q$ ) terms of the form:  $\alpha_s^n \ln(Q^2 / Q_T^2) = \alpha_s^n L$  are large.

Thus, the following terms are effectively of the same order::

$$\alpha_s(1+L) \sim \alpha_s^2(L^2+L^3) \sim \alpha_s^3(L^4+L^5)$$

We need to **re-order the terms of the perturbative expansion**:

fixed-order  
calculation

$$\frac{d\sigma}{dQ_T^2} = Q_T^{-2} \left\{ \alpha_s(1+L) + \alpha_s^2(L^2+L^3) + \alpha_s^3(L^4+L^5) \right. \\ \left. + \alpha_s^2(1+L) + \alpha_s^3(L^2+L^3) \right. \\ \left. + \alpha_s^3(1+L) + \dots \right\}$$

Log  
resummation

$$\frac{d\sigma}{dQ_T^2} = Q_T^{-2} \left\{ \alpha_s(1+L) + \alpha_s^2(L^2+L^3) + \alpha_s^3(L^4+L^5) \right. \\ \left. + \alpha_s^2(1+L) + \alpha_s^3(L^2+L^3) \right. \\ \left. + \alpha_s^3(1+L) + \dots \right\}$$

# NLO + soft gluon emission example: $t\bar{t}$

## 1) Parton-shower (LO):

Soft & collinear gluon emission via Monte Carlo

**Good:** low- $p_T$  distribution

**Bad:** it misses total x-section and high- $p_T$


## 2) NLO:

**Good:** total x-section & high- $p_T$

**Bad:** Artificially large distribution at low- $p_T$  ( $t\bar{t}$  often produced at  $\sim$ rest)

## 3) NLO+parton-shower:

**Good:** everywhere...




# Goals of study and theoretical tools

**Goal:** Study low part of  $p_T$  distributions for various heavy-particles: DY, W, Z, H and  $t\bar{t}$ .

**How:** Studying the evolution of the peak of  $d\sigma/dp_T$  (whose position is dominated by soft-gluon resummation effects) as a function of the mass of the particle and  $\sqrt{s}$ .

**Tools:**









**PO**Positive **W**eight **H**ardest **E**mission **G**enerator  
 Avoids double-counting of parton-shower contributions based on  $p_T$  of parton

(\*) Tune: models semi-hard and non-perturbative part of collision: MPI, UE, hadronization, ...



$d\sigma / dp_T$  for Z, PDF set: CT10nlo &  $(\mu_f, \mu_R) = (m_X, m_X)$


Local fit (e.g. polynomial 3) for finding peak of  $p_T$  distribution :









- ✓ The  $p_T$  -distribution gets harder with increasing  $\sqrt{s}$ .
- ✓ The peak position increases slowly (logarithmically) with energy.
- ✓ Similar generic behaviour found for W and DY.



$d\sigma / dp_T$  for  $gg \rightarrow \text{Higgs}(125 \text{ GeV})$ , PDF set: CT10nlo &  $(\mu_f, \mu_R) = (m_X, m_X)$



- ✓ The  $p_T$ -distribution gets harder with increasing  $\sqrt{s}$ .
- ✓ The peak position increases logarithmically with energy, but (much) faster than for Z: Gluons ( $gg \rightarrow H$ ) radiate more than quarks ( $q, \bar{q} \rightarrow Z$ ).



$d\sigma / dp_T$  for  $t\bar{t}$ , PDF set: CT10nlo &  $(\mu_f, \mu_R) = (m_X, m_X)$ 


- ✓ The  $p_T$ -distribution gets harder with increasing  $\sqrt{s}$ .
- ✓ The peak position increases faster than for H or Z:  $t\bar{t}$  is heavier than Higgs and it's mostly produced by gluons (which radiate more than  $q, \bar{q} \rightarrow Z$ ).

# $\sqrt{s}$ - evolution of maximum peak (DY, W, Z, H, $t\bar{t}$ )

POWHEG + PYTHIA

POWHEG + HERWIG



❑ HERWIG tends to predict smaller  $p_T$  (peak) than PYTHIA.


❑ Logarithmic and power-law fits

I. At threshold, minimum  $p_T^{\text{peak}} = 2 \text{ GeV}$  (intrinsic parton  $k_T$ ).

II. The Slope increases as  $\log(\sqrt{s})$  for DY, W, Z, H but faster, as a power law ( $s^n$ ), for  $t\bar{t}$ .

III. The slope is higher for heavier particles (higher virtuality to radiate) and for gluon-induced processes (compared to quark-induced)

Soft gluon radiation is larger for gluons (two colours) than for quarks (1 colour) by a factor:





$$2N_c^2 / N_c^2 - 1 = 2.25$$

# $\sqrt{s} / m_X$ - evolution of maximum peak (DY, W, Z, H, $t\bar{t}$ )

Evolution as a function of normalized  $\sqrt{s} / m_X$  factorizes out the effects due to different masses of the produced systems:


**POWHEG + PYTHIA**

**POWHEG + HERWIG**



Logarithmic and power-law fits



- I. At threshold, minimum  $p_T^{\text{peak}} = 2 \text{ GeV}$  (intrinsic parton  $k_T$ ).
- II. The slope increases as  $\log(\sqrt{s})$  for DY, W, Z, H but faster, as a power law ( $s^n$ ), for  $t\bar{t}$ .
- III. The slope is higher for heavier particles and for gluon-induced processes (compared to quark-induced).



Soft gluon radiation is larger for gluons (two colours) than for quarks (1 colour) by a factor:


$$2N_c^2 / N_c^2 - 1 = 2.25$$

$d\sigma/dp_T$  for  $gg \rightarrow \text{Higgs}(125 \text{ GeV})$  for  $\sqrt{s} = 8 \text{ TeV}$  for different PDF sets and scales





Higgs peak position robust wrt. changes of scales.

Higgs peak position is a robust observable, consistent with other calculations (HRes, MadGraph, aMCatNLO and SHERPA+NLO, arXiv:1307.1347 [hep-ph]).




$d\sigma / dp_T$  for  $t\bar{t}$  for  $\sqrt{s} = 8$  TeV for different PDF sets and scales




$t\bar{t}$  Peak position robust wrt. changes of scales.


Experimental spectrum should be binned more finely to study low  $p_T$  peak position.




$\sqrt{s} / m_X$  - evolution of maximum peak of H for different PDF sets and scales




*Position of the peak is robust with respect to scale & PDF uncertainties.*



$\sqrt{s} / m_X$  - evolution of maximum peak of W for different PDF sets and scales



Position of the peak is robust with respect to scale & PDF uncertainties.





# Summary

- ✓ The peak of  $d\sigma / dp_T$  distribution of a heavy particle produced in p-p collisions is mostly determined by soft-gluon emission of the colliding partons.
- ✓ We study the evolution of the peak of  $d\sigma / dp_T$  as a function of the mass of the particle and  $\sqrt{s}$  using NLO+"NLL" soft-gluon resummation tools (POWHEG + parton-shower: PYTHIA or HERWIG) for the following systems: DY, W, Z, H,  $t\bar{t}$  in the range  $\sqrt{s} \sim \text{threshold} - 100 \text{ TeV}$ .
- ✓ The  $p_T$ -distribution for all particles gets harder with increasing  $\sqrt{s}$
- ✓ HERWIG tends to predict smaller  $p_T$  (peak) than PYTHIA.
- ✓ The peak position increases logarithmically with energy and with the mass of the system.
- ✓ The speed of increase of  $p_T$  (peak) is faster for gluon-dominated processes (Higgs,  $t\bar{t}$ ) than for quark-induced ones (DY, W, Z) as expected: gluons radiate 2.25 times more than quarks.
- ✓ Position of the peak is robust with respect to scale & PDF uncertainties.

# Outlook

- We want to cross-check the POWHEG + parton-shower results with those obtained from analytical NNLO+NNLL calculations (e.g. RESBOS and DYRES for Z boson).
- We'll try to determine the quantitative scaling-law governing the increase of  $p_T$  (peak) with  $\sqrt{s}$  and  $m_X$ .

THANKS

# Backup slides

## Analytical $Q_T$ resummation

## Parton showering programs (Pythia, MC@NLO, Sherpa...)

evaluate(s) effects of multiple parton radiation in hadronic scattering

applies to a restricted class of processes and observables (e.g., lepton distributions in Drell-Yan-like processes); inclusive with respect to hadronic radiation

apply to a wide range of observables; exclusive with respect to hadronic radiation

is proved to all orders in the QCD coupling by special factorization theorems devised for each qualified observable

no factorization proofs for individual observables

streamlined computation of higher-order corrections and high- $p_T$  contributions

beyond the leading order, radiative contributions and high- $p_T$  tails may be difficult to implement

resummation of all logarithms  $\ln Q_T^2/Q^2$

resummation of leading logarithms  $\ln Q_T^2/Q^2$


nonperturbative contributions are constrained by invoking their universality in the considered class of processes

nonperturbative scattering is evaluated in one of several available models

more strict and precise; relies on first principles of perturbative QCD


more flexible; more parameters to tune to describe various hadronic scattering effects

## POsitive W eight H ardest E mission G enerator

- It's a method to generate the hardest emission first, with NLO accuracy, independently from the subsequent shower.
  - It generates events with **positive weights** only. **No negative weights** to handle.
  - It can be interfaced with any SMC program (HERWIG,PYTHIA,SHERPA,...) which comply with the Les Houches User Process Interface and has the capability to veto emissions harder than the first (SCALUP). It is thus possible to compare different outputs.
  - Can use existing NLO calculations with little effort. No need to be a SMC expert to implement them.
- 

# Shower Monte Carlo

- Initial or final-state **collinear** and **soft emission** (always with  $k_T > \Lambda_{QCD}$ ) are **strongly enhanced**, due to the vanishing denominator in the propagator of the parent.



$$\frac{1}{(q+g)^2 - m_q^2} = \frac{1}{2E_g E_q (1 - \beta_q \cos \theta_{qg})}$$

soft divergence if  $E_g \rightarrow 0$

collinear if  $\theta_{qg} \rightarrow 0$  (only if  $m_q = 0$ )


- Shower algorithms evaluate all these enhanced contributions at all orders
- They give a description of hard collisions up to a distance scale of the order  $1/\Lambda_{QCD}$ .
- At larger distance perturbation theory breaks down and we need to rely on non perturbative methods (i.e. lattice).

# Shower Monte Carlo

- SMC's contain a large library of hard SM and BSM cross sections.
- They dress the hard event with QCD radiation that enhances the cross section in the soft or collinear limit. **From this the name shower.**
- They contain models for hadron formation.
- They handle unstable particle decays.

Thanks to factorization theorem we can separate “hard” physics from “soft” one.  
Parton shower is the link between the two.

Moreover, theoretical basis of the shower are process and energy independent. In principle, once tuned at a certain energy, SMC's have predictive power to all other energies.





## Differences between HERWIG and PYTHIA

They share many common features but


- Their difference is in particular in the treatment of the non-perturbative processes. The PYTHIA philosophy, in fact, is to describe also the hadronisation processes in as much detail as possible.
- They are slightly different also in the description of the hard sub-process.
- Another difference is the scale of the hard scattering,  $\mu^2$  ; PYTHIA sets it to the transverse mass of the two outgoing partons, whereas the scale used by HERWIG is given by the Formula:

$$\mu^2 = \frac{2 \hat{s} \hat{t} \hat{u}}{\hat{s}^2 + \hat{t}^2 + \hat{u}^2}$$

- I will complete this slide when I understand well.


we should include full NLO calculation in SMC's, avoiding double-counting of radiation. This would merge benefits of both approaches:

- ❖ Observables integrated over singular regions are reasonably described by both approaches.
- ❖ At **low**  $p_T$ , exclusive observables, sensitive to IR singularities, are well described by **SMC's**). NLO calculation fails in this region because large logarithms are not properly resummed.
- ❖ If interested in **high  $p_T$**  (LHC, Tevatron) **NLO calculation** is more reliable.




# Results

(POWHEG+PYTHIA)




(POWHEG+HERWIG)




# Results

Local fit for finding peak: Polynomial 3

$W^+ (7 \text{ TeV})$




# Results




POWHEG + PYTHIA

POWHEG + HERWIG




# Results



POWHEG + PYTHIA

POWHEG + HERWIG



$\sqrt{s}/m$

# Conclusions

- All of them start at 2 GeV. **Except  $h\nu q$** . Why?
- As the particles become heavier, the slope becomes greater. It is not the case for  $W^{+/-}$  and  $Z$ . Why?
- Gluon processes have higher peaks than quark processes.

