

ϕ_1 measurements at Belle

Veronika Chobanova

veronika@mpp.mpg.de

May 26th, 2014

Flavor Physics & CP Violation, Marseille




MAX-PLANCK-GESELLSCHAFT

- This talk was supposed to be about $D^* D^* K_S^0$ but analysis not finished yet
- Instead, review of ϕ_1 results with focus on the latest results of

$$B^0 \rightarrow \eta' K^0$$

$$B \rightarrow \omega K$$





Outline

Introduction



$$B^0 \rightarrow \eta' K^0$$

$$B \rightarrow \omega K$$



Measurement of ϕ_1 : $b \rightarrow c\bar{c}s$ Measurement of ϕ_1 in $b \rightarrow c\bar{c}s$ transitions

- Theoretically clean, penguins strongly suppressed
- Experimentally clean, rather free from background



Assuming only tree contribution:

$$\mathcal{A}_{CP} \equiv \frac{|\lambda_{CP}|^2 - 1}{|\lambda_{CP}|^2 + 1} = 0, \quad \mathcal{S}_{CP} \equiv \frac{2\Im(\lambda_{CP})}{|\lambda_{CP}|^2 + 1} = \sin 2\phi_1$$


Measurement of ϕ_1 : $b \rightarrow c\bar{c}s$



$$\sin(2\beta) \equiv \sin(2\phi_1)$$

HFAG

Moriond 2014

PRELIMINARY



Measurement of ϕ_1 : $b \rightarrow sq\bar{q}$ 


- Main contribution from loop-processes, tree contributions CKM suppressed
- $S_{CP} = \sin 2\phi_1$, same as $b \rightarrow c\bar{c}s$
- Penguin amplitudes highly sensitive to new physics
- Could be affected by a heavy unknown particle in the loop that can distort the measured $\sin 2\phi_1$ and branching fractions
 \Rightarrow Deviations from the SM predictions could be a hint at new physics



Outline

Introduction

$B^0 \rightarrow \eta' K^0$

$B \rightarrow \omega K$




Phenomenology of $B^0 \rightarrow \eta' K^0$ 

- Neglecting the tree contribution,

$$\mathcal{A}_{CP} = 0 \quad \mathcal{S}_{CP} = -\eta \sin 2\phi_1$$

- Taking into account pollution from a tree process, $\mathcal{S}_{CP} = -\eta \sin 2\phi_1^{eff}$
- SM prediction: $\Delta \mathcal{S}_{\eta' K^0} = \mathcal{S}_{\eta' K^0} - \sin 2\phi_1 \approx (-0.05 - 0.09)$
- $B^0 \rightarrow \eta' K^0$ consists of two CP final states:
 $\eta = 1$ ($\eta' K_L^0$) and $\eta = -1$ ($\eta' K_S^0$)

$B^0 \rightarrow \eta' K^0$ event reconstruction

- K_S^0
 $\rightarrow \pi^+ \pi^-$ and $\pi^0 \pi^0$
- K_L^0
 ECL and/or KLM clusters without associated charged tracks
- η'
 $\rightarrow \rho^0 \gamma$
 $\rightarrow \eta(\gamma\gamma) \pi^+ \pi^-$
 $\rightarrow \eta(3\pi) \pi^+ \pi^-$

Measurement of TCPV in $B^0 \rightarrow \eta' K^0$

- Main background $q\bar{q}$, further contribution from $b \rightarrow u, d, s, c$
- Suppression of $q\bar{q}$ BG with a likelihood ratio $\mathcal{R}_{s/b} = \mathcal{L}_{sig}/(\mathcal{L}_{sig} + \mathcal{L}_{bkg})$
- \mathcal{L} contains a Fisher discriminant built from event-shape variables
- Pre-selection cut at
 - $\mathcal{R}_{s/b} > 0.5$ (K_L^0 modes)
 - $\mathcal{R}_{s/b} > 0.1$ ($K_S^0 \eta' [\rho^0 \gamma]$ mode)

Fit observables for signal yield extraction from full Belle data set

$$B^0 \rightarrow \eta' K_S^0$$

$$5.23 \text{ GeV}/c^2 \leq M_{bc} \leq 5.30 \text{ GeV}/c^2$$

$$-0.2 \text{ GeV} \leq \Delta E \leq 0.2 \text{ GeV}$$

$$0 \leq \mathcal{R}_{s/b} \leq 1$$

$$M_{bc} = \sqrt{E_{beam}^2 - p_B^2}$$

$$\Delta E = E_B - E_{beam}$$

$$B^0 \rightarrow \eta' K_L^0$$




$$0 \text{ GeV}/c \leq p_B^* \leq 2 \text{ GeV}/c$$

$$0.5 \leq \mathcal{R}_{s/b} \leq 1$$

p_B^* : B momentum calculated assuming $\Delta E = 0$ and nominal $m_{K_L^0}$



Measurement of TCPV in $B^0 \rightarrow \eta' K^0$

$$B^0 \rightarrow \eta' K_S^0$$




- total PDF
- $b \rightarrow u, d, s, c$
- BG
- $q\bar{q}$ BG

$$B^0 \rightarrow \eta' K_L^0$$



- total PDF
- comb. BG with fake η'
- comb. BG with fake K_L^0
- comb. BG with real η' and K_L^0


Signal yield: 3941 ± 90.5 events

Measurement of TCPV in $B^0 \rightarrow \eta' K^0$ 

Decay mode	$-\xi_f \mathcal{S}_f$	\mathcal{A}_f
$\eta' K_S^0$	$+0.71 \pm 0.07$	$+0.02 \pm 0.05$
$\eta' K_L^0$	$+0.46 \pm 0.21$	$+0.09 \pm 0.14$
$\eta' K^0$	$+0.68 \pm 0.07 \pm 0.03$	$+0.03 \pm 0.05 \pm 0.04$

Systematic uncertainties $B^0 \rightarrow \eta' K^0$

Source	$\mathcal{S}_{\eta' K^0}$	$\mathcal{A}_{\eta' K^0}$
Vertexing	± 0.014	± 0.033
Δt resolution	± 0.025	± 0.006
$\eta' K_S^0$ Signal fraction	± 0.013	± 0.006
$\eta' K_L^0$ Signal fraction	± 0.005	± 0.004
Background Δt PDF	± 0.001	< 0.001
Physics parameters	± 0.001	< 0.001
Possible fit bias	± 0.001	± 0.001
Flavor tagging	± 0.003	± 0.003
Tag-side interference	± 0.001	± 0.020
Total	± 0.032	± 0.040

Measurement of TCPV in $B^0 \rightarrow \eta' K^0$ 

- Results consistent with previous Belle measurement

$$\mathcal{A}_{CP} = 0.01 \pm 0.07(\text{stat.}) \pm 0.05(\text{sys.})$$

$$\mathcal{S}_{CP} = 0.64 \pm 0.10(\text{stat.}) \pm 0.04(\text{sys.})$$


- World's most precise measurement to date of $B^0 \rightarrow \eta' K^0$ CP parameters
- \mathcal{A}_{CP} and \mathcal{S}_{CP} values consistent with SM predictions


Outline

Introduction

$B^0 \rightarrow \eta' K^0$

$B \rightarrow \omega K$



Phenomenology of $B \rightarrow \omega K$ 

- Taking into account pollution from a tree process, $\mathcal{S}_{CP} = -\eta \sin 2\phi_1^{\text{eff}}$
- SM prediction: $\Delta S_{\omega K_S^0} = S_{\omega K_S^0} - \sin 2\phi_1 \approx (0.1 - 0.2)$
- $B^0 \rightarrow \omega K_S^0$ is a CP final state with $\eta = -1$

$B \rightarrow \omega K$ measurement

$B \rightarrow \omega K$ reconstruction

$B^0 \rightarrow \omega K_S^0$ and $B^+ \rightarrow \omega K^+$

$\omega \rightarrow \pi^+ \pi^- \pi^0$, $\pi^0 \rightarrow \gamma \gamma$

$K_S^0 \rightarrow \pi^+ \pi^-$


- Main BG contribution from $q\bar{q}$ events
- Suppressed through \mathcal{L} that contains a Fisher discriminant with event-shape variables
- After cut $\mathcal{L} > 0.2$, transformation






$$\mathcal{F}_{B\bar{B}/q\bar{q}} = \log \frac{\mathcal{L} - 0.2}{1 - \mathcal{L}}$$

Extraction of $\mathcal{B}(\omega K^0)$, $\mathcal{B}(\omega K^+)$, $\mathcal{A}_{\omega K_S^0}$, $\mathcal{S}_{\omega K_S^0}$, $\mathcal{A}_{\omega K^+}$ from the full Belle data set.

Simultaneous extended 7D ML fit of

- ΔE
- M_{bc}
- $\mathcal{F}_{B\bar{B}/q\bar{q}}$
- $m(3\pi)$: mass of ω candidates
- $\mathcal{H}(3\pi)$: helicity of ω candidates
- Δt , q



$B^0 \rightarrow \omega K_S^0$ results

- total PDF
- - qq \bar{q} BG
- . . . BB \bar{B} BG
- /// signal

$B \rightarrow \omega K$ results

Black font: previous measurements

Blue font: Full Belle data set of 772×10^6 $B\bar{B}$ pairs


	$B\bar{B}$ -pairs	$\mathcal{BR}(B^0 \rightarrow \omega K^0)$	\mathcal{A}_{CP}	S_{CP}
Belle	388×10^6	$(4.4^{+0.8}_{-0.7} \pm 0.4) \times 10^{-6}$	-	-
Belle	535×10^6	-	$-0.09 \pm 0.29 \pm 0.06$	$0.11 \pm 0.46 \pm 0.07$
BaBar	467×10^6	$(5.4 \pm 0.8 \pm 0.3) \times 10^{-6}$	$0.52^{+0.22}_{-0.20} \pm 0.03$	$0.55^{+0.26}_{-0.29} \pm 0.02$
Belle	772×10^6	$(4.5 \pm 0.4 \pm 0.3) \times 10^{-6}$	$-0.36 \pm 0.19 \pm 0.05$	$0.91 \pm 0.32 \pm 0.05$

	$B\bar{B}$ -pairs	$\mathcal{BR}(B^+ \rightarrow \omega K^+)$	\mathcal{A}_{CP}
Belle	388×10^6	$(8.1 \pm 0.6 \pm 0.6) \times 10^{-6}$	$0.05^{+0.08}_{-0.07} \pm 0.01$
BaBar	383×10^6	$(6.3 \pm 0.5 \pm 0.3) \times 10^{-6}$	$-0.01 \pm 0.07 \pm 0.01$
Belle	772×10^6	$(6.8 \pm 0.4 \pm 0.4) \times 10^{-6}$	$-0.03 \pm 0.04 \pm 0.01$


Paper submitted to PRD, in review; arXiv 1311.6666

$B \rightarrow \omega K$ systematic uncertainties

Category	$\delta\mathcal{B}(\omega K_S^0)$ (%)	$\delta\mathcal{A}_{\omega K_S^0}$ (10^{-2})	$\delta\mathcal{S}_{\omega K_S^0}$ (10^{-2})	$\delta\mathcal{B}(\omega K^+)$ (%)	$\delta\mathcal{A}_{\omega K^+}$ (10^{-2})
$N_{B\bar{B}}$	1.4	N/A	N/A	1.4	N/A
π^0 reconstruction	4.0	N/A	N/A	4.0	N/A
K_S^0 reconstruction	0.8	N/A	N/A	N/A	N/A
PID	1.8	N/A	N/A	2.8	N/A
Tracking	0.7	N/A	N/A	1.1	N/A
Vertex quality selection	0.9	0.3	0.5	0.9	N/A
Δt resolution function	0.6	2.6	4.4	0.8	0.7
Flavor-tagging	0.0	0.3	0.8	0.0	N/A
Misreconstruction	0.9	0.1	0.3	0.7	0.1
$B\bar{B}$ background yields	0.8	0.2	0.5	0.9	0.3
Parametric shape	1.8	0.5	1.5	1.0	0.5
Nonparametric shape	0.1	0.1	0.2	0.1	0.3
Fit bias	0.6	0.7	0.1	0.9	0.3
Background CP violation	N/A	1.5	1.4	N/A	0.1
Tag-side interference	N/A	3.2	0.2	N/A	N/A
Total	5.5	4.6	5.2	5.6	1.0

Measurement of ϕ_1 : $B^0 \rightarrow \omega K_S^0$ First evidence of CP violation in $B^0 \rightarrow \omega K_S^0$ 

Clear asymmetry can be seen in the difference between the B^0 and \bar{B}^0 distributions




CP conservation $(\mathcal{A}_{CP}, \mathcal{S}_{CP}) = (0, 0)$ ruled out by 3.1 standard deviations

Measurement of ϕ_1 : $b \rightarrow sq\bar{q}$

$$\sin(2\beta^{\text{eff}}) \equiv \sin(2\phi_1^{\text{eff}})$$

HFAG
Moriond 2014
PRELIMINARY



Naïve $b \rightarrow sq\bar{q}$ average 0.655 ± 0.032

Conclusion

- Most precise measurements of ϕ_1 in $b \rightarrow c\bar{c}s$ transitions
- $b \rightarrow sq\bar{q}$ decays also sensitive to ϕ_1 , measured value could be affected by new physics
- $B^0 \rightarrow \eta' K^0$ CP parameters results are world's most precise measurement, consistent with previous measurements
- $B \rightarrow \omega K$ measurements of branching fractions and CP parameters mostly consistent with previous results, four out of five parameters world's most precise results
- $b \rightarrow sq\bar{q}$ results consistent with ϕ_1 from $b \rightarrow c\bar{c}s$ transitions, more precision needed
- More on $b \rightarrow sq\bar{q}$ transitions in the talk of Eugenia Puccio

Thank you for your attention


BACK UP

Measurement of TCPV in $B^0 \rightarrow \eta' K^0$

Fit performed in each tagging-quality bin and for each decay mode separately

K^0 mode	η' mode	signal region		$+\mathcal{R}_{s/b} > 0$.
		N_{sig}	purity	purity
$K_S^0 \rightarrow \pi^+ \pi^-$	$\rho^0 \gamma$	1410.5 ± 48.5	0.19	0.59
	$\eta(\gamma\gamma)\pi^+\pi^-$	648.3 ± 27.9	0.49	0.89
	$\eta(3\pi)\pi^+\pi^-$	174.3 ± 13.5	0.65	0.94
$K_S^0 \rightarrow \pi^0 \pi^0$	$\rho^0 \gamma$	162.2 ± 21.4	0.04	0.13
	$\eta(\gamma\gamma)\pi^+\pi^-$	104.0 ± 14.2	0.16	0.65
K_L	$\eta(\gamma\gamma)\pi^+\pi^-$	829.2 ± 54.0	0.30	
	$\eta(3\pi)\pi^+\pi^-$	612.5 ± 36.1	0.19	
Total		3941.0 ± 90.5		

Higher signal yield than previous analysis (+ 20%) due to improved tracking and K_S^0 selection

$\omega K_S C_{CP}$
HFAG
 Morioud 2014
 PRELIMINARY
 $\omega K_S S_{CP}$
HFAG
 Morioud 2014
 PRELIMINARY
