

# Agravity

- 0) From the weak scale...
- 1) ...to the Planck scale...
- 2) ...and above...
- 3) ...up to infinite energy.


Alessandro Strumia  
Talk at Planck 2014

# 0) What was found

But should not have been found


# Only the Higgs

$m_h = 125.6$  GeV




# The SM Higgs


Fit to Higgs couplings




New physics only in loops





Composite Higgs



MSSM fit ( $\tan\beta \gg 1$ )






Higgs or dilaton?



**And nothing else**

# Maybe up to the Planck scale

For the measured  $M_h$ ,  $M_t$  the SM can be extrapolated up to  $M_{Pl}$ .  
And is close to vacuum meta-stability.



For the measured masses even the  $\beta$ -function of  $\lambda \sim$  vanishes around  $M_{Pl}$

$$\lambda = \beta_\lambda = 0 \text{ at } M_{Pl}$$

# The SM parameters at NNLO

SM parameters extracted with data at 2 loop accuracy: at  $\bar{\mu} = M_t$

$$g_2 = 0.64822 + 0.00004 \left( \frac{M_t}{\text{GeV}} - 173.10 \right) + 0.00011 \frac{M_W - 80.384 \text{ GeV}}{0.014 \text{ GeV}}$$

$$g_Y = 0.35761 + 0.00011 \left( \frac{M_t}{\text{GeV}} - 173.10 \right) - 0.00021 \frac{M_W - 80.384 \text{ GeV}}{0.014 \text{ GeV}}$$

$$y_t = 0.9356 + 0.0055 \left( \frac{M_t}{\text{GeV}} - 173.10 \right) - 0.0004 \frac{\alpha_3(M_Z) - 0.1184}{0.0007} \pm 0.0005_{\text{th}}$$

$$\lambda = 0.1271 + 0.0021 \left( \frac{M_h}{\text{GeV}} - 125.66 \right) - 0.00004 \left( \frac{M_t}{\text{GeV}} - 173.10 \right) \pm 0.0003_{\text{th}}$$

$$\frac{m}{\text{GeV}} = 132.03 + 0.94 \left( \frac{M_h}{\text{GeV}} - 125.66 \right) + 0.17 \left( \frac{M_t}{\text{GeV}} - 173.10 \right) \pm 0.15_{\text{th}}.$$


Renormalization to large energies is done with 3 loop RGE.

[Buttazzo, Degrassi, Giardino, Giudice, Sala, Salvio, Strumia, 1307.3536]

# What is this talk about?

In the past decades, theory was driven by the naturalness principle:  
 “light fundamental scalars cannot exist, unless they are accompanied by new physics that protects their mass from quadratically divergent corrections”.

Theorists proposed a beautiful plausible scenario with beautiful LHC signals:



**But LHC found the higgs and nothing else so far.**

I assume that this will be the final outcome and reconsider the basic question.

The goal of this talk is presenting an alternative: a renormalizable theory valid above  $M_{\text{Pl}}$  such that  $M_h$  is naturally smaller than  $M_{\text{Pl}}$  without new physics at the weak scale. It naturally gives inflation and an anti-graviton ghost-like.



# 1) Finite Naturalness

# The good, the bad, the ugly

The **good possibility** of naturalness is in trouble.

The **bad possibility** is that the Higgs is light because of ant\*\*pic selection.

The **ugly possibility** is that **quadratic divergences vanish and a modified Finite Naturalness applies.**

Power divergences are unphysical, nobody knows if they vanish or not. The answer is chosen by the ultimate unknown physical cut-off. Surely it is not a Lorentz-breaking lattice. Maybe it behaves like dimensional regularization.

To start, I explore if this heresy can work and find its consequences and tests.




“Finite naturalness is here considered only as a pure mathematical hypothesis without any pretence of truth”



# Iipse undixt

Wilson proposed the usual naturalness attributing a physical meaning to momentum shells of power-divergent loop integrals, used in the 'averaged action'.




*“The final blunder was a claim that scalar elementary particles were unlikely to occur in elementary particle physics at currently measurable energies unless they were associated with some kind of broken symmetry. The claim was that, otherwise, their masses were likely to be far higher than could be detected. The claim was that it would be unnatural for such particles to have masses small enough to be detectable soon.*

*But this claim makes no sense”*

*Kenneth G. Wilson — Dec. 2004*

# The SM satisfies Finite Naturalness

Quantum corrections to the dimensionful parameter  $m^2 \simeq M_h^2$  in the SM Lagrangian  $\frac{1}{2}m^2|H|^2 - \lambda|H|^4$  are small for the measured values of the parameters



From arXiv:1303.7244

$$M_h = 125.6 \text{ GeV} \Rightarrow m(\bar{\mu} = M_t) = 132.7 \text{ GeV} \Rightarrow m(\bar{\mu} = M_{Pl}) = 140.9 \text{ GeV}$$

# Finite Naturalness and new physics

**FN would be ruined by new heavy particles too coupled to the SM.**

Unlike in the other scenarios, high-scale model building is very constrained. Imagine there is no GUT. No flavour models too. Above us only sky.

FN holds if the top really is the top — if the weak scale is the highest scale.

**Data demand some new physics: DM, neutrino masses, maybe axions...**

**FN still holds if such new physics lies not much above the weak scale.**

Is this possible? If yes what are the signals?

# Finite Naturalness and new physics

**Neutrino mass** models add extra particles with mass  $M$


$$M \lesssim \begin{cases} 0.7 \cdot 10^7 \text{ GeV} \times \sqrt[3]{\Delta} & \text{type I see-saw model,} \\ 200 \text{ GeV} \times \sqrt{\Delta} & \text{type II see-saw model,} \\ 940 \text{ GeV} \times \sqrt{\Delta} & \text{type III see-saw model.} \end{cases}$$

**Leptogenesis** is compatible with FN only in type I.

**Axion** and LHC usually are like fish and bicycle because  $f_a \gtrsim 10^9 \text{ GeV}$ . Axion models can satisfy FN, e.g. KSVZ models employ heavy quarks with mass  $M$

$$M \lesssim \sqrt{\Delta} \times \begin{cases} 0.74 \text{ TeV} & \text{if } \Psi = Q \oplus \bar{Q} \\ 4.5 \text{ TeV} & \text{if } \Psi = U \oplus \bar{U} \\ 9.1 \text{ TeV} & \text{if } \Psi = D \oplus \bar{D} \end{cases}$$

**Inflation**: flatness implies small couplings. Gravity gives an upper bound on  $H_I$  and on any mass [Arvintataki, Dimopoulos..]



$$\delta m^2 \sim \text{[Diagram]} \sim \frac{y_t^2 M^6}{M_{\text{Pl}}^4 (4\pi)^6} \quad \text{so} \quad M \lesssim \Delta^{1/6} \times 10^{14} \text{ GeV}$$

**Dark Matter**: extra scalars/fermions with/without weak gauge interactions.

# DM with EW gauge interactions

Consider a Minimal Dark Matter  $n$ -plet. 2-loop quantum corrections to  $M_h^2$ :


$$\delta m^2 = \frac{cnM^2}{(4\pi)^4} \left( \frac{n^2 - 1}{4} g_2^4 + Y^2 g_Y^4 \right) \times \begin{cases} 6 \ln \frac{M^2}{\Lambda^2} - 1 & \text{for a fermion} \\ \frac{3}{2} \ln^2 \frac{M^2}{\Lambda\mu^2} + 2 \ln \frac{M^2}{\Lambda^2} + \frac{7}{2} & \text{for a scalar} \end{cases}$$

Quantum numbers $SU(2)_L$ $U(1)_Y$ Spin	DM could decay into	DM mass in TeV	$m_{DM^\pm} - m_{DM}$ in MeV	Finite naturalness bound in TeV, $\Lambda \sim M_{Pl}$	$\sigma_{SI}$ in $10^{-46} \text{ cm}^2$
2    1/2    0	$EL$	0.54	350	$0.4 \times \sqrt{\Delta}$	$(2.3 \pm 0.3) 10^{-2}$
2    1/2    1/2	$EH$	1.1	341	$1.9 \times \sqrt{\Delta}$	$(2.5 \pm 0.8) 10^{-2}$
3    0    0	$HH^*$	2.5	166	$0.22 \times \sqrt{\Delta}$	$0.60 \pm 0.04$
3    0    1/2	$LH$	2.7	166	$1.0 \times \sqrt{\Delta}$	$0.60 \pm 0.04$
3    1    0	$HH, LL$	1.6+	540	$0.22 \times \sqrt{\Delta}$	$0.06 \pm 0.02$
3    1    1/2	$LH$	1.9+	526	$1.0 \times \sqrt{\Delta}$	$0.06 \pm 0.02$
4    1/2    0	$HHH^*$	2.4+	353	$0.14 \times \sqrt{\Delta}$	$1.7 \pm 0.1$
4    1/2    1/2	$(LHH^*)$	2.4+	347	$0.6 \times \sqrt{\Delta}$	$1.7 \pm 0.1$
4    3/2    0	$HHH$	2.9+	729	$0.14 \times \sqrt{\Delta}$	$0.08 \pm 0.04$
4    3/2    1/2	$(LHH)$	2.6+	712	$0.6 \times \sqrt{\Delta}$	$0.08 \pm 0.04$
5    0    0	$(HHH^*H^*)$	9.4	166	$0.10 \times \sqrt{\Delta}$	$5.4 \pm 0.4$
5    0    1/2	stable	10	166	$0.4 \times \sqrt{\Delta}$	$5.4 \pm 0.4$
7    0    0	stable	25	166	$0.06 \times \sqrt{\Delta}$	$22 \pm 2$


# DM without EW gauge interactions

DM coupling to the Higgs determines  $\Omega_{\text{DM}}$ ,  $\sigma_{\text{SI}}$  and Finite Naturalness  $\delta m^2$

scalar DM singlet



Fermion DM singlet ( $m_S=300$  GeV)



Observable DM satisfies Finite Naturalness if lighter than  $\approx 1$  TeV



## **2) A new principle**

Finite Naturalness is phenomenologically viable, what about its theory?

# Nature has no scale

FN needs something different from the effective field theory ideology

$$\mathcal{L} \sim \Lambda^4 + \Lambda^2 |H|^2 + \mathcal{L}_4 + \frac{H^6}{\Lambda^2} + \dots$$

that leads to the hierarchy problem. Nature is singling out  $\mathcal{L}_4$ . Why?

**Principle: “Nature has no fundamental scales  $\Lambda$ ”.**

Then, the fundamental QFT is described by  $\mathcal{L}_4$ : only a-dimensional couplings.

Power divergences vanish simply because they have mass dimension, and there are no masses. Scale invariance at tree level is an accidental symmetry, like baryon number. [Other authors assume scale or conformal invariance as quantum symmetries and argue that the regulator must respect them].

**Quantum corrections break scale invariance and should generate  $M_h, M_{\text{Pl}}$**

Can this happen? I apply this principle first to  $M_h$  and later to  $M_{\text{Pl}}$ .

# What is the weak scale?

- Could be the only scale of particle physics. Just so.
- Could be generated from nothing by heavier particles.
- Could be generated from nothing by weak-scale dynamics. Like QCD.

# Dynamical generation of the weak scale

## Goals:

- 1) **Dynamically generate** the weak scale and weak scale DM
- 2) **Preserve** the successful automatic features of the SM:  $B, L...$
- 3) **Get DM stability** as one extra automatic feature.

## Model:

$G_{\text{SM}} \otimes \text{SU}(2)_X$  with one extra scalar  $S$ , doublet under  $\text{SU}(2)_X$  and potential

$$V = \lambda_H |H|^4 - \lambda_{HS} |HS|^2 + \lambda_S |S|^4.$$

[Hambye, Strumia, 1306.2329]

# Dynamical generation of the weak scale

1)  $\lambda_S$  runs negative at low energy:

$$\lambda_S \simeq \beta_{\lambda_S} \ln \frac{s}{s_*} \quad \text{with} \quad \beta_{\lambda_S} \simeq \frac{9g_X^4}{8(4\pi)^2}$$

$$S(x) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ w + s(x) \end{pmatrix} \quad w \simeq s_* e^{-1/4}$$


$$H(x) = \frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ v + h(x) \end{pmatrix} \quad v \simeq w \sqrt{\frac{\lambda_{HS}}{2\lambda_H}}$$

Problem: vacuum energy must be negative???

2) No new Yukawas.

3)  $SU(2)_X$  vectors get mass  $M_X = \frac{1}{2}g_X w$  and are automatically stable.

4) Bonus: threshold effect stabilises  $\lambda_H = \lambda + \lambda_{HS}^2 / \beta_{\lambda_S}$ .





# Experimental implications

- 1) New scalar  $s$ : like another  $h$  with suppressed couplings;  $s \rightarrow hh$  if  $M_s > 2M_h$ .
- 2) Dark Matter coupled to  $s, h$ . Assuming that DM is a thermal relict

$$\sigma v_{\text{ann}} + \frac{1}{2}\sigma v_{\text{semi-ann}} = \frac{11g_X^2}{1728\pi w^2} + \frac{g_X^2}{64\pi w^2} \approx 2.2 \times 10^{26} \frac{\text{cm}^3}{\text{s}}$$

fixes  $w = g_X \times 2 \text{ TeV}$ , so all is predicted in terms of one parameter  $g_X$ :





(Insignificant hint in  $ZZ$  and  $\gamma\gamma$  data around 143 GeV)

# Dark/EW phase transition

The model predicts a first order phase transition for  $s$

The universe remains trapped at  $s = 0$  until the potential energy  $\Delta V$  is violently released via thermal tunnelling:  $\Gamma \sim T^4 e^{-S/T}$  with  $S \propto g_X^4$ .



- For the critical value  $g_X \approx 1.2$  one has  $\Delta V \approx \rho$  such that

$$f_{\text{peak}} \approx 0.3 \text{ mHz} \quad \Omega_{\text{peak}} h^2 \approx 2 \cdot 10^{-11} \quad \text{detectable at LISA}$$

- For  $g_X > 1.2$  gravitational waves become weaker.
- For  $g_X < 1.2$  the universe gets trapped in a (too long?) inflationary phase.

## 3) Agravity

[Salvio, Strumia, 1403.4226]



# What about gravity?

Does quantum gravity give  $\delta M_h^2 \sim M_{\text{Pl}}^2$  ruining Finite Naturalness?

Maybe  $M_{\text{Pl}}^{-1}$  is just a small coupling and there are no new particles around  $M_{\text{Pl}}$ .

Quantum gravity would be very different from what strings suggest...

# Adimensional gravity

Applying the adimensional principle to the SM plus gravity and a scalar  $S$  gives:

$$\mathcal{S} = \int d^4x \sqrt{|\det g|} \mathcal{L}$$

$$\mathcal{L} = \mathcal{L}_{\text{SM}} + \frac{R^2}{3f_0^2} + \frac{R^2 - 3R_{\mu\nu}^2}{3f_2^2} - \xi_H |H|^2 R + |D_\mu S|^2 - \xi_S |S|^2 R - \lambda_S |S|^4 + \lambda_{HS} |HS|^2$$

where  $f_0, f_2$  are the adimensional 'gauge couplings' of gravity and  $R \sim \partial_\mu \partial_\nu g_{\mu\nu}$ .

Of course the theory is renormalizable, and indeed the graviton propagator is:

$$\frac{-i}{k^4} \left[ 2f_2^2 P_{\mu\nu\rho\sigma}^{(\text{spin } 2)} - f_0^2 P_{\mu\nu\rho\sigma}^{(\text{spin } 0)} + \text{gauge-fixing} \right].$$

The Planck scale should be generated dynamically as  $\xi_S \langle S \rangle^2 = \bar{M}_{\text{Pl}}^2/2$ .

Then, the spin-0 part of  $g_{\mu\nu}$  gets a mass  $M_0 \sim f_0 M_{\text{Pl}}$  and the spin 2 part splits into the usual graviton and an **anti-graviton** with mass  $M_2 = f_2 \bar{M}_{\text{Pl}}/\sqrt{2}$  that acts as a Pauli-Villars in view its **negative kinetic term** [Stelle, 1977].

# A ghost?

Classically, higher derivatives are bad [Ostrogradski, 1850]:

$\partial^4 \Rightarrow$  unbounded negative kinetic energy  $\Rightarrow$  the theory is dead.

The dispersion relation  $P^4 = m^4$  has 4 solutions:  $E = \pm m$  and  $E = \pm im$ .

In presence of masses,  $\partial^4$  can be decomposed as 2 fields with 2 derivatives:

$$\frac{1}{k^4} \rightarrow \frac{1}{k^4 - M_2^2 k^2} = \frac{1}{M_2^2} \left[ \frac{1}{k^2} - \frac{1}{k^2 - M_2^2} \right]$$

Quantistically, the state with negative kinetic term can be reinterpreted as **positive energy and negative norm** by swapping  $a \leftrightarrow a^\dagger$ .


This is the  $i\epsilon$  choice that makes the theory renormalizable.

Lee, Wick, Cutkosky... claim that, it gives a slightly acausal unitary  $S$  matrix.


Without masses,  $\partial^4$  cannot be decomposed. Such crackpotton field has its own quantisation rules, I do not yet understand what they mean.

This is what happened with anti-particles: sometimes we have the right equations before understanding what they mean. I ignore the issue and compute.

# A ghost?



## A ghost?



If we knew  
what we were doing  
it wouldn't be research

*Albert Einstein*

# A ghost?



Me ne frego !

# Quantum Gravity...


The quantum behaviour of a renormalizable theory is encoded in its RGE. The unusual  $1/k^4$  makes easy to get signs wrong. Literature is contradictory.

Preliminary results at one loop:

- $f_2$  is asymptotically free:

$$(4\pi)^2 \frac{df_2^2}{d \ln \mu} = -f_2^4 \left[ \frac{133}{10} + \frac{N_V}{5} + \frac{N_f}{20} + \frac{N_s}{60} \right]$$

- Gravity does not affect running of gauge couplings: these two diagrams cancel



presumably because abelian  $g$  is undefined without charged particles.

- $f_0$  is not asymptotically free unless  $f_0^2 < 0$

$$(4\pi)^2 \frac{df_0^2}{d \ln \mu} = \frac{5}{3} f_2^4 + 5 f_2^2 f_0^2 + \frac{5}{6} f_0^4 + \frac{f_0^4}{12} \sum_s (1 + 6\xi_s)^2$$

# ...Quantum Agravity

- Yukawa couplings get an extra multiplicative RGE correction:

$$(4\pi)^2 \frac{dy_t}{d \ln \mu} = \frac{9}{2} y_t^3 - y_t \left( 8g_3^2 - \frac{15}{8} f_2^2 \right)$$

- The RGE for  $\xi$  is perturbative up to  $\xi_H \lesssim 1/f_0$

$$(4\pi)^2 \frac{d\xi_H}{d \ln \mu} = -\frac{5 f_2^4}{3 f_0^2} \xi_H + f_0^2 \xi_H (6\xi_H + 1) \left( \xi_H + \frac{2}{3} \right) + (6\xi_H + 1) \left[ 2y_t^2 - \frac{3}{4} g_2^2 + \dots \right]$$

- Agravity makes quartics small at low energy:

$$(4\pi)^2 \frac{d\lambda_H}{d \ln \mu} = \xi_H^2 [5f_2^4 + f_0^4 (1 + 6\xi_H)^2] - 6y_t^4 + \frac{9}{8} g_2^4 + \dots$$

- Agravity creates a mixed quartic:

$$(4\pi)^2 \frac{d\lambda_{HS}}{d \ln \mu} = \frac{\xi_H \xi_S}{2} [5f_2^4 + f_0^4 (6\xi_S + 1)(6\xi_H + 1)] + \text{multiplicative}$$



# Generation of the Planck scale

Some mechanisms can generate dynamically the Planck scale

a)  $\lambda_S$  runs negative below  $M_{\text{Pl}}$

or

b)  $f_2$  or  $\xi_S$  run non-perturbative.

Focus on a): scalar Planckion.  $\xi_S$  makes the vacuum equations non-standard:

$$\frac{\partial V}{\partial S} - \frac{4V}{S} = 0 \quad \text{i.e.} \quad \frac{\partial V_E}{\partial S} = 0$$

where  $V_E = V/(\xi S^2)^2 \sim \lambda_S(S)/\xi_S^2(S)$  is the Einstein-frame potential. The vev

$$\langle S \rangle = \bar{M}_{\text{Pl}}/\sqrt{2\xi_S}$$

needs a condition different from the usual Coleman-Weinberg:

$$\frac{\beta_{\lambda_S}(\bar{\mu} \sim \langle S \rangle)}{\lambda_S(\bar{\mu} \sim \langle S \rangle)} - 2 \frac{\beta_{\xi_S}(\bar{\mu} \sim \langle S \rangle)}{\xi_S(\bar{\mu} \sim \langle S \rangle)} = 0$$

Furthermore, the cosmological constant is fine-tuned to zero by imposing

$$\lambda_S(\bar{\mu} \sim \langle S \rangle) = 0$$



So the minimum equation simplifies to

$$\beta_{\lambda_S}(\bar{\mu} \sim \langle S \rangle) = 0$$

$\lambda_S = \beta_{\lambda_S} = 0$  around  $M_{\text{Pl}}$ : is this running possible?

# Yes, this is how $\lambda_H$ can run in the SM!

RGE running of the  $\overline{\text{MS}}$  quartic Higgs coupling in the SM



$H$  cannot get a Planck-scale vev. Model: add a mirror copy of the SM, broken by the fact that  $S$ , the Higgs mirror, lies in the Planck minimum:  $\xi_S \sim 10^{1\div 2}$ .

# Inflation = perturbative agravity

Inflation needs special theories with flat potential and/or super-Planckian vevs.

A successful class of models is  $\xi$ -inflation: a scalar  $S$  with  $-\frac{1}{2}f(S)R + V(S)$ .  
Redefine  $g_{\mu\nu} = g_{\mu\nu}^E \times \bar{M}_{\text{Pl}}^2/f$  to the Einstein frame to make the graviton canonical

$$\sqrt{\det g} \left[ -\frac{f}{2}R + \frac{(\partial_\mu s)^2}{2} - V \right] = \sqrt{\det g_E} \left[ -\frac{\bar{M}_{\text{Pl}}^2}{2}R_E + \bar{M}_{\text{Pl}}^2 \left( \frac{1}{f} + \frac{3f'^2}{2f^2} \right) \frac{(\partial_\mu s)^2}{2} - V_E \right]$$

where  $V_E = \bar{M}_{\text{Pl}}^4 V/f^2$  is flat (good for inflation) if  $V(S) \propto f^2(S)$  **above**  $M_{\text{Pl}}$ .

In general, this restriction is unmotivated and uncontrollable.

In quantum agravity  $f(S) = \xi_S(\bar{\mu} \sim S)|S|^2$  and  $V(S) = \lambda_S(\bar{\mu} \sim S)|S|^4!$

Inflation is a typical phenomenon in agravity: the slow-roll parameters are the  $\beta$ -functions, which are small if the theory is perturbative. In the Einstein frame

$$\epsilon \equiv \frac{\bar{M}_{\text{Pl}}^2}{2} \left( \frac{1}{V_E} \frac{\partial V_E}{\partial s_E} \right)^2 = \frac{1}{2} \frac{\xi_S}{1 + 6\xi_S} \left[ \frac{\beta_{\lambda_S}}{\lambda_S} - 2 \frac{\beta_{\xi_S}}{\xi_S} \right]^2,$$

$$\eta \equiv \frac{\bar{M}_{\text{Pl}}^2}{V_E} \frac{\partial^2 V_E}{\partial s_E^2} = \frac{\xi_S}{1 + 6\xi_S} \left[ \frac{\beta(\beta_{\lambda_S})}{\lambda_S} - 2 \frac{\beta(\beta_{\xi_S})}{\xi_S} + \frac{5 + 36\xi_S \beta_{\xi_S}^2}{1 + 6\xi_S} \frac{1}{\xi_S^2} - \frac{7 + 48\xi_S \beta_{\lambda_S} \beta_{\xi_S}}{1 + 6\xi_S} \frac{1}{2\lambda_S \xi_S} \right].$$

# Approximating a gravity inflation

If the inflaton is the Planckion  $s$ , its potential is approximately logarithmic

$$\lambda_S(\bar{\mu} \approx s) \approx 0 + 0 \ln s + \frac{g^4}{2(4\pi)^4} \ln^2 \frac{s}{\langle s \rangle}, \quad \xi_S(\bar{\mu}) \approx \xi_S$$

The canonical Einstein-frame field is

$$s_E = \bar{M}_{\text{Pl}} \sqrt{\frac{1 + 6\xi_S}{\xi_S}} \ln \frac{s}{\langle s \rangle}$$

and its potential is:

$$V_E = \frac{\bar{M}_{\text{Pl}}^4 \lambda_S}{4 \xi_S^2} \approx \frac{M_s^2}{2} s_E^2 \quad \text{with} \quad M_s = \frac{g^2 \bar{M}_{\text{Pl}}}{2(4\pi)^2} \frac{1}{\sqrt{\xi_S(1 + 6\xi_S)}}$$

Inflation occurs at  $s_E \approx 2\sqrt{N} \bar{M}_{\text{Pl}}$  for  $N \approx 60$ : **above** the Planck scale:

$$A_s \approx \frac{g^4 N^2}{24\pi^2 \xi_S (1 + 6\xi_S)} \quad n_s \approx 1 - \frac{2}{N} \approx 0.967, \quad r = \frac{A_t}{A_s} \approx \frac{8}{N} \approx 0.13,$$

In general: (3 predictions) – (2 parameters  $\xi_S$  and  $g$ ) = (1 prediction).

In the SM-mirror model  $g \approx 1.0$  so  $\xi_S \approx 230$  i.e.  $\langle s \rangle \approx 1.6 \cdot 10^{17}$  GeV: ok.

# Generation of the Weak scale

RGE running generates  $M_h$  from  $M_{\text{Pl}}$ . 3 regimes:

1) below  $M_{0,2}$ : ignore gravity,  $M_h$  runs logarithmically as in the SM

$$(4\pi)^2 \frac{dM_h^2}{d \ln \bar{\mu}} = \beta_{\text{SM}} M_h^2 \quad \beta_{\text{SM}} = 12\lambda_H + 6y_t^2 - \frac{9g_2^2}{2} - \frac{9g_1^2}{10}$$

2) between  $M_{0,2}$  and  $M_{\text{Pl}}$ : the apparent masses run:

$$(4\pi)^2 \frac{dM_h^2}{d \ln \bar{\mu}} = \left[ \beta_{\text{SM}} + 5f_2^2 + \frac{5f_2^4}{3f_0^2} + \dots \right] M_h^2 - \xi_H \left[ 5f_2^4 + f_0^4(1 + 6\xi_H) \right] \bar{M}_{\text{Pl}}^2$$

3) above  $M_{\text{Pl}}$  couplings are adimensional:  $\lambda_{HS}|H|^2|S|^2$  leads to  $M_h^2 = \lambda_{HS}\langle s \rangle^2$ :

$$(4\pi)^2 \frac{d\lambda_{HS}}{d \ln \bar{\mu}} = -\xi_H \xi_S [5f_2^4 + f_0^4(6\xi_S + 1)(6\xi_H + 1)] + \dots$$

The weak scale arises if  $f_{0,2} \sim \sqrt{M_h/M_{\text{Pl}}} \sim 10^{-8}$  i.e.  $M_{0,2} \sim 10^{11}$  GeV

All small parameters such as  $f_{0,2}$  and  $\lambda_{HS} \sim f_{0,2}^4$  are naturally small

The Planckion  $s$  can have any mass between  $M_h$  and  $M_{\text{Pl}}$

# Black holes

Perturbative Quantum Gravity cannot convert a small coupling  $1/M_{\text{Pl}}$  into a big mass. Non-perturbative QG, a black hole with mass  $M_{\text{BH}}$ , could give

$$\delta M_h^2 \sim M_{\text{BH}}^2 e^{-M_{\text{BH}}^2/M_{\text{Pl}}^2}.$$

The black holes possibly dangerous for FN have mass  $M_{\text{BH}} \sim M_{\text{Pl}}$ .

Such black holes do not exist if the fundamental coupling of gravity is small.

The minimal mass of a black hole is  $M_{\text{BH}} > M_{\text{Pl}}/f_{0,2}$  because of

$$V_{\text{Newton}} = -\frac{Gm}{r} \left[ 1 - \frac{4}{3}e^{-M_2 r} + \frac{1}{3}e^{-M_0 r} \right]$$

Non-perturbative QG corrections  $\delta M_h^2 \propto e^{-1/f_{0,2}^2}$  can be neglected for  $f_{0,2} \ll 1$

# Landau poles

[Giudice, Isidori, Salvio, Strumia, to appear]

# Landau poles

We have the RGE above  $M_{\text{Pl}}$ , can the theory reach infinite energy?

**Problem:** Landau poles for  $g_Y$ , possibly  $\lambda$ ,  $y_t$ ,  $y_b$ ,  $y_\tau$ ? To analyse any QFT:

1) Get 1-loop RGE, asymptotically approximate

$$g_i = c_i / \ln \bar{\mu} \ll 1$$

2) Get a system of ordinary equations in  $c_i$ .

3) Find multiple sets of solutions  $c_i^1, c_i^2, \dots$


4) Check if at least one physical solution exists, such that all couplings are real.

5) If yes, extrapolate down to low energy.

6) Perturb: UV fixed points admit deformations; IR fixed points are predicted.

In the SM there is one acceptable solution and it predicts  $g_Y, y_\tau = 0$  and, in this limit,  $y_t$  ( $M_t \approx 185.8 \text{ GeV}$ ) and an acceptable range for  $M_h$ .

But  $g_Y \neq 0$  gives a Landau pole at  $10^{43} \text{ GeV}$ .





# Landau poles

Can the SM be extended into a theory valid up to infinite energy?

Idea: **avoid Landau poles by making hypercharge non abelian**. The best possibilities — SU(5)-like GUTs — are not compatible with finite naturalness.

FN demands extensions at the weak scale. There are two possibilities:

$$SU(4)_c \otimes SU(2)_L \otimes SU(2)_R \quad \text{and} \quad SU(3)_c \otimes SU(3)_L \otimes SU(3)_R$$

Flavor, precision data, LHC... imply multi-TeV bounds on some particles ( $H'$ ,  $Z'$ ,  $W'$ ...). Difficult attempt to reconcile bounds with naturalness is underway.

# Conclusions



The exploration is still in progress.  
The truth can be somewhere along this set of ideas.

Of course, going from Higgs and no SUSY to modified naturalness to an anti-graviton ghost at  $10^{11}$  GeV is risky.

Of course, it is much more reasonable to imagine anthropic selection within a multiverse of branes wrapped on 6 or 7 extra dimensions compactified on...