



Contribution ID: 252

Type: **Presentation**

## Observations of gamma-ray bursts with the HAWC observatory

*Monday 23 June 2014 17:40 (15 minutes)*

The temporal evolution and end of GRB spectra have important implications for the acceleration mechanisms of gamma-ray bursts (GRBs). Above  $\approx 10$  GeV the effective area of *Fermi*-LAT is approximately constant and since the photon flux is steeply decreasing with energy, an insufficient number of photons is detected. The High Altitude Water Cherenkov (HAWC) observatory is a gamma-ray detector in the 100 GeV - 100 TeV range currently under construction in central Mexico that has the potential to measure GRB spectra beyond the LAT energy range. Unlike Imaging Atmospheric Cherenkov Telescopes, it has a large field of view and near 100 % duty cycle that will allow for observations of the prompt GRB phase. In this presentation I will show first results of GRB observations with the partially completed HAWC array, including GRB 130427A, the most energetic GRB so far detected at a redshift  $z < 0.5$ .

**Author:** LENNARZ, Dirk (Georgia Tech)

**Co-author:** FOR THE, HAWC collaboration (HAWC collaboration)

**Presenter:** LENNARZ, Dirk (Georgia Tech)

**Session Classification:** Gamma-Ray Astrophysics

**Track Classification:** Gamma-Ray Astrophysics