

Introduction to Transverse Beam Dynamics

Bernhard Holzer, CERN-LHC

The „ not so ideal world “


IV.) Scaling Laws, Mini Beta Insertions,
and all the rest



Lattice Design ... in 10 seconds ... the Matrices

Transformation of the coordinate vector (x, x') in a lattice

$$\begin{pmatrix} x(s) \\ x'(s) \end{pmatrix} = M_{1 \rightarrow 2} \begin{pmatrix} x_0 \\ x'_0 \end{pmatrix}$$



Matrix expressed as function of focusing properties

$$M_{1 \rightarrow 2} = M_{qf} * M_{kf} * M_B * M_{qd} * M_{...}$$

Transformation of the coordinate vector (x, x')
expressed as a function of the twiss parameters


$$M_{1 \rightarrow 2} = \begin{pmatrix} \sqrt{\frac{\beta_2}{\beta_1}} (\cos \psi_{12} + \alpha_1 \sin \psi_{12}) & \sqrt{\beta_1 \beta_2} \sin \psi_{12} \\ \frac{(\alpha_1 - \alpha_2) \cos \psi_{12} - (1 + \alpha_1 \alpha_2) \sin \psi_{12}}{\sqrt{\beta_1 \beta_2}} & \sqrt{\frac{\beta_1}{\beta_2}} (\cos \psi_{12} - \alpha_2 \sin \psi_{12}) \end{pmatrix}$$

And both descriptions are equivalent !!

Lattice Design ... in 10 seconds ... the β -function

Transformation Matrix for half a FODO

$$M_{\text{half cell}} = M_{QF/2} * M_D * M_{QD/2} = \begin{pmatrix} 1 - l_D/\bar{f} & l_D \\ -l_D/\bar{f}^2 & 1 + l_D/\bar{f} \end{pmatrix}$$



nota bene: $\bar{f} = 2 * f$... it is a half quad !



Compare to the twiss parameter form of M

$$M_{1 \rightarrow 2} = \begin{pmatrix} \sqrt{\beta_2} (\cos \psi_{12} + \alpha_1 \sin \psi_{12}) & \sqrt{\beta_1 \beta_2} \sin \psi_{12} \\ \frac{(\alpha_1 - \alpha_2) \cos \psi_{12} - (1 + \alpha_1 \alpha_2) \sin \psi_{12}}{\sqrt{\beta_1 \beta_2}} & \sqrt{\beta_1} (\cos \psi_{12} - \alpha_2 \sin \psi_{12}) \end{pmatrix}$$

In the middle of a foc (defoc) quadrupole of the FoDo we always have $\alpha = 0$, and the half cell will lead us from β_{max} to β_{min}

$$M_{\text{half cell}} = \begin{pmatrix} \sqrt{\frac{\beta}{\beta_0}} \cos \frac{\psi_{\text{cell}}}{2} & \sqrt{\beta \beta_0} \sin \frac{\psi_{\text{cell}}}{2} \\ \frac{-1}{\sqrt{\beta \beta_0}} \sin \frac{\psi_{\text{cell}}}{2} & \sqrt{\frac{\beta}{\beta_0}} \cos \frac{\psi_{\text{cell}}}{2} \end{pmatrix}$$


Scaling law for a FODO cell:



$$\hat{\beta} = \frac{(1 + \sin \frac{\psi_{\text{cell}}}{2})L}{\sin \psi_{\text{cell}}} !$$

$$\bar{\beta} = \frac{(1 - \sin \frac{\psi_{\text{cell}}}{2})L}{\sin \psi_{\text{cell}}} !$$

$$\langle \hat{\beta}(\psi) + \bar{\beta}(\psi) \rangle$$




The maximum and minimum values of the β -function are solely determined by the phase advance and the length of the cell.

Longer cells lead to larger β ... and there is an optimum phase !!

* see advanced CAS lectures, lattice design (e.g. CERN-2006-002)

Lattice Design ... in 10 seconds ... Dispersion

Dispersion in a FoDo Cell:




$$M_{\text{halfcell}} = \begin{pmatrix} C & S \\ C' & S' \end{pmatrix} = \begin{pmatrix} 1 - l_D/\tilde{f} & l_D \\ -l_D/\tilde{f}^2 & 1 + l_D/\tilde{f} \end{pmatrix}$$

rule to calculate the dispersion

$$D(s) = S(s) \int_{s_0}^{s1} \frac{1}{\rho} C(\tilde{s}) d\tilde{s} - C(s) \int_{s_0}^{s1} \frac{1}{\rho} S(\tilde{s}) d\tilde{s}$$

$$\hat{D} = \frac{\ell^2}{\rho} \cdot \frac{(1 + \frac{1}{2} \sin \frac{\psi_{\text{cell}}}{2})}{\sin^2 \frac{\psi_{\text{cell}}}{2}}$$

$$\check{D} = \frac{\ell^2}{\rho} \cdot \frac{(1 - \frac{1}{2} \sin \frac{\psi_{\text{cell}}}{2})}{\sin^2 \frac{\psi_{\text{cell}}}{2}}$$




19.) Chromaticity:

A Quadrupole Error for $\Delta p/p \neq 0$

Influence of external fields on the beam: prop. to magn. field & prop. zu 1/p

dipole magnet $\alpha = \frac{\int B dl}{p/e}$



$$x_D(s) = D(s) \frac{\Delta p}{p}$$

focusing lens $k = \frac{g}{p/e}$




Figure 29: FODO cell

particle having ...
to high energy
to low energy
ideal energy

Chromaticity: Q'

$$k = \frac{g}{p/e} \quad p = p_0 + \Delta p$$

in case of a momentum spread:

$$k = \frac{eg}{p_0 + \Delta p} \approx \frac{e}{p_0} \left(1 - \frac{\Delta p}{p_0}\right) g = k_0 + \Delta k$$

$$\Delta k = -\frac{\Delta p}{p_0} k_0$$

... which acts like a quadrupole error in the machine and leads to a tune spread:


$$\Delta Q = -\frac{1}{4\pi} \frac{\Delta p}{p_0} k_0 \beta(s) ds$$

definition of chromaticity:

$$\Delta Q = Q' \frac{\Delta p}{p} ; \quad Q' = -\frac{1}{4\pi} \oint k(s) \beta(s) ds$$

Where is the Problem ?

Tunes and Resonances



avoid resonance conditions:

$$m Q_x + n Q_y + l Q_s = \text{integer}$$

... for example: $1 Q_x = 1$

... and now again about Chromaticity:

Problem: chromaticity is generated by the lattice itself !!

Q' is a number indicating the size of the tune spot in the working diagram,

Q' is always created if the beam is focussed

→ it is determined by the focusing strength k of all quadrupoles

$$Q' = -\frac{1}{4\pi} \oint k(s) \beta(s) ds$$

k = quadrupole strength


β = betafunction indicates the beam size ... and even more the sensitivity of the beam to external fields

Example: LHC

$$\left. \begin{aligned} Q' &= 250 \\ \Delta p/p &= \pm 0.2 \cdot 10^{-3} \\ \Delta Q &= 0.256 \dots 0.36 \end{aligned} \right\}$$


→ Some particles get very close to resonances and are lost

in other words: the tune is not a point it is a pancake




Tune signal for a nearly uncompensated chromaticity ($Q' \approx 20$)

Ideal situation: chromaticity well corrected, ($Q' \approx 1$)



Tune and Resonances

$$m \cdot Q_x + n \cdot Q_y + l \cdot Q_s = \text{integer}$$



RA e Tune diagram up to 3rd order

... and up to 7th order

*Homework for the operateurs:
find a nice place for the tune
where against all probability
the beam will survive*


Correction of Q':

Need: additional quadrupole strength for each momentum deviation $\Delta p/p$

1.) sort the particles according to their momentum $x_D(s) = D(s) \frac{\Delta p}{p}$



... using the dispersion function




2.) apply a magnetic field that rises quadratically with x (sextupole field)

$$\left. \begin{aligned} B_x &= \tilde{g}xz \\ B_z &= \frac{1}{2} \tilde{g}(x^2 - z^2) \end{aligned} \right\} \frac{\partial B_x}{\partial z} = \frac{\partial B_z}{\partial x} = \tilde{g}x \quad \text{linear rising „gradient“:}$$

Correction of Q':

Sextupole Magnets:




k_1 normalised quadrupole strength

k_2 normalised sextupole strength

$$k_1(\text{sext}) = \frac{\tilde{g} x}{p/e} = k_2 * x$$

$$k_1(\text{sext}) = k_2 * D * \frac{\Delta p}{p}$$




Combined effect of „natural chromaticity“ and Sextupole Magnets:

$$Q' = -\frac{1}{4\pi} \left\{ \int k_1(s) \beta(s) ds + \int k_2 * D(s) \beta(s) ds \right\}$$

You only should not forget to correct Q' in both planes ...
and take into account the contribution from quadrupoles of both polarities.

corrected chromaticity


considering an arc built out of single cells:



$$Q_x = -\frac{1}{4\pi} \left\{ \sum_{F \text{ quad}} k_{yF} \hat{\beta}_x l_{yF} - \sum_{D \text{ quad}} k_{yD} \bar{\beta}_x l_{yD} \right\} + \frac{1}{4\pi} \sum_{F \text{ sext}} k_2^F l_{\text{sext}}^F D_x^F \beta_x^F - \frac{1}{4\pi} \sum_{D \text{ sext}} k_2^D l_{\text{sext}}^D D_x^D \beta_x^D$$

$$Q_y = -\frac{1}{4\pi} \left\{ -\sum_{F \text{ quad}} k_{yF} \bar{\beta}_y l_{yF} + \sum_{D \text{ quad}} k_{yD} \hat{\beta}_y l_{yD} \right\} - \frac{1}{4\pi} \sum_{F \text{ sext}} k_2^F l_{\text{sext}}^F D_x^F \beta_x^F + \frac{1}{4\pi} \sum_{D \text{ sext}} k_2^D l_{\text{sext}}^D D_x^D \beta_x^D$$

20.) Insertions



Insertions

... the most complicated one: the drift space

Question to the audience: what will happen to the beam parameters α , β , γ if we stop focusing for a while ...?

$$\begin{pmatrix} \beta \\ \alpha \\ \gamma \end{pmatrix}_s = \begin{pmatrix} C^2 & -2SC & S^2 \\ -CC' & SC' + S'C & -SS' \\ C'^2 & -2S'C' & S'^2 \end{pmatrix} * \begin{pmatrix} \beta \\ \alpha \\ \gamma \end{pmatrix}_0$$

transfer matrix for a drift:

$$M = \begin{pmatrix} C & S \\ C' & S' \end{pmatrix} = \begin{pmatrix} 1 & s \\ 0 & 1 \end{pmatrix} \longrightarrow \begin{matrix} \beta(s) = \beta_0 - 2\alpha_0 s + \gamma_0 s^2 \\ \alpha(s) = \alpha_0 - \gamma_0 s \\ \gamma(s) = \gamma_0 \end{matrix}$$

β -Function in a Drift:

let's assume we are at a symmetry point in the center of a drift.

$$\beta(s) = \beta_0 - 2\alpha_0 s + \gamma_0 s^2$$


as $\alpha_0 = 0$, $\rightarrow \gamma_0 = \frac{1 + \alpha_0^2}{\beta_0} = \frac{1}{\beta_0}$

and we get for the β function in the neighborhood of the symmetry point


$$\beta(s) = \beta_0 + \frac{s^2}{\beta_0} \quad !!!$$

**At the end of a long symmetric drift space the beta function reaches its maximum value in the complete lattice.
-> here we get the largest beam dimension.**

-> keep l as small as possible



21.) Luminosity



Example: Luminosity run at LHC

$$\beta_{x,y} = 0.55 \text{ m} \quad f_0 = 11.245 \text{ kHz}$$

$$\varepsilon_{x,y} = 5 \cdot 10^{-10} \text{ rad m} \quad n_b = 2808$$

$$\sigma_{x,y} = 17 \text{ } \mu\text{m}$$

$$I_p = 584 \text{ mA}$$

$$L = \frac{1}{4\pi e^2 f_0 n_b} * \frac{I_{p1} I_{p2}}{\sigma_x \sigma_y}$$


$$L = 1.0 \cdot 10^{34} \text{ } 1/\text{cm}^2 \text{ s}$$

Mini- β Insertions: some guide lines

- * calculate the periodic solution in the arc
- * introduce the drift space needed for the insertion device (detector ...)
- * put a quadrupole doublet (triplet ?) as close as possible
- * introduce additional quadrupole lenses to match the beam parameters to the values at the beginning of the arc structure

parameters to be optimised & matched to the periodic solution: $\alpha_x, \beta_x, D_x, D'_x$
 $\alpha_y, \beta_y, Q_x, Q_y$

8 individually powered quad magnets are needed to match the insertion (... at least)



Mini- β Insertions: Betafunctions

A mini- β insertion is always a kind of special symmetric drift space.


→ greetings from Liouville

$$\alpha^* = 0$$

$$\gamma^* = \frac{1 + \alpha^2}{\beta} = \frac{1}{\beta^*}$$



$$\sigma'^* = \sqrt{\frac{\varepsilon}{\beta^*}}$$

$$\beta^* = \frac{\sigma^*}{\sigma'^*}$$




at a symmetry point β is just the ratio of beam dimension and beam divergence.

The LHC Insertions




mini β optics



... and now back to the Chromaticity

$$Q' = -\frac{1}{4\pi} \oint k(s) \beta(s) ds$$



Clearly there is another problem ...

... if it were easy everybody could do it



Again: the phase space ellipse

for each turn write down - at a given

position „s” in the ring - the

single particle amplitude x

and the angle x' ... and plot it. $\begin{pmatrix} x \\ x' \end{pmatrix}_{s1} = M_{turn} * \begin{pmatrix} x \\ x' \end{pmatrix}_{s0}$




A beam of 4 particles

- each having a slightly different emittance:

25.) Particle Tracking Calculations

particle vector:

$$\begin{pmatrix} x \\ x' \end{pmatrix}$$



Idea: calculate the particle coordinates x, x' through the linear lattice ... using the matrix formalism.
if you encounter a nonlinear element (e.g. sextupole): stop
calculate explicitly the magnetic field at the particles coordinate

$$B = \left(\begin{array}{c} g'xz \\ \frac{1}{2} g' (x^2 - z^2) \end{array} \right)$$

calculate kick on the particle

$$\Delta x'_1 = \frac{B_x l}{p/e} = \frac{1}{2} \frac{g'}{p/e} l (x_1^2 - z_1^2) = \frac{1}{2} m_{\text{sext}} l (x_1^2 - z_1^2)$$

$$\begin{pmatrix} x_1 \\ x'_1 \end{pmatrix} \rightarrow \begin{pmatrix} x_1 \\ x'_1 + \Delta x'_1 \end{pmatrix}$$



$$\Delta z'_1 = \frac{B_z l}{p/e} = \frac{g' x_1 z_1}{p/e} l = m_{\text{sext}} l x_1 z_1$$

$$\begin{pmatrix} z_1 \\ z'_1 \end{pmatrix} \rightarrow \begin{pmatrix} z_1 \\ z'_1 + \Delta z'_1 \end{pmatrix}$$

and continue with the linear matrix transformations



Installation of a weak (!!!) sextupole magnet

The good news: sextupole fields in accelerators cannot be treated with conventional methods.
→ no equations; instead: Computer simulation
„ particle tracking “



Effect of a strong (!!!) Sextupole ...

→ **Catastrophy !**



„dynamic aperture“

Resume':

quadrupole error: tune shift

$$\Delta Q \approx \int_{s_0}^{s_0+l} \frac{\Delta k(s) \beta(s)}{4\pi} ds \approx \frac{\Delta k(s) l_{quad} \bar{\beta}}{4\pi}$$

beta beat

$$\Delta \beta(s_0) = \frac{\beta_0}{2 \sin 2\pi Q} \int_{s_1}^{s_1+l} \beta(s_1) \Delta k \cos(2(\psi_{s_1} - \psi_{s_0}) - 2\pi Q) ds$$

chromaticity

$$\Delta Q = Q' \frac{\Delta p}{p}$$

$$Q' = -\frac{1}{4\pi} \oint k(s) \beta(s) ds$$

momentum compaction

$$\frac{\delta l_e}{L} = \alpha_p \frac{\Delta p}{p}$$

$$\alpha_p \approx \frac{2\pi}{L} \langle D \rangle \approx \frac{\langle D \rangle}{R}$$

beta function in a symmetric drift

$$\beta(s) = \beta_0 + \frac{s^2}{\beta_0}$$

Appendix I:

Dispersion: Solution of the inhomogeneous equation of motion

Ansatz:
$$D(s) = S(s) \int_{s_0}^{s1} \frac{1}{\rho} C(\tilde{s}) d\tilde{s} - C(s) \int_{s_0}^{s1} \frac{1}{\rho} S(\tilde{s}) d\tilde{s}$$

$$D'(s) = S' * \int \frac{1}{\rho} C dt + S \frac{1}{\rho} C - C' * \int \frac{1}{\rho} S dt - C \frac{1}{\rho} S$$

$$D'(s) = S' * \int \frac{C}{\rho} dt - C' * \int \frac{S}{\rho} dt$$

$$\begin{aligned} D''(s) &= S'' * \int \frac{C}{\rho} d\tilde{s} + S' \frac{C}{\rho} - C'' * \int \frac{S}{\rho} d\tilde{s} - C' \frac{S}{\rho} \\ &= S'' * \int \frac{C}{\rho} d\tilde{s} - C'' * \int \frac{S}{\rho} d\tilde{s} + \frac{1}{\rho} (CS' - S C') \\ & \qquad \qquad \qquad = \det M = 1 \end{aligned}$$

remember: for Cs and S(s) to be independent solutions the Wronski determinant has to meet the condition

$$W = \begin{vmatrix} C & S \\ C' & S' \end{vmatrix} \neq 0$$

and as it is independent of the variable „s“

$$\frac{dW}{ds} = \frac{d}{ds} (CS' - SC') = CS'' - SC'' = -K(CS - SC) = 0$$

we get for the initial conditions that we had chosen ...

$$\left. \begin{array}{l} C_0 = 1, \quad C'_0 = 0 \\ S_0 = 0, \quad S'_0 = 1 \end{array} \right\} W = \begin{vmatrix} C & S \\ C' & S' \end{vmatrix} = 1$$

$$D'' = S'' * \int \frac{C}{\rho} d\tilde{s} - C'' * \int \frac{S}{\rho} d\tilde{s} + \frac{1}{\rho}$$

remember: S & C are solutions of the homog. equation of motion:

$$S'' + K * S = 0$$

$$C'' + K * C = 0$$

$$D'' = -K * S * \int \frac{C}{\rho} d\tilde{s} + K * C * \int \frac{S}{\rho} d\tilde{s} + \frac{1}{\rho}$$

$$D'' = -K * \left\{ S \int \frac{C}{\rho} d\tilde{s} + C \int \frac{S}{\rho} d\tilde{s} \right\} + \frac{1}{\rho}$$

= D(s)

$$D'' = -K * D + \frac{1}{\rho}$$

... or

$$\underline{\underline{D'' + K * D = \frac{1}{\rho}}}$$


qed

Appendix II:


Quadrupole Error and Beta Function

a change of quadrupole strength in a synchrotron leads to tune shift:

$$\Delta Q \approx \int_{s_0}^{s_0+l} \frac{\Delta k(s) \beta(s)}{4\pi} ds \approx \frac{\Delta k(s) * l_{quad} * \bar{\beta}}{4\pi}$$



tune spectrum ...



tune shift as a function of a gradient change

But we should expect an error in the β -function as well ...
... shouldn't we ???

Quadrupole Errors and Beta Function


a quadrupole error will not only influence the oscillation frequency ... „tune“
... but also the amplitude ... „beta function“

split the ring into 2 parts, described by two matrices
A and B

$$M_{turn} = B * A$$

$$A = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$$

$$B = \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix}$$



matrix of a quad error
between A and B

$$M_{dist} = \begin{pmatrix} m_{11}^* & m_{12}^* \\ m_{21}^* & m_{22}^* \end{pmatrix} = B \begin{pmatrix} 1 & 0 \\ -\Delta k ds & 1 \end{pmatrix} A$$

$$M_{dist} = B \begin{pmatrix} a_{11} & a_{12} \\ -\Delta k ds a_{11} + a_{12} & -\Delta k ds a_{12} + a_{22} \end{pmatrix}$$

$$M_{dist} = \begin{pmatrix} \sim & b_{11} a_{12} + b_{12} (-\Delta k ds a_{12} + a_{22}) \\ \sim & \sim \end{pmatrix}$$

the beta function is usually obtained via the matrix element „m12“, which is in Twiss form for the undistorted case

$$m_{12} = \beta_0 \sin 2\pi Q$$

and including the error:

$$m_{12}^* = b_{11}a_{12} + b_{12}a_{22} - b_{12}a_{12}\Delta kds$$

$$\underbrace{\hspace{10em}}_{m_{12} = \beta_0 \sin 2\pi Q}$$

$$(1) \quad m_{12}^* = \beta_0 \sin 2\pi Q - a_{12}b_{12}\Delta kds$$

As M^* is still a matrix for one complete turn we still can express the element m_{12} in twiss form:

$$(2) \quad m_{12}^* = (\beta_0 + d\beta) \sin 2\pi(Q + dQ)$$

Equalising (1) and (2) and assuming a small error

$$\beta_0 \sin 2\pi Q - a_{12}b_{12}\Delta kds = (\beta_0 + d\beta) \sin 2\pi(Q + dQ)$$

$$\beta_0 \sin 2\pi Q - a_{12}b_{12}\Delta kds = (\beta_0 + d\beta) \sin 2\pi Q \underbrace{\cos 2\pi dQ}_{\approx 1} + \cos 2\pi Q \underbrace{\sin 2\pi dQ}_{\approx 2\pi dQ}$$

~~$$\beta_0 \sin 2\pi Q - a_{12}b_{12}\Delta kds = \beta_0 \sin 2\pi Q + \beta_0 2\pi dQ \cos 2\pi Q + d\beta_0 \sin 2\pi Q + d\beta_0 2\pi dQ \cos 2\pi Q$$~~

ignoring second order terms

$$-a_{12}b_{12}\Delta kds = \beta_0 2\pi dQ \cos 2\pi Q + d\beta_0 \sin 2\pi Q$$

remember: tune shift dQ due to quadrupole error: $dQ = \frac{\Delta k \beta_1 ds}{4\pi}$
(index „1“ refers to location of the error)

$$-a_{12}b_{12}\Delta kds = \frac{\beta_0 \Delta k \beta_1 ds}{2} \cos 2\pi Q + d\beta_0 \sin 2\pi Q$$

solve for $d\beta$

$$d\beta_0 = \frac{-1}{2 \sin 2\pi Q} \{2a_{12}b_{12} + \beta_0 \beta_1 \cos 2\pi Q\} \Delta kds$$

express the matrix elements a_{12} , b_{12} in Twiss form

$$M = \begin{pmatrix} \sqrt{\frac{\beta_s}{\beta_0}} (\cos \psi_s + \alpha_0 \sin \psi_s) & \sqrt{\beta_s \beta_0} \sin \psi_s \\ \frac{(\alpha_0 - \alpha_s) \cos \psi_s - (1 + \alpha_0 \alpha_s) \sin \psi_s}{\sqrt{\beta_s \beta_0}} & \sqrt{\frac{\beta_0}{\beta_s}} (\cos \psi_s - \alpha_s \sin \psi_s) \end{pmatrix}$$

$$d\beta_0 = \frac{-1}{2 \sin 2\pi Q} \{2a_{12}b_{12} + \beta_0\beta_1 \cos 2\pi Q\} \Delta k ds$$

$$a_{12} = \sqrt{\beta_0\beta_1} \sin \Delta\psi_{0 \rightarrow 1}$$

$$b_{12} = \sqrt{\beta_1\beta_0} \sin(2\pi Q - \Delta\psi_{0 \rightarrow 1})$$

$$d\beta_0 = \frac{-\beta_0\beta_1}{2 \sin 2\pi Q} \underbrace{\{2 \sin \Delta\psi_{12} \sin(2\pi Q - \Delta\psi_{12}) + \cos 2\pi Q\}}_{\dots}$$

... after some TLC transformations ... = $\cos(2\Delta\psi_{01} - 2\pi Q)$

$$\Delta\beta(s_0) = \frac{-\beta_0}{2 \sin 2\pi Q} \int_{s_1}^{s_1+l} \beta(s_1) \Delta k \cos(2(\psi_{s_1} - \psi_{s_0}) - 2\pi Q) ds$$

Nota bene: ! the beta beat is proportional to the strength of the error Δk

!! and to the β function at the place of the error ,

!!! and to the β function at the observation point,
(... remember orbit distortion !!!)

!!!! there is a resonance denominator