



High Precision Measurement of Transversity using Di-hadron Correlations in $p^\uparrow + p$ Collisions at $\sqrt{s} = 500$ GeV at STAR

Michael Skoby

For the STAR Collaboration

Center for Exploration of Energy and Matter




INDIANA UNIVERSITY

Overview

- Why measure $\pi^+\pi^-$ correlations?
- Some analysis details
- Asymmetry measurements vs η , p_T and M_{Inv}
- Conclusions

Motivation


Bacchetta, Courtoy,
Radici, JHEP **1303**
(2013) 119



- Di-hadron correlations allow point-to-point transversity measurements in SIDIS
- High precision data lacking at relatively high x
- Measuring transversity from polarized p+p data
 - collinear framework
 - high precision, reduced u-quark dominance
 - test of universality (SIDIS vs p+p)
 - new kinematic regime


STAR

- 2011 polarized p+p collisions at 500 GeV with 25 pb^{-1} integrated luminosity
- $P_{\text{beam}} = 53\%$
- Solenoidal Tracker at RHIC (STAR)
- Charged pions measured in Time Projection Chamber
 - 2π azimuthal coverage
 - $-1 < \eta < 1$
- Endcap and Barrel electromagnetic calorimeters and vertex position detector used to select events



Charged Pion Purity Estimates

p_T range (GeV/c)	Pion purity
1.5 – 2.0	0.97
2.0 – 3.0	0.94
3.0 – 4.0	0.88
4.0 – 6.0	0.83
6.0 – 8.0	0.86
> 8.0	0.97





- Use dE/dx to identify pions

- $n\sigma(\pi) \approx \# \text{ of } \sigma \text{ in } z = \ln\left(\frac{dE/dx_{\text{measured}}}{dE/dx_{\text{parameterized}}}\right)$ distribution


- Excellent pion purity samples

Asymmetry Observable

- Calculated for \vec{P}_B as incident beam, \vec{P}_A as target
- Incident beam is polarized and target unpolarized by summing over bunches
- Pion separation = $\sqrt{(\Delta\eta^2 + \Delta\phi^2)} < 0.7$
- $A_{UT} \propto h_1 \cdot H_1^<$
 - Transversity (h_1)
 - Interference Fragmentation Function ($H_1^<$)
- A_{UT} is expected to depend on the invariant mass (M_{Inv}) and p_T of the pion pair



Extract A_{UT}




- Particle $p_T > 1.5$ GeV/c
- Pair $p_T > 3.75$ GeV/c
- For a given M_{Inv} , p_T bin the asymmetry is calculated for 8 ϕ_{RS} bins
- The asymmetry is the amplitude extracted from a single-parameter fit
- Example shown here is one M_{Inv} , p_T bin

$$A_{UT}(\varphi_{RS}) = \frac{1}{P} \frac{\sqrt{N \uparrow(\varphi_{RS}) N \downarrow(\varphi_{RS} + \pi)} - \sqrt{N \downarrow(\varphi_{RS}) N \uparrow(\varphi_{RS} + \pi)}}{\sqrt{N \uparrow(\varphi_{RS}) N \downarrow(\varphi_{RS} + \pi)} + \sqrt{N \downarrow(\varphi_{RS}) N \uparrow(\varphi_{RS} + \pi)}}$$


Asymmetry (η, p_T)

- A_{UT} as a function of η plotted for 5 p_T bins
- Significant asymmetry seen at high η and high $\langle p_T \rangle$




Asymmetry (η, p_T)

- A_{UT} as a function of η plotted for 5 p_T bins
- Significant asymmetry seen at high η and high $\langle p_T \rangle$




Asymmetry (M_{Inv}, p_T)

- A_{UT} as a function of M_{Inv} plotted for 5 p_T bins
- Avg M_{Inv} in each M_{Inv} bin decreases with decreasing $\langle p_T \rangle$
- Significant asymmetry seen at mid- M_{Inv} and high $\langle p_T \rangle$




Asymmetry (M_{Inv}, p_T)

- A_{UT} as a function of M_{Inv} plotted for 5 p_T bins
- Avg M_{Inv} in each M_{Inv} bin decreases with decreasing $\langle p_T \rangle$
- Significant asymmetry seen at mid- M_{Inv} and high $\langle p_T \rangle$




Asymmetry (p_T, M_{Inv})

- A_{UT} as a function of p_T plotted for 5 M_{Inv} bins
- Avg p_T in each p_T bin slightly decreases with decreasing $\langle M_{Inv} \rangle$
- Asymmetry rises significantly for high p_T and high M_{Inv}





Asymmetry (p_T, M_{Inv})

- A_{UT} as a function of p_T plotted for 5 M_{Inv} bins
- Avg p_T in each p_T bin slightly decreases with decreasing $\langle M_{Inv} \rangle$
- Asymmetry rises significantly for high p_T and high M_{Inv}




Measurement Bias





- The events we choose to record are biased towards pions that fragment from quarks
- There should be no asymmetry for pion pairs that come from gluons
- To account for the bias a dilution correction is estimated in the top panel
 - Quarks/partons ratio of biased data over the quarks/partons ratio of unbiased sample
- **Correction not applied to data**

Results for $\eta^{\pi^+\pi^-} < 0$



$\langle x \rangle$ Coverage at STAR



Bacchetta, Courtoy, Radici, JHEP **1303** (2013) 119

- High precision asymmetries measured at relatively high $\langle x \rangle$ and high effective Q^2

Conclusions

- Preliminary STAR data show high precision pion pair correlation asymmetries at large p_T and M_{Inv} for $\eta^{\pi^+\pi^-} > 0$
- These results are at much higher Q^2 and sample a different mixture of quark flavors than SIDIS
- Results may be used to test universality of transverse polarization dependent quantities (SIDIS vs p+p)
- STAR results from 2012 polarized p+p collisions at $\sqrt{s} = 200$ GeV coming soon (higher precision than 2006)