

Lattice calculation of hadronic light-by-light scattering contribution to the muon $g-2$

Tom Blum (UConn / RIKEN BNL Research Center)

Confinement XI, St. Petersburg, September 9, 2014

Based on arXiv:1407.2923, TB, Saumitra Chowdhury, Masashi Hayakawa, and Taku Izubuchi

For HVP see Mainz work shop mini-proceedings arXiv:1407.4021 and Lattice 2014 talks

Work on $g-2$ done in collaboration with

	HVP	HLbL
Christopher Aubin (Fordham U)		Saumitra Chowdhury (UConn)
Maarten Golterman (SFSU)		Norman Christ (Columbia)
Santiago Peris (Barcelona)		Masashi Hayakawa (Nagoya)
		Taku Izubuchi (BNL/RBRC)
RBC/UKQCD Collaboration		Luchang Jin (Columbia)
		Christoph Lehner (BNL)
		Norikazu Yamada (KEK)

The magnetic moment of the muon

Interaction of particle with static magnetic field

$$V(\vec{x}) = -\vec{\mu} \cdot \vec{B}_{\text{ext}}$$

The magnetic moment $\vec{\mu}$ is proportional to its spin ($c = \hbar = 1$)

$$\vec{\mu} = g \left(\frac{e}{2m} \right) \vec{S}$$


The Landé ***g*-factor** is predicted from the **free Dirac eq.** to be

$$g = 2$$

for elementary fermions

The magnetic moment of the muon

In interacting **quantum** (field) theory g gets corrections



$$\gamma^\mu \rightarrow \Gamma^\mu(q) = \left(\gamma^\mu F_1(q^2) + \frac{i \sigma^{\mu\nu} q_\nu}{2m} F_2(q^2) \right)$$


which results from Lorentz and gauge invariance when the muon is on-mass-shell.

$$F_2(0) = \frac{g-2}{2} \equiv a_\mu \quad (F_1(0) = 1)$$

(the anomalous magnetic moment, or anomaly)

The magnetic moment of the muon

Compute corrections to $g-2$ in pert. theory in $\alpha = \frac{e^2}{4\pi} = \frac{1}{137} + \dots$



(leading) Schwinger term = $\frac{\alpha}{2\pi} = 0.0011614\dots$

hadronic contributions $\sim 6 \times 10^{-5}$, 10^{-6} times smaller


dominate error, ~ 0.4 ppm (exp 0.54 ppm)

QED		116 584 71.8 951 (9)(19)(7)(77)	Aoyama (2012)
EW		15.4 (2)	Czarnecki (2002)
QCD	LO HVP	692.3 (4.2)	Davier (2010)
		694.91 (3.72) (2.10)	Hagiwara (2011)
	NLO HVP	701.5 (4.7)	Davier (2010)
		-9.79 (9)	Hagiwara (2006), Kurz (2014)
	HLbL	10.5 (2.6)	Prades (2009)
	NNLO HVP	1.24 (1)	Kurz (2014)

New experiments + new theory = new physics?

- Fermilab E989 (~ 2 years away) and J-PARC E34: 0.14 ppm
- $a_\mu(\text{Expt}) - a_\mu(\text{SM}) = 287(63)(51) (\times 10^{-11})$, or $\sim 3.6\sigma$ (or 2.9)
- If both central values stay the same,
 - E989 ($\sim 4\times$ smaller error) $\rightarrow \sim 5\sigma$
 - E989+new HLBL theory (models+lattice, 10%) $\rightarrow \sim 6\sigma$
 - E989+new HLBL +new HVP (50% reduction) $\rightarrow \sim 8\sigma$
- **Big discrepancy!** (New Physics $\sim 2\times$ Electroweak)
- Lattice calculations crucial
- a_μ good for constraining and explaining BSM physics

The hadronic light-by-light amplitude



Model estimates: about $(10-12) \times 10^{-10}$ with a 25-40% uncertainty (difficult to quantify)


Lattice calculation: model independent, approximations (non-zero a , finite V , ...) systematically improvable

Compute directly on lattice, using **QCD and QED**

Lattice field theory calculation reminder

- Compute QFT path integrals numerically and stochastically
 - Fields live on **finite, discrete**, 4d (Euclidean) space-time lattice
 - Generate ensemble of field configurations using monte carlo
 - Average over configurations
- Typically **compute correlation function** of fields, extract (Minkowski) matrix element or amplitude
- Computation dominated by quark propagators, **inverse of large, sparse matrix**.
- Extrapolate to continuum, infinite volume, physical quark masses (now directly accessible)

Lattice QCD: conventional approach (*c.f.*, HVP)




Correlation of 4 EM currents
 $\Pi^{\mu\nu\rho\sigma}(q, p_1, p_2)$

Two independent momenta
+ external mom q

Compute for all possible
values of p_1 and p_2 ($O(V^2)$)
four index tensor

several q (extrap $q \rightarrow 0$),
fit, plug into perturbative QED
two-loop integrals

Alternate approach: Lattice QCD+QED



Average over combined gluon *and* photon gauge configurations

Quarks coupled to gluons and photons


muon coupled to photons

[Hayakawa, *et al.* hep-lat/0509016;

Chowdhury *et al.* (2008);

Chowdhury Ph. D. thesis (2009)]

Alternate approach: Lattice QCD+QED



Attach one photon by hand
(see why in a minute)

Correlation of hadronic loop
and muon line


[Hayakawa, *et al.* hep-lat/0509016;

Chowdhury *et al.* (2008);


Chowdhury Ph. D. thesis (2009)]

Formally expand in α electromagnetic

The leading and next-to-leading contributions in α to magnetic part of correlation function come from



Subtraction of lowest order piece




Subtraction term is product of separate averages of the loop and line

Gauge configurations identical in both, so two are **highly correlated**

In PT, correlation function and subtraction have **same contributions except the light-by-light** term which is absent in the subtraction

Subtraction of lowest order piece: two photons?




- absent in subtraction term, but vanishes due to **Furry's theorem**
- Only after averaging over gauge fields, potentially large error ($O(\alpha^2)$ compared to signal of $O(\alpha^3)$)
- **Exact symmetry** under $\mathbf{p} \rightarrow -\mathbf{p}$
 $e \rightarrow -e$ on muon line only
- If e unchanged, only effect is to flip the sign of all diagrams with two photons, so these cancel on each configuration.
- Observe large reductions in statistical errors after \pm momentum averaging

LbL contribution from lattice QED – a test

- LbL calculation: quenched (no vac. pol.), non-compact QED
 - $m_{\text{lepton}} = 0.1$, loop and line
 - 16^3 and $24^3 \times 64 (\times 8)$ Domain Wall Fermions
 - $(L/4)^3 = 64$ and 216 propagators for the lepton loop to enhance statistics
 - incoming muon at rest, $\vec{p} = \pm(2\pi/L, 0, 0)^T$, and permutations, at external vertex
 - several source/sink separations for muon (8-32) to project on to ground states

LbL contribution from lattice QED – a test



Blum, Chowdhury, Hayakawa, and Izubuchi (arXiv:1407.2923)

- lowest non-trivial momentum only
- Stat. errors only
- Several source/sink separations (t_{sep}) for muon (loop is same, only line differs)
- Significant excited state contamination
- $m_{\mu} = 0.1$ loop and line
- Consistent with expectations from continuum PT


HLbL contribution from lattice QCD+QED

Calculation is almost the same, just take

$$U_\mu(x) = U_\mu^{QED}(x)U_\mu(x)^{QCD} \text{ for the combined gauge field}$$

- HLbL calculation: RBC/UKQCD 2+1f DWF ensemble
 - $m_\pi = 329$ MeV (light quarks heavier than u,d; s is physical)
 - lattice size $24^3 \times 64(\times 16)$ DWF
 - lattice spacing $a = 0.114$ fm
 - quenched QED for now (sea quarks not charged)
 - $(L/4)^3 = 216$ propagators for the quark loop/configuration.
 - All mode averaging (AMA) technique crucial to make computation tractable


HLbL contribution from lattice QCD+QED



Blum, Chowdhury, Hayakawa, and Izubuchi (arXiv:1407.2923)

- Stat. errors only, lowest non-trivial momentum
- Several source/sink separations (t_{sep}) for muon
- Significant excited state contamination
- $m_\pi = 329$ MeV
- Model value/error is “Glasgow Consensus”
(arXiv:0901.0306 [hep-ph], physical masses)
- Subtracted result shows correct e^4 behavior (✓)

Momentum dependence



- $t_{\text{sep}} = 10$
- Stat. errors only, lowest non-trivial momentum
- $m_\pi = 329$ MeV
- Model value/error is “Glasgow Consensus”

(arXiv:0901.0306 [hep-ph], physical masses)

Blum, Chowdhury, Hayakawa, and Izubuchi (arXiv:1407.2923)

Simplest model: π^0 exchange + charged pion loop.


Roughly, $\sim m_\mu^2$, $\sim 1/m_\pi^2$, $\sim m_\rho$ (cutoff in form factor)

- m_{π^0} 139.57 MeV, m_{π^\pm} 134.98 MeV, $m_\mu = 105$ MeV,
 $m_\rho = 770$ MeV
5.65-1.63=4.02 ($\times 10^{-10}$) (less than 50% of total)
- m_π 330 MeV, $m_\mu = 200$ MeV, $m_\rho = 900$ MeV
8.35-0.51=7.84 ($\times 10^{-10}$)
- the rest comes mostly from η, η' exchange

Lattice value (lowest q^2 and un-physical masses) is roughly the total model value, evaluated at physical masses, 10.5.

Momentum extrapolation to zero probably increases this

“Disconnected” diagrams






(and similar) **not calculated yet**

Omission due to use of quenched QED, i.e., sea quarks not electrically charged. Two possibilities,

- 1 Re-weight in α T. Ishikawa, *et al.*, Phys.Rev.Lett. 109 (2012) 072002 or
- 2 dynamical QED(+QCD) in HMC BMWc arXiv:1406.4088, QCDSF

Use same non-perturbative method as for quenched QED

Disconnected quark loop diagrams



Disconnected quark loop diagrams in non-pert. method

One additional correlation function (+dynamical QED)

$$\mathcal{M}_C = \left\langle \text{Diagram} \right\rangle_{\text{QCD+f-QED}} \quad \mathcal{S}_C = \left\langle \text{Diagram} \right\rangle_{\text{QCD+f-QED}} \left\langle \text{Diagram} \right\rangle_{\text{f-QED}}$$

The diagram for \mathcal{M}_C shows a quark loop with a gluon self-energy correction (a gluon loop) on the bottom quark line. The diagram for \mathcal{S}_C shows a quark loop with a photon self-energy correction (a fermion loop) on the bottom quark line.

$$\mathcal{M}_D = \left\langle \text{Diagram} \right\rangle_{\text{QCD+f-QED}} \quad \mathcal{S}_D = \left\langle \text{Diagram} \right\rangle_{\text{QCD+f-QED}} \left\langle \text{Diagram} \right\rangle_{\text{f-QED}}$$

The diagram for \mathcal{M}_D shows two quark loops connected by a gluon line. The diagram for \mathcal{S}_D shows two quark loops connected by a photon line.

Need to address


- quark and muon masses (model results: significant dep.)
- quenched QED (reweight QCD ensemble, or dynamical QED)
- Finite volume
- continuum limit $a \rightarrow 0$
- $q^2 \rightarrow 0$ exptrap
- QED renormalization
- excited states/“around the world” effects
- ...

Computationally demanding, but straightforward

Alternative non-pert. method (Luchang Jin)

see talk at Lattice 2014

No subtraction, use 2 independent stochastic photons, one exact (similar to Rome123)



No clear advantage so far, but good check

Summary and Outlook

- First lattice QCD calculation of HLbL contribution to $g-2$. Promising, method is feasible with current resources.
- Checked QCD+QED method in pure QED (✓)
- Crucial to leverage FNAL E989, J-PARC E34 experiments, search for new physics
- Next HLbL calculations:
 - ① RBC/UKQCD 2+1f DWF ensemble
 $m_\pi = 170 \text{ MeV}, 32^3 \times 64 \times 32$
 - ② 2+1f DWF dynamical or reweighted QCD+QED
 $m_\pi = 300 \text{ MeV}, 24^3 \times 64$
 - ③ RBC/UKQCD 2+1f Möbius-DWF
 $m_\pi = 140 \text{ MeV}, 48^3 \times 96 \times 24$

Acknowledgments

- This research is supported in part by the US DOE
- Computational resources provided by the RIKEN BNL Research Center and USQCD Collaboration
- Lattice computations done on
 - Ds cluster at FNAL
 - q-series clusters at JLab
 - QCDOC at BNL