

J/ψ and ψ(2S) measurements in p+p collisions at $\sqrt{s} = 200$ and 500 GeV in the STAR experiment

Barbara Trzeciak

for the STAR Collaboration

*Faculty of Nuclear Sciences and Physical Engineering
Czech Technical University in Prague*



*Hot Quarks 2014
September 21-28, 2014
Les Negras, Andalucia, Spain*



INVESTICE DO ROZVOJE VZDĚLÁVÁNÍ




Charmonia in p+p collisions



Charmonia - $c\bar{c}$ bound states

$J/\psi, \psi(2S), \chi_c$




Charmonia in $p+p$ collisions



Charmonia - $c\bar{c}$ bound states

$J/\psi, \psi(2S), \chi_c$



- **Production mechanism** in elementary collisions is not fully understood

→ Color singlet vs color octet

- Color-singlet process: J/ψ is produced via intermediate color-neutral $c\bar{c}$ state with the same quantum numbers as the final J/ψ state
- Color-octet process: J/ψ is produced via intermediate colored $c\bar{c}$ state of any possible quantum numbers


→ **Quarkonium measurements - tests of different production models, help to understand QCD**

Charmonia in $p+p$ collisions



Charmonia - $c\bar{c}$ bound states

$J/\psi, \psi(2S), \chi_c$



Feed-down

Inclusive J/ψ production:


- ▶ prompt J/ψ
- ▶ direct J/ψ ($\sim 60\%$), feed down from $\psi(2S)$ ($\sim 10\%$) and χ_c ($\sim 30\%$) decays
- ▶ non-prompt J/ψ : B-mesons feed-down ($10\text{-}25\%$ at 4-12 GeV/c, STAR: Phys. Lett. B722 (2013) 55)

STAR Experiment at RHIC



$J/\psi, \psi(2S) \rightarrow e^+ e^-$

Large acceptance: $|\eta| < 1, 0 < \phi < 2\pi$




- ✓ VPD - minimum bias trigger
- ✓ TPC
- ✓ TOF
- ✓ BEMC

STAR Experiment at RHIC



$J/\psi, \psi(2S) \rightarrow e^+ e^-$

Large acceptance: $|\eta| < 1, 0 < \phi < 2\pi$




- ✓ VPD - minimum bias trigger
- ✓ TPC - tracking, **PID**: dE/dx
- ✓ TOF
- ✓ BEMC

STAR Experiment at RHIC




$J/\psi, \psi(2S) \rightarrow e^+ e^-$

Large acceptance: $|\eta| < 1, 0 < \phi < 2\pi$




- ✓ VPD - minimum bias trigger
- ✓ TPC - tracking, **PID**: dE/dx
- ✓ TOF - time resolution < 100 ps **PID**: $1/\beta$
- ✓ BEMC

STAR Experiment at RHIC




$J/\psi, \psi(2S) \rightarrow e^+ e^-$

Large acceptance: $|\eta| < 1, 0 < \phi < 2\pi$




- ✓ VPD - minimum bias trigger
- ✓ TPC - tracking, **PID**: dE/dx
- ✓ TOF - time resolution < 100 ps **PID**: $1/\beta$
- ✓ BEMC

STAR Experiment at RHIC



$J/\psi, \psi(2S) \rightarrow e^+ e^-$



Large acceptance: $|\eta| < 1, 0 < \phi < 2\pi$





- ✓ VPD - minimum bias trigger
- ✓ TPC - tracking, **PID**: dE/dx
- ✓ TOF - time resolution $< 100 \text{ ps}$ **PID**: $1/\beta$
- ✓ BEMC - trigger, **PID**: E/p (~ 1 for electron)

Excellent eID

J/Ψ and $\Psi(2S)$ signals



$p+p 200 \text{ GeV}$




$p+p 500 \text{ GeV}$

J/Ψ p_T spectrum in $p+p$ 200 GeV



- Test of charmonium production models


STAR 2005&2006: Phys. Rev. C80, 041902(R) (2009)
 PHENIX: Phys. Rev. D 85, 092004 (2012)
 direct NNLO CS: P.Artoisenet et al., Phys. Rev. Lett. 101, 152001 (2008) and J.P.Lansberg private communication
 NLO CS+CO: Y.-Q.Ma, K.Wang, and K.T.Chao, Phys. Rev. D 84, 51 114001 (2011) and private communication
 CEM: A.D.Frawley, T.Ullrich, R.Vogt, Phys. Rept. 462 (2008) 125, and R.Vogt private communication





STAR EMC : Phys. Lett. B 722 (2013) 55

STAR MB: Acta Phys. Polonica B Vol.5, No 2 (2012), 543

- ✓ *prompt NLO CS+CO* model describes the data for $p_T > 4$ GeV/c
- ✓ *direct NNLO* CS* model misses high- p_T part
- ✓ *prompt CEM* model can reasonably well describe the p_T spectra (overpredicts the data at $p_T \sim 3$ GeV/c)




J/ψ p_T spectrum in p+p 500 GeV



- ✓ Precise J/ψ measurement at new beam energy, up to $p_T = 20 \text{ GeV}/c$

J/Ψ x_T scaling



$$x_T = 2p_T / \sqrt{s}$$

$$\frac{d^2\sigma}{2\pi p_T dp_T dy} = g(x_T) / (\sqrt{s})^n$$

✓ $p_T > 5 \text{ GeV}/c$ - J/ψ production follows the x_T scaling of cross-section at mid-rapidity, with $n = 5.6$ (Phys. Rev. C 80, 041902 (2009))


→ x_T scaling breaking - transition from hard to soft process

n - number of constituents taking an active role in hadron production

$\Psi(2S)$ in $p+p$ 500 GeV



- Further test of charmonium production models
- Constrain $\Psi(2S)$ feed-down contribution to inclusive J/ψ production





- ✓ First measurement of $(\Psi(2S) / J/\psi)$ ratio in $p+p$ at 500 GeV
 - Consistent with other experiments
 - No collision energy dependence observed

$B \rightarrow J/\Psi$ fraction in $p+p$ 200 GeV




- ✓ Measurement based on azimuthal angular correlations between high- p_T J/ψ and charged hadrons
 - B-hadron feed-down contribution: 10-25%, in the range $4 < p_T < 12$ GeV/c
 - Agreement with FONLL + CEM prediction



$B \rightarrow J/\Psi$ fraction in $p+p$ 200 GeV




- ✓ Measurement based on azimuthal angular correlation between high- p_T J/ψ and charged hadrons



Phys. Rev. C 80, 041902 (2009)

→ B-hadron feed-down contribution: 10-25%, in the range $4 < p_T < 12$ GeV/c

→ Agreement with FONLL + CEM prediction and with measurements from other experiments



Phys. Lett. B 722 (2013) 55


J/ψ polarization



- Discrimination power between different J/ψ production models
 - Competing mechanisms domination in different theoretical approaches lead to different expected polarization

J/ψ polarization can be analyzed via the angular distribution of the decay lepton pair

$$\frac{d\sigma}{d(\cos\theta)d\phi} \propto 1 + \lambda_\theta \cos^2\theta + \lambda_{\theta\phi} \sin(2\theta)\cos\phi + \lambda_\phi \sin^2\theta\cos(2\phi)$$



- ✓ θ - polar angle between momentum of a positive lepton in the J/ψ rest frame and the polarization axis z
- ✓ ϕ - corresponding azimuthal angle


In the **helicity frame** z axis is defined along the J/ψ momentum in the center of mass frame

J/Ψ polarization




- Discrimination power between different J/Ψ production models
 - Competing mechanisms domination in different theoretical approaches lead to different expected polarization

$J=1 \rightarrow$ three J_z eigenstates $| 1, +1 \rangle, | 1, 0 \rangle, | 1, -1 \rangle$




P. Faccioli, C. Laorenco, J. Seixas, H.K. Wohri, Eur. Phys. J. C 69, 657 (2010)

J/ Ψ polarization in p+p 200 GeV



PHENIX: Phys. Rev. D 82, 012001 (2010)
 COM: Phys. Rev. D 81, 014020 (2010)
 CSM NLO⁺: Phys. Lett. B, 695, 149 (2011)
 and private communication



The angular distribution integrated over the azimuthal angle:

$$W(\cos\theta) \propto 1 + \lambda_\theta \cos^2\theta$$

λ_θ – polarization parameter

$\lambda_\theta = -1$ – longitudinal polarization



$\lambda_\theta = 1$ – transverse polarization

- ✓ Polarization parameter λ_θ is measured in the helicity frame at $|y| < 1$ and $2 < p_T < 6$ GeV/c
 - RHIC data indicate trend towards longitudinal polarization with increasing p_T
 - The result is consistent with NLO⁺ CSM


J/ψ polarization in p+p 500 GeV



- Information about full decay angular distribution
 - ✓ Measurement in progress, larger statistics $\sim 22 \text{ pb}^{-1}$ vs $\sim 1.8 \text{ pb}^{-1}$



- ✓ Reconstruction of both θ and ϕ angles
- ✓ J/ψ signal up to $p_T \sim 15 \text{ GeV}/c$, can be divided into $\sim 5 p_T$ bins




Muon Telescope Detector (MTD)

Precision quarkonium measurements via di- μ channel


μ advantages over e :

- No γ conversion
- Much less Dalitz decay contribution
- Less affected by radiative loses in the detector material



- ✗ Acceptance: 45% at $|ln\ell| < 0.5$
- ✗ Multi-gap Resistive Plate Chamber (MRPC) - gas detector
- ✗ Long-MRPCs

Heavy Flavor Tracker (HFT)



Inner tracking system with 3 sub-systems

Precise pointing resolution

$$B \rightarrow J/\psi + X$$


- NLO CS+CO and CEM models describe the J/ψ p_T spectrum in $p+p$ 200 GeV
- New J/ψ measurement in $p+p$ 500 GeV – production follows x_T scaling at high p_T
- First $\psi(2S)$ / J/ψ measurement in $p+p$ at 500 GeV – no collision energy dependence observed
- J/ψ polarization in $p+p$ 200 GeV consistent with the NLO⁺ CSM prediction
- Measurement of J/ψ polarization in $p+p$ 500 GeV in progress
 - ➔ *HFT and MTD since 2014 – significant improvement of quarkonium measurements*

Thank you !

J/Ψ production mechanism - CSM




- Comparison of CSM to RHIC data




J.P. Lansberg, Phys.Lett.B 695 (2011) 149

J/ Ψ production mechanism - NRQCD



Each color singlet and octet term has a specific polarization associated



colour-singlet 3S_1

$\lambda_\theta = +1$ at LO to $\lambda_\theta = -1$ at NLO

tiny fraction of the total cross section

octet ${}^1S_0 \rightarrow \lambda_\theta = 0$ at LO, NLO

octet ${}^3S_1 \rightarrow \lambda_\theta = +1$ at LO, NLO, at high p_T

octet ${}^3P_J \rightarrow \lambda_\theta \gg +1$ at NLO at LO it is 0


Dominance of the 3S_1 and 3P_J octet terms

$\rightarrow \lambda_\theta \approx +1$,

for high- p_T S-wave quarkonia


Color-octet contributions have fixed shape but adjustable normalizations (LDMEs)

J/Ψ polarization - observation frames




P. Faccioli, C. Laorenco, J. Seixas, H.K. Wohri, Eur. Phys. J. C 69, 657 (2010)

J/Ψ polarization - frame invariant approach



- Frame invariant quantity:

$$\tilde{\lambda} = \frac{\lambda_\theta + 3\lambda_\phi}{1 - \lambda_\phi}$$



Any arbitrary choice of the experimental observation frame will give the same value of this quantity