

# Identified Two-particle Correlations and Quantum Number Conservation in p-p, p-Pb and Pb-Pb Collisions at LHC Energies

Gyula Bencédi<sup>1</sup>,  
Gergely Gábor Barnaföldi<sup>1</sup>,  
Levente Molnár<sup>2</sup>




<sup>1</sup>Wigner Research Centre for Physics, Budapest, Hungary

<sup>2</sup>Institut Pluridisciplinaire Hubert Curien, Strasbourg, France

September 12, 2014

**10<sup>th</sup> International Workshop on High-pT Physics in the RHIC/LHC era**

**9-12 September 2014, SUBATECH Nantes**



# Outline

## 1. Motivation

- importance of particle correlations at high- $p_T$


## 2. Monte Carlo studies of two-particle correlations in terms of quantum number conservations

- Focusing on identified associated particle spectra
- Identified particle yields from like-sign and unlike-sign correlations
- Comparison of triggered relative yields in p-p, Pb-Pb and p-Pb
  - Collision energy (p-p)
  - Event multiplicity (p-p)
  - Centrality (Pb-Pb)

## 3. Summary, Outlook

# 1. Importance of particle correlations at high $p_T$

- Hadronization in QGP
  - High- $p_T$  hadrons, particle ratios, collective effects
  - High- $p_T$  factorization holds: PDF  $\times$  pQCD  $\times$  FF + jet quenching HI (PID?)
  - Intermediate  $p_T$ : recombination rises
- Fragmentation effects, including PID: recombination effects in intermediate  $p_T$  ( $R_{CP}, R_{AA}, \text{Barion/Meson}, v_2$ )
- Jet-like correlations: ridge (AuAu, dAu, pPb)
  - PID could shed light on formation mechanism...



1.

# Importance of particle correlations at high $p_T$


## Identified two-particle azimuthal correlations

- Identified **triggers**, identified **associateds** integrated in mid-rapidity region

near

- Trigger  $p_T > Assoc p_T$
- Trigger, Assoc  $|\eta| < 1$
- Trigger, Associated species selection both on near and away sides

$\pi^+, \pi^-, K^+, K^-, p^+, p^-, \dots$



$\pi^+, \pi^-, K^+, K^-, p^+, p^-, \dots$

away

**Expectation: conservation of quantum numbers**

$\pi, K, p$  – momentum  $p$ , charge  $Q$  (+K strangeness, +p baryon number)

Fragmentation / hadronization for different particle flavours

## 2.

## Quantum number conservation in correlations

## Monte Carlo datasets and analysis cuts

Collision system	$\sqrt{s} (\sqrt{s_{NN}})$	Statistics
p-p	14 TeV	500 M
p-p	7 TeV	100 M
p-p	2.76 TeV	100 M
p-p	200 GeV	500 M
Pb-Pb (0-10%)	2.76 TeV	4 M
Pb-Pb (30-40%)	2.76 TeV	2 M
Pb-Pb (80-90%)	2.76 TeV	10 M
p-Pb (min.bias)	5.02 TeV	10 M


- p-p events generated by PYTHIA8 (tune 4C)
- Pb-Pb and p-Pb events generated by HIJING
- only charged, final-state particles ( $\pi^\pm$ ,  $K^\pm$ ,  $p$ ,  $\bar{p}$ , and the charged hadrons ( $h^\pm$ ))
- $|\eta_{trig}| > 1$  and  $|\eta_{assoc}| > 1$  were rejected
- transverse momenta of the trigger particles selected within  $2 \text{ GeV}/c < p_{T,trig} < 25 \text{ GeV}/c$
- $p_{T,assoc} < p_{T,trig}$  to avoid double counting
- $p_T$  spectra were projected within  $|\Delta\eta| < 1$

## 2.

## Quantum number conservation in correlations

## Identified associated particle spectra I.

- PID-PID associated  $p_T$  spectra up to high- $p_T$
- p-p @ 7TeV, Pythia (tune320)
- Same side
- Example:  
Trigger particle: **proton**  
in  $2 < p_T [\text{GeV}/c] < 25$  range
- Associated:  $\pi^+, \pi^-, K^+, K^-, p, \bar{p}$
- Acceptance:  $|\eta| < 1$ .



## 2.


## Quantum number conservation in correlations

## Identified associated particle spectra I.

- PID-PID associated  $p_T$  spectra up to high- $p_T$
- p-p @ 7TeV, Pythia (tune320)
- Same side
- Example:
  - Trigger particle: **proton**
  - in  $2 < p_T \text{ [GeV/c]} < 25$  range
- Associated:  $\pi^+, \pi^-, K^+, K^-, p, pbar$
- Acceptance:  $|\eta| < 1$ .

Observation:  
differences between  
proton/anti-proton triggered  
associated yields

Strong effect on the near side




## 2.

# Quantum number conservation in correlations

## Identified associated particle spectra II.

- Quantum number conservation can be tested by plotting the identified associated spectra for identified trigger hadrons
- Effect is on the near side
- The yields of the PID-associated spectra significantly decrease with the selection of charged pion, kaon and proton triggers, respectively
- The basic conservation laws of the quantum numbers – such as **charge (Q)**, **baryon number (B)**, and **strangeness (S)** – are fulfilled and could be reflected in the PID-associated spectra after the hadronization process






## 2.

# Quantum number conservation in correlations

## Identified associated particle spectra II.

- Quantum number conservation can be tested by plotting the identified associated spectra for identified trigger hadrons
- Effect is on the near side
- The yields of the PID-associated spectra significantly decrease with the selection of charged pion, kaon and proton triggers, respectively
- The basic conservation laws of the quantum numbers – such as **charge (Q)**, **baryon number (B)**, and **strangeness (S)** – are fulfilled and could be reflected in the PID-associated spectra after the hadronization process.
- Strength of the effects**



Trigger / Assoc	Strength of effect	
	Near side	Away side
$\pi^+ / (\pi^+, \pi^-)$	$\sim 2$	$\sim 1$
$K^+ / (K^+, K^-)$	$\sim 5$	$\sim 1$
$p / (p, \bar{p})$	$\sim 10$	$\sim 1$

## 2.


# Quantum number conservation in correlations

## Identified associated particle ratios – a closer look

$$\frac{1}{N_{\text{trig}}^i} \times \frac{dN_{\text{assoc}}}{dp_{T,\text{assoc}}}, \quad i \in \{\pi^\pm, K^\pm, p^\pm, h^\pm\}, \quad j \in \{h^\pm\}$$

$$\frac{1}{N_{\text{trig}}^j} \times \frac{dN_{\text{assoc}}}{dp_{T,\text{assoc}}},$$

- pp @ 7TeV, PYTHIA 6 (tune 320)
- $|\Delta\phi| < \pi/2, |\Delta\eta| < 2$ .
- Trigger particle: **proton**
  - in  $2 < p_T \text{ (GeV/c)} < 25$
- Associateds:  $\pi^+, \pi^-, K^+, K^-, p, pbar$
- Acceptance:  $|\eta| < 1$
- In an unmodified fragmentation process the baryon number and charge is conserved and leads to highly correlated distributions in the same phase space



## 2.


# Quantum number conservation in correlations

## Identified associated particle ratios – a closer look

$$\frac{1}{N_{\text{trig}}^i} \times \frac{dN_{\text{assoc}}}{dp_{T,\text{assoc}}}, \quad i \in \{\pi^\pm, K^\pm, p^\pm, h^\pm\}, \quad j \in \{h^\pm\}$$

$$\frac{1}{N_{\text{trig}}^j} \times \frac{dN_{\text{assoc}}}{dp_{T,\text{assoc}}},$$

- pp @ 7TeV, PYTHIA 6 (tune 320)
- $|\Delta\phi| < \pi/2, |\Delta\eta| < 2$ .
- Trigger particle: **proton**
  - in  $2 < p_T \text{ (GeV/c)} < 25$
- Associateds:  $\pi^+, \pi^-, K^+, K^-, p, pbar$
- Acceptance:  $|\eta| < 1$
- In an unmodified fragmentation process the baryon number and charge is conserved and leads to highly correlated distributions in the same phase space



## 2.


# Quantum number conservation in correlations

## Identified particle yields from like-sign and unlike-sign correlations in pp and PbPb collisions

- To extract and enhance the expected quantum number conservation effects, the **ratio** of the PID-triggered-to-charged **hadron-triggered associated spectra** have been plotted

$$\frac{dN_{assoc}^{PID}}{dp_{T,assoc}} \bigg/ \frac{dN_{assoc}^{hadron}}{dp_{T,assoc}}$$

- The splitting effect can be observed for any of the trigger species both in p-p and Pb-Pb collisions. The **splitting is larger on the near-side** than on the away-side.
- Qualitatively the strength of the quantum numbers:  
**charge (Q) < strangeness (S) < baryon number (B)**
- The **largest splitting** effect can be seen for unlike-sign correlations of **protons and antiprotons**.



## 2.


# Quantum number conservation in correlations

## Identified particle yields from like-sign and unlike-sign correlations in pp and PbPb collisions

- To extract and enhance the expected quantum number conservation effects, the **ratio** of the PID-triggered-to-charged **hadron-triggered associated spectra** have been plotted

$$\frac{\frac{dN_{assoc}^{PID}}{dp_{T,assoc}}}{\frac{dN_{assoc}^{hadron}}{dp_{T,assoc}}}$$

- The splitting effect can be observed for any of the trigger species both in p-p and Pb-Pb collisions. The **splitting is larger on the near-side** than on the away-side.
- Qualitatively the strength of the quantum numbers: **charge (Q) < strangeness (S) < baryon number (B)**
- The **largest splitting** effect can be seen for unlike-sign correlations of **protons and antiprotons**.




## 2.

# Quantum number conservation in correlations

## Identified particle yields from like-sign and unlike-sign correlations in pp, pPb, and PbPb collisions

By plotting the differences between the (+/-) unlike-sign and (+/+) like-sign trigger/associated particles pairs for the PID-triggered-to/charged hadron triggered yields

$$\left( \frac{dN_{assoc}^{PID}}{dp_{T,assoc}} / \frac{dN_{assoc}^{hadron}}{dp_{T,assoc}} \right)^{+/-} / \left( \frac{dN_{assoc}^{PID}}{dp_{T,assoc}} / \frac{dN_{assoc}^{hadron}}{dp_{T,assoc}} \right)^{+/+} = \left( \frac{dN_{assoc}^{PID}}{dp_{T,assoc}} \right)^{+/-} / \left( \frac{dN_{assoc}^{PID}}{dp_{T,assoc}} \right)^{+/+}$$



## 2.


# Quantum number conservation in correlations

## Identified particle yields from like-sign and unlike-sign correlations in pp, pPb, and PbPb collisions

By plotting the differences between the (+/-) unlike-sign and (+/+) like-sign trigger/associated particles pairs for the PID-triggered-to/charged hadron triggered yields

$$\left( \frac{dN_{assoc}^{PID}}{dp_{T,assoc}} / \frac{dN_{assoc}^{hadron}}{dp_{T,assoc}} \right)^{+/-} / \left( \frac{dN_{assoc}^{PID}}{dp_{T,assoc}} / \frac{dN_{assoc}^{hadron}}{dp_{T,assoc}} \right)^{+/+} = \left( \frac{dN_{assoc}^{PID}}{dp_{T,assoc}} \right)^{+/-} / \left( \frac{dN_{assoc}^{PID}}{dp_{T,assoc}} \right)^{+/+}$$

NEAR



## 2.

# Quantum number conservation in correlations


## Identified particle yields from like-sign and unlike-sign correlations in pp, pPb, and PbPb collisions

By plotting the differences between the (+/-) unlike-sign and (+/+) like-sign trigger/associated particles pairs for the PID-triggered-to/charged hadron triggered yields

$$\left( \frac{dN_{assoc}^{PID}}{dp_{T,assoc}} / \frac{dN_{assoc}^{hadron}}{dp_{T,assoc}} \right)^{+/-} / \left( \frac{dN_{assoc}^{PID}}{dp_{T,assoc}} / \frac{dN_{assoc}^{hadron}}{dp_{T,assoc}} \right)^{+/+} = \left( \frac{dN_{assoc}^{PID}}{dp_{T,assoc}} \right)^{+/-} / \left( \frac{dN_{assoc}^{PID}}{dp_{T,assoc}} \right)^{+/+}$$

NEAR

**PION:** The PID-triggered associated spectra for the like-sign and unlike-sign trigger/associated particle pairs is **similar for pions** in p-p, p-Pb and Pb-Pb collisions





## 2.

# Quantum number conservation in correlations

## Identified particle yields from like-sign and unlike-sign correlations in pp, pPb, and PbPb collisions

By plotting the differences between the (+/-) unlike-sign and (+/+) like-sign trigger/associated particles pairs for the PID-triggered-to/charged hadron triggered yields


$$\left( \frac{dN_{assoc}^{PID}}{dp_{T,assoc}} / \frac{dN_{assoc}^{hadron}}{dp_{T,assoc}} \right)^{+/-} / \left( \frac{dN_{assoc}^{PID}}{dp_{T,assoc}} / \frac{dN_{assoc}^{hadron}}{dp_{T,assoc}} \right)^{+/+} = \left( \frac{dN_{assoc}^{PID}}{dp_{T,assoc}} \right)^{+/-} / \left( \frac{dN_{assoc}^{PID}}{dp_{T,assoc}} \right)^{+/+}$$

NEAR

The PID-triggered associated spectra for the like-sign and unlike-sign trigger/associated particle pairs is **similar for pions** in p-p, p-Pb and Pb-Pb collisions

### KAON / PROTON

In contrast, there is **clear difference** between the p-p, p-Pb and Pb-Pb systems for kaons and protons at the near side.



## 2.

# Quantum number conservation in correlations

## Identified particle yields from like-sign and unlike-sign correlations in pp, pPb, and PbPb collisions

By plotting the differences between the (+/-) unlike-sign and (+/+) like-sign trigger/associated particles pairs for the PID-triggered-to/charged hadron triggered yields


$$\left( \frac{dN_{assoc}^{PID}}{dp_{T,assoc}} / \frac{dN_{assoc}^{hadron}}{dp_{T,assoc}} \right)^{+/-} / \left( \frac{dN_{assoc}^{PID}}{dp_{T,assoc}} / \frac{dN_{assoc}^{hadron}}{dp_{T,assoc}} \right)^{+/+} = \left( \frac{dN_{assoc}^{PID}}{dp_{T,assoc}} \right)^{+/-} / \left( \frac{dN_{assoc}^{PID}}{dp_{T,assoc}} \right)^{+/+}$$

The PID-triggered associated spectra for the like-sign and unlike-sign trigger/associated particle pairs is **similar for pions** in p-p, p-Pb and Pb-Pb collisions.

In contrast, there is clear **difference** between the p-p, p-Pb and Pb-Pb systems for **kaons and protons** at the near side.

**Less deviance on the away side.**

09/12/14




# 2.

## Quantum number conservation in correlations

### Yield dependence on collision energy in p-p

Near side:  
the splitting  
decreases with  
increasing  $\sqrt{s}$




# 2.

## Quantum number conservation in correlations

### Yield dependence on collision energy in p-p

**Near side:**  
the splitting  
decreases with  
increasing  $\sqrt{s}$

**Away side:**  
no obvious energy  
dependence, kaons  
show small  
asymmetry for all  $\sqrt{s}$




## 2.

# Quantum number conservation in correlations

## Yield dependence on event multiplicity in p-p

**Near side:**  
the higher the event multiplicity the smaller the size of the splitting due to increasing underlying event contribution at higher multiplicities




## 2. Quantum number conservation in correlations

### Yield dependence on event multiplicity in p-p

**Near side:**  
the higher the event multiplicity the smaller the size of the splitting due to increasing underlying event contribution at higher multiplicities

**Away side:**  
no obvious dependence with event multiplicity




## 2.

# Quantum number conservation in correlations

## Yield dependence on centrality in Pb-Pb

**Near side:**  
reverse evolution  
pattern is observed  
for kaons and  
protons as a  
function of centrality




# 2.

## Quantum number conservation in correlations

### Yield dependence on centrality in Pb-Pb

**Near side:**  
reverse evolution  
pattern is observed  
for kaons and  
protons as a  
function of centrality

**Away side:**  
kaons and protons  
show similar  
reversal trend as on  
the near side






## 2.

## Quantum number conservation in correlations

 $\Delta\phi$  projections

p-p 7TeV, PYTHIA 6  
(Perugia0)

trigger  $p_T$  bins




# 2.

# Quantum number conservation in correlations

$\Delta\phi$  projections

p-p 7TeV, PYTHIA 6  
(Perugia0)

trigger  $p_T$  bins




# 2.

# Quantum number conservation in correlations

## $\Delta\phi$ projections

Pb-Pb 2.76TeV,  
HIJING




# 2.

# Quantum number conservation in correlations

$\Delta\phi$  projections


Pb-Pb 2.76TeV,  
HIJING



$\pi^+/\pi^-$

$K^+/K^-$

$p/\bar{p}$



2.

# Quantum number conservation in correlations

$\Delta\phi$  projections

$\pi^+/\pi^-$

$K^+/K^-$

$p/pbar$

$\pi^+/\pi^-$

$K^+/K^-$

$p/pbar$


**PYTHIA – pp / NEAR side**

**$p/pbar$  is more correlated than  $K^+/K^-$**

**HIJING – PbPb / NEAR side**

**No difference between  $K^+/K^-$  and  $p/pbar$**

trigger  $p_T$  bins



**p-p 7TeV, PYTHIA 6  
(Perugia0)**

**Pb-Pb 2.76TeV,  
HIJING**

# 3.

## Summary and Outlook

---

MC simulations and analysis have been performed to study PID-PID azimuthal correlations at mid-rapidity

The observed splitting effect of the PID-triggered-to-charged hadron-triggered associated spectra shows

- **a peculiar pattern which**
  - The the most prominent for p-p unlike-sign associated particle pairs (compared to K+/K-, +- pairs) on the near- and away side;
  - has a reverse behaviour in Pb-Pb and in p-Pb on the away-side and remains unchanged on the near-side in Pb-Pb;
  - has similar behaviour in the p-p, peripheral Pb-Pb and minimum bias p-Pb on the near-side in terms of the conservation of baryon numbers;
  - decreases towards higher event multiplicities in p-p;
  - decreases with increasing collision energies from RHIC to LHC in p-p.
- **No experimental measurements to contrast with the observed MC analysis**
    - Further Monte Carlo checks needed to have a better understanding...
  - Analysis has been started to perform the same analysis exploiting the PID capabilities of ALICE at the LHC
  - The observed interesting patterns can be measured at higher momentum in ALICE
    - In principle this can be done by the TPC, statistical method: relativistic dE/dx
    - Purity PID cuts

# 3.

## Summary and Outlook

---





### References:

- [1] ALICE Collaboration, T. Acconcia et al., “A Very High Momentum Particle Identification Detector,” <http://arxiv.org/abs/1309.5880>arXiv:1309.5880 [nucl-ex]
- [2] G. Bencédi, G. G. Barnaföldi, and L. Molnár, “Monte Carlo Studies of Identified Two-particle Correlations in p-p and Pb-Pb Collisions,” <http://arxiv.org/abs/1403.0117>arXiv:1403.0117 [hep-ex]

# Backups...





NLO pQCD AKK FF : p+p collisions at 200 GeV



09/12/14

# Azimuthal correlations





**ZYAM: Zero Yield At Minimum**

$$\Delta\phi = \phi_{\text{trigger}} - \phi_{\text{assoc}}, \quad \Delta\eta = \eta_{\text{trigger}} - \eta_{\text{assoc}}$$

## 2.

## Quantum number conservation in correlations

 $\Delta\phi$  projections (flavor conservations, K)

- **Near side** flavor and charge correlations decrease as a function of  $p_{T,\text{assoc}}$  and  $p_{T,\text{trig}}$  (the width narrows) compared to the away side correlations
- **Away side** stays roughly constant when the trigger particle momentum is increased

# 2. Quantum number conservation in correlations

## $\Delta\phi$ projections

$\pi^+/\pi^-$


$K^+/K^-$

$p/p\bar{b}$

$\pi^+/\pi^-$

$K^+/K^-$

$p/p\bar{b}$



**p-p 7TeV, PYTHIA 6  
(Perugia0)**

**Pb-Pb 2.76TeV,  
HIJING**