



# Talk Outline

- Vector-like fermions LHC (Indirect) signatures
  - simple vector-like extensions of SM
  - in electroweak precision observables
  - find regions favored by  $hgg$ ,  $h\gamma\gamma$  at the LHC
- Vector-like fermions LHC (Direct) signatures
  - $b'_{(-1/3)}$ ,  $t'_{(2/3)}$ ,  $\chi_{(5/3)}$
  - single and double resonant channels
  - find luminosity required for discovery


# Vectorlike $\psi$

- Theory with Vectorlike fermions:
  - both  $\chi$  and  $\chi^c$  present
  - can write vectorlike mass term  $\mathcal{L} \supset -M \chi \chi^c + h.c.$ 
    - contrast with SM (chiral theory):
 

$q_L = (3, 2)_{1/6}$	No $(\bar{3}, \bar{2})_{-1/6}$
$U_R = (3, 1)_{+2/3}$	No $(\bar{3}, 1)_{-2/3}$
$D_R = (3, 1)_{-1/3}$	No $(\bar{3}, 1)_{1/3}$
- For a VL pair, define a Dirac state  $\mathcal{X} \equiv \begin{pmatrix} \chi_\alpha \\ \chi^{c\dot{\alpha}} \end{pmatrix}$ 
  - in terms of which the mass term is:  $\mathcal{L} \supset -M \bar{\mathcal{X}} \mathcal{X}$
  - Eg: SU(2) doublet  $\mathcal{X} \equiv \begin{pmatrix} \mathcal{X}_1 \\ \mathcal{X}_2 \end{pmatrix}$

# VL fermions Decoupling

- Independent source of mass  $M$  (not given by  $m = \lambda v$ )
  - Can make  $M$  arbitrarily large
    - without hitting Landau pole in Yukawa coupling (4th Gen)
  - $M$  could be related to EW scale (or not)
    - Eg. of connection: ExtraDim Th  $M = M_{KK} \sim TeV$ , SUSY solutions to  $\mu$  problem
  - Nice decoupling behavior :  $S, T, U, h \rightarrow \gamma\gamma, gg \rightarrow h, \dots$ 
    - For instance  $h\gamma\gamma, ggh$  couplings



# Implications of VL Theory

Vectorlike fermions	Chiral (4-gen) fermions
$M$ allowed by EW symmetry	$m$ only after EWSB, $= \lambda \langle H \rangle$
can be arbitrarily heavy	Landau pole in Yukawa coupling $\lambda$
CC + NC tree-level decays	only CC tree-level decays
loops decoupling	some loops nondecoupling

# VECTOR-LIKE FERMIONS IN ELECTROWEAK PRECISION & HIGGS OBSERVABLES

[S.Ellis, R.Godbole, SG, J.Wells; To Appear]

# Simple VL extensions of SM

- $1\bar{1}$  :  $SU(2)$  singlet VL pair
- $2\bar{2}$  :  $SU(2)$  doublet VL pair
- $2\bar{2} + 1\bar{1}$  : MVSM
- $2\bar{2} + 1\bar{1} + 1\bar{1}$  : Vector-like extension of the SM (VSM)

$2\bar{2} + 1\bar{1} + 1\bar{1} : \text{VSM}$ 

- $VSM \equiv VLQ (\mathcal{X}_Q, \xi_U, \Upsilon_D) \oplus VLL (\mathcal{X}_L, \xi_N, \Upsilon_E)$ 
  - where  $\mathcal{X} = (2, Y_{\mathcal{X}})$ ;  $\Upsilon = (1, Y_{\mathcal{X}} - 1/2)$ ;  $\xi = (1, Y_{\mathcal{X}} + 1/2)$

$$\mathcal{L}_{\text{Yuk}} \supset -\lambda_{\xi} \bar{\mathcal{X}} \cdot H^* \xi - \lambda_{\Upsilon} \bar{\mathcal{X}} H \Upsilon + h.c.$$

$Y_{\mathcal{X}} = \pm Y_{SM}$  assignments:

$Y_{\mathcal{X}}$	-1/2	-1/6	1/6	1/2
$Q_1, Q_4$	0	1/3	2/3	1
$Q_2, Q_3$	-1	-2/3	-1/3	0

$$\mathcal{L}_{\text{mass}} \supset -(\bar{\mathcal{X}}_1 \quad \bar{\xi}) \begin{pmatrix} M_{\mathcal{X}} & \tilde{m} \\ \tilde{m} & M_{\xi} \end{pmatrix} \begin{pmatrix} \mathcal{X}_1 \\ \xi \end{pmatrix} - (\bar{\mathcal{X}}_2 \quad \bar{\Upsilon}) \begin{pmatrix} M_{\mathcal{X}} & m \\ m & M_{\Upsilon} \end{pmatrix} \begin{pmatrix} \mathcal{X}_2 \\ \Upsilon \end{pmatrix}$$

Diagonalize and obtain  $W_{\mu}^a$ ,  $B_{\mu}$  and  $h$  couplings

We assume tiny VL-SM mixing Yukawa terms



# Mixing with SM fields?

$$\mathcal{L}_{\text{Yuk}} \supset -\lambda'_\xi \bar{Q} \cdot H^* \xi - \lambda'_\Upsilon \bar{Q} H \Upsilon - \lambda'_U \bar{\chi} \cdot H^* U - \lambda'_D \bar{\chi} H D + h.c.$$


similarly for the VL-leptons

- EWSB  $\langle H \rangle = v/\sqrt{2}$  will mix SM  $\leftrightarrow$  VL fermions
  - Here, take  $\lambda'$  small
    - such that flavor constraints are satisfied
    - $Zb\bar{b}$  coupling is not shifted too much
    - but big enough to have prompt decays
    - no significant effect in Higgs observables

For sizable mixing case, see: [Dawson, Furlan '12] [Aguilar-Saveedra '13] [Fajfer et al. '13]  
[Dermisek, Raval '13]


# Observables

Precision electroweak observables ( $S, T, U$ )





Modifications to  $hgg$ ,  $h\gamma\gamma$  couplings:

$\sigma(gg \rightarrow h)$



$\Gamma(h \rightarrow \gamma\gamma)$




We compute ratios  $\frac{\Gamma_{h \rightarrow gg}}{SM}$ ,  $\frac{\Gamma_{h \rightarrow \gamma\gamma}}{SM}$


using leading-order expressions


QCD corrections to ratios small: [Furlan '11] [Gori, Low '13]



$$\mu_{\gamma\gamma}^{VBF} \approx \frac{\Gamma_{\gamma\gamma}}{\Gamma_{\gamma\gamma}^{SM}}; \quad \mu_{ZZ}^{ggh} \approx \frac{\Gamma_{gg}}{\Gamma_{gg}^{SM}}; \quad \mu_{\gamma\gamma}^{ggh} \approx \frac{\Gamma_{gg}}{\Gamma_{gg}^{SM}} \frac{\Gamma_{\gamma\gamma}}{\Gamma_{\gamma\gamma}^{SM}}; \quad \frac{\mu_{\gamma\gamma}^{ggh}}{\mu_{ZZ}^{ggh}} \approx \frac{\Gamma_{\gamma\gamma}}{\Gamma_{\gamma\gamma}^{SM}} \approx \mu_{\gamma\gamma}^{VBF}$$

$2\bar{2} + 1\bar{1} : MVLE$ 



$\lambda_E = 1, M_E = M_L, Y_L = (-1/2, 1/2)$  (solid,dashed)






$2\bar{2} + 1\bar{1} : \text{MVLE}$ 

$2\bar{2} + 1\bar{1} : MVQD$ 

$\lambda_D = 1, M_D = M_Q, Y_Q = (1/6, -1/6)$  (solid, dashed)





$2\bar{2} + 1\bar{1} : MVQD$ 

$2\bar{2} + 1\bar{1} + 1\bar{1}$  : VSM

Category	$M_q$ (GeV)	$M_\ell$ (GeV)	Color
Light	$\leq 700$	$\leq 450$	Red
Medium	(700, 1000)	(450, 750)	Blue
Heavy	$> 1000$	$> 750$	Green

$$M_{\{Q,U,D,L,E,N\}} = M_{VL}, Y_Q = 1/6, Y_L = -1/2$$

## EWPT &amp; Higgs Observables

 $2\bar{2} + 1\bar{1} + 1\bar{1} : \text{VSM}$ 

$$M_Q = M_U = M_D, \quad M_L = M_E = M_N, \quad \lambda_U = \lambda_D \equiv \lambda_Q, \quad \lambda_E = \lambda_N \equiv \lambda_L, \quad Y_Q = 1/6, \quad Y_L = -1/2$$





$\chi^2$  fit to the LHC Data


[ATLAS arXiv:1307.1427] [CMS-PAS-HIG-13-005, 2013]

Coupling	ATLAS	CMS
$\kappa_g$	$1.04 \pm 0.14$	$0.83 \pm 0.11$
$\kappa_\gamma$	$1.2 \pm 0.15$	$0.97 \pm 0.18$

$$\chi^2 = \sum_{i=1}^4 \left( \kappa_i^{\text{Exp}} - \kappa_i^{\text{Th}} \right)^2 / \left( \sigma_i^{\text{Exp}} \right)^2$$

$\chi^2$  fit to LHC Higgs Data $2\bar{2} + 1\bar{1} + 1\bar{1}$  : VSM  $\chi^2$  fit

$$Y_Q = 1/6, Y_L = -1/2, \lambda_q = 1, \lambda_\ell = 1$$

$\chi^2$  fit to LHC Higgs Data $2\bar{2} + 1\bar{1} + 1\bar{1}$  : VSM  $\chi^2$  fit

$$Y_Q = 1/6, Y_L = -1/2, M_q = 1000 \text{ GeV}, M_\ell = 500 \text{ GeV}$$

# DIRECT PRODUCTION OF $t'$ , $b'$ , $\chi_{5/3}$ VECTORLIKE FERMIONS AT THE LHC

Model independent analysis.

Benchmark points from *Warped extra dimensions*

[SG, T.Mandal, S.Mitra, R.Tibrewala, arXiv:1107.4306, PRD84 (2011) 055001]

[SG, T.Mandal, S.Mitra, G.Moreau : arXiv:1306.2656]

See Also: [SG, Moreau, Singh, '10]

[Dennis et al, '07] [Carena et al, '07] [Contino, Servant, '08]

[Atre et al, '08, '09, '11] [Han et al. '10] [Aguilar-Saavedra, '09] [Mrazek, Wulzer, '09]

[SG, Moreau, Singh, '10][Bini et al. '12][Buchkremer et al. '13]

# Decay Modes of $t'$ , $b'$ , $\chi$

EWSB induced mixing  $\implies$  Tree-level NC Couplings

- as usual will have  $t'_L b_L W^\pm$  and  $b'_L t_L W^\pm$  CC couplings
- also, from Yukawa coupling  $\langle \Sigma \rangle = v \implies t \leftrightarrow t'$ ,  $b \leftrightarrow b'$  mixing

$$\mathcal{L} \supset ( \begin{array}{cc} b & b' \end{array} ) \gamma^\mu \left( \begin{array}{cc} g_Z & 0 \\ 0 & g'_Z \end{array} \right) \left( \begin{array}{c} b \\ b' \end{array} \right)_{L,R} Z_\mu + ( \begin{array}{cc} b_L & b'_L \end{array} ) \left( \begin{array}{cc} m_b & 0 \\ \tilde{m}_b & M_{b'} \end{array} \right) \left( \begin{array}{c} b_R \\ b'_R \end{array} \right) + h.c.$$

- Diagonalize to go to mass basis
  - $v \rightarrow v(1 + h/v)$  leads to  $b'bh$  coupling
  - $g_Z \neq g'_Z$  leads to  $b'bZ$  coupling
  - Similarly  $t'tZ$ ,  $t'th$  couplings also, in addition to  $t'bw$

# Decay Modes of $t'$ , $b'$ , $\chi$

EWSB induced mixing  $\implies$  Tree-level NC Couplings

- as usual will have  $t'_L b_L W^\pm$  and  $b'_L t_L W^\pm$  CC couplings
- also, from Yukawa coupling  $\langle \Sigma \rangle = v \implies t \leftrightarrow t'$ ,  $b \leftrightarrow b'$  mixing

$$\mathcal{L} \supset ( \begin{matrix} b & b' \end{matrix} ) \gamma^\mu \begin{pmatrix} g_Z & 0 \\ 0 & g'_Z \end{pmatrix} \begin{pmatrix} b \\ b' \end{pmatrix}_{L,R} Z_\mu + ( \begin{matrix} b_L & b'_L \end{matrix} ) \begin{pmatrix} m_b & 0 \\ \tilde{m}_b & M_{b'} \end{pmatrix} \begin{pmatrix} b_R \\ b'_R \end{pmatrix} + h.c.$$


- Diagonalize to go to mass basis
  - $v \rightarrow v(1 + h/v)$  leads to  $b'bh$  coupling
  - $g_Z \neq g'_Z$  leads to  $b'bZ$  coupling
  - Similarly  $t'tZ$ ,  $t'th$  couplings also, in addition to  $t'bw$
- VL Tree-level Decays
  - $b' \rightarrow tW$ ,  $b' \rightarrow bZ$ ,  $b' \rightarrow bh$
  - $t' \rightarrow bW$ ,  $t' \rightarrow tZ$ ,  $t' \rightarrow th$
  - $\chi \rightarrow tW$

## $b'$ Phenomenology at the LHC

[SG, T.Mandal, S.Mitra, R.Tibrewala, arXiv:1107.4306, PRD84 (2011) 055001]

[SG, T.Mandal, S.Mitra, G.Moreau : arXiv:1306.2656]

See Also: [Contino, Servant '08][Mrazek, Wulzer '10]

$b'$  Single & Double Resonant channels



... followed by  $b_2 \rightarrow bZ$

- Both  $b_2$  on-shell : **Double Resonant (DR)** channel
- Only one  $b_2$  on-shell : **Single Resonant (SR)** channel
  - $|M(bZ) - M_{b_2}| \geq \alpha_{cut} M_{b_2}; \quad \alpha_{cut} = 0.05$



$b'$  Double Resonant

Pair Production :  $pp \rightarrow b'\bar{b}' \rightarrow bZ\bar{b}Z \rightarrow bj\bar{j}l\bar{l}$




**Cuts:**  
*Rapidity:*  $-2.5 < y_{b,j,Z} < 2.5$ ,  
*Transverse momentum:*  $p_{T,b,j,Z} > 25$  GeV,  
*Invariant mass cuts:*  
 $M_Z - 10$  GeV  $< M_{jj} < M_Z + 10$  GeV,  
 $0.95M_{b_2} < M_{(bZ)} < 1.05M_{b_2}$ .

$b'$  Single Resonant - I



Single Resonant :  $bg \rightarrow b'bZ \rightarrow bZbZ \rightarrow bbJJ\ell\ell$

Model Independent LHC-14 reach



Brown dots : DT Model

Green dots : TT Model

$b'$  Single Production - IISingle Production :  $bg \rightarrow b'Z \rightarrow bZZ \rightarrow bj\ell\ell$ 

Cuts:

Rapidity:  $-2.5 < y_{b,j,Z} < 2.5$ ,Transverse momentum:  $p_{T,b,j,Z} > 0.1M_{b_2}$ ,


Invariant mass cuts:

 $M_Z - 10$  GeV  $< M_{jj} < M_Z + 10$  GeV, $0.95M_{b_2} < M(bZ)$  OR  $(bjj) < 1.05M_{b_2}$ .

## $\chi$ Phenomenology at the LHC


[SG, T.Mandal, S.Mitra, G.Moreau : arXiv:1306.2656]

[Contino, Servant '08][Mrazek, Wulzer '10][Cacciapaglia et al. '12]

$\chi$  Double and Single Resonant channels

$$pp \rightarrow \chi t W \rightarrow t W t W \rightarrow t W t \nu$$

$X$	$M_\chi$ (GeV)	$\sigma_{tot}$ (fb)	$\sigma_{SR}$ (fb)	cuts	$S$ (fb)	BG (fb)	$\mathcal{L}$ ( $fb^{-1}$ )
$X_1$	500	2406	261.5	Basic	977.5	3.257	-
				Disc.	146.1	0.115	0.826
$X_2$	750	235.5	29.31	Basic	99.99	3.257	-
				Disc.	42.74	0.115	2.824
$X_3$	1000	39.19	5.198	Basic	17.92	3.257	-
				Disc.	11.36	0.115	10.63
$X_4$	1250	8.576	1.231	Basic	4.305	3.257	-
				Disc.	3.226	0.115	37.42
$X_5$	1500	2.188	0.364	Basic	1.235	3.257	-
				Disc.	1.010	0.115	119.5
$X_6$	1750	0.613	0.121	Basic	0.393	3.257	-
				Disc.	0.339	0.115	355.8

$\chi$  Single Resonant Channel


Blue Dots - ST Model

Green Dots - TT Model

## $t'$ Phenomenology at the LHC

[SG, Tanumoy Mandal, Subhadip Mitra, Gregory Moreau : arXiv:1306.2656]


See also: [Harigaya et al., '12] [Giridhar, Mukhopadhyaya, 2012] [Azatov et al., '12]  
[Berger, Hubisz, Perelstein, '12] [Cacciapaglia et al., '10, '12] [Aguilar-Saavedra et al. '05]

$t'$  Double and Single Resonant channels

$pp \rightarrow t_2 th \rightarrow thth \rightarrow tbbtbb \rightarrow 6 b 4 j$  (4 b-tags)

$T$	$M_{t_2}$ (GeV)	$\sigma_{tot}$ (fb)	$\sigma_{SR}$ (fb)	cuts	$S$ (fb)	BG (fb)	$\mathcal{L}$ ( $fb^{-1}$ )
$T_1$	500	1207	223.0	Basic	237.4	102.7	-
				Disc.	52.38	0.389	6.379
$T_2$	750	115.2	18.30	Basic	22.67	102.7	-
				Disc.	13.25	0.389	25.22
$T_3$	1000	18.38	2.715	Basic	3.088	102.7	-
				Disc.	2.421	0.389	138.0
$T_4$	1250	3.821	0.590	Basic	0.477	102.7	-
				Disc.	0.415	0.389	1889.2



$t'$  Single Resonant channel

Blue Dots - ST Model

Green Dots - TT Model

# Conclusions

- Considered Vector-like fermion extensions of the SM
- Indirect probes
  - $hgg$ ,  $h\gamma\gamma$  couplings may show deviations in the future
  - otherwise constraint on parameter space
- Direct LHC signatures
  - have NC (and CC) tree-level decays
  - Identified promising DR and SR channels
    - SR can probe EW couplings
  - 14 TeV LHC with  $\approx 300 \text{ fb}^{-1}$  reach about 1.5 - 2 TeV in DR
    - Already ATLAS & CMS limits around 700 GeV

# BACKUP SLIDES

BACKUP SLIDES

# Warped Model

SM in background 5D warped AdS space

[Randall, Sundrum '99]



$$ds^2 = e^{-2k|y|}(\eta_{\mu\nu} dx^\mu dx^\nu) + dy^2$$

$Z_2$  orbifold fixed points:

- Planck (UV) Brane
- TeV (IR) Brane

$R$  : radius of Ex. Dim.

$k$  : AdS curvature scale ( $k \lesssim M_{pl}$ )



Hierarchy prob soln:


- IR localized Higgs :  $M_{EW} \sim ke^{-k\pi R}$  : Choose  $k\pi R \sim 34$ 
  - Gauge-theory dual is a composite Higgs model

# Explaining SM mass hierarchy

Bulk Fermions explain SM mass hierarchy [Gherghetta, Pomarol 00][Grossman, Neubert '00]

$$\mathcal{L}_{Yuk}^{(5)} \supset \sqrt{|g|} \{ c_L k \bar{\psi}_L \psi_L + c_R k \bar{\psi}_R \psi_R + (\lambda_5 \bar{\psi}_R \psi_L H + h.c.) \}$$

$$\Psi_L(x, y) = \frac{e^{(2-c)ky}}{\sqrt{2\pi R N_0}} \Psi_L^{(0)}(x) + \dots$$



FCNC largely under control, but still strong constraints

# AdS/CFT Correspondence

## AdS/CFT Correspondence

[Maldacena, 1997]

- A classical supergravity theory in  $AdS_5 \times S_5$  at weak coupling is **dual** to a 4D large-N CFT at strong coupling
- The CFT is at the boundary of  $AdS$  [Witten 1998; Gubser, Klebanov, Polyakov 1998]

$$Z_{CFT}[\phi_0] = e^{-\Gamma_{AdS}[\phi_0]}$$

$\mathcal{L} \supset \int d^4x \mathcal{O}_{CFT}(x) \phi_0(x)$ Eg: $\langle \mathcal{O}(x_1) \mathcal{O}(x_2) \rangle = \frac{\delta^2 Z_{CFT}[\phi_0]}{\delta \phi_0(x_1) \delta(x_2)}$ with $Z_{CFT}$ given by the RHS	$\Gamma_{AdS}[\phi]$ supergravity eff. action $\phi(y, x)$ is a solution of the EOM ( $\delta\Gamma = 0$ ) for given bndry value $\phi_0(x) = \phi(y = y_0, x)$
--	---

# 4D Duals of Warped Models

[Arkani-Hamed, Porrati, Randall, 2000; Rattazzi, Zaffaroni, 2001]

- Dual of Randall-Sundrum model **RS1 (SM on IR Brane)**
  - Planck brane  $\implies$  UV Cutoff; Dynamical gravity in the 4D CFT
  - TeV (IR) brane  $\implies$  IR Cutoff; Conformal invariance broken below a TeV
    - All SM fields are composites of the CFT
  
- Dual of Warped Models with **Bulk SM**
  - UV localized fields are elementary
  - IR localized fields (Higgs) are composite
    - 4D dual is Composite Higgs model [Georgi, Kaplan 1984]
    - Shares many features with Walking Extended Technicolor
  - Partial Compositeness
    - AdS dual is weakly coupled and hence calculable!
  - KK states are dual to composite resonances

# Warped Bulk Gauge Group

[Agashe, Delgado, May, Sundrum '03]

Bulk gauge group :  $SU(3)_{QCD} \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_X$

- Gauge Symmetry breaking:

- By Boundary Condition (BC):

$A_{-+}(x, y)$  BC:  $A|_{y=0} = 0$ ;  $\partial_y A|_{y=\pi R} = 0$

- $SU(2)_R \times U(1)_X \rightarrow U(1)_Y$

- By VEV of IR localized Higgs

Higgs  $\Sigma = (2, 2)_0$

- $SU(2)_L \times U(1)_Y \rightarrow U(1)_{EM}$



# Warped Fermions


- SM fermions :  $(+, +)$  BC  $\rightarrow$  zero-mode
- “Exotic” fermions :  $(-, +)$  BC  $\rightarrow$  No zero-mode
  - 1<sup>st</sup> KK vectorlike fermion

- Typical  $c_{t_R}, c_{t_L}$  :  $(-, +)$  top-partners “light”

$c$  : Fermion bulk mass parameter

[Choi, Kim, 2002] [Agashe, Delgado, May, Sundrum, 03]  
 [Agashe, Perez, Soni, 04] [Agashe, Servant 04]

- Look for it at the LHC



[Dennis et al, '07] [Carena et al, '07] [Contino, Servant, '08]  
 [Atre et al, '09, '11] [Aguilar-Saavedra, '09] [Mrazek, Wulzer, '09]  
 [SG, Moreau, Singh, '10] [SG, Mandal, Mitra, Tibrewala, '11] [SG, Mandal, Mitra, Moreau : '13]

# Fermion rep : $Zb\bar{b}$ not protected (DT model)

[Agashe, Delgado, May, Sundrum '03]

- Complete  $SU(2)_R$  multiplet
  - $Q_L \equiv (\mathbf{2}, \mathbf{1})_{1/6} = (t_L, b_L)$
  - $\psi_{t_R} \equiv (\mathbf{1}, \mathbf{2})_{1/6} = (t_R, b')$
  - $\psi_{b_R} \equiv (\mathbf{1}, \mathbf{2})_{1/6} = (T, b_R)$ 
    - "Project-out"  $b'$ ,  $T$  zero-modes by  $(-, +)$  B.C.
    - New  $\psi_{VL} : b', T$
- $b \leftrightarrow b'$  mixing
  - $Zb\bar{b}$  coupling shifted
    - So LEP constraint quite severe

# Fermion rep : $Zb\bar{b}$ protected (ST & TT models)

- $Q_L = (2, 2)_{2/3} = \begin{pmatrix} t_L & \chi \\ b_L & T \end{pmatrix}$  [Agashe, Contino, DaRold, Pomarol '06]
- $Zb_L\bar{b}_L$  protected by custodial  $SU(2)_{L+R} \otimes P_{LR}$  invariance  
 $Wt_L b_L, Zt_L t_L$  not protected, so shifts

Fermion rep :  $Zb\bar{b}$  protected (ST & TT models)

- $Q_L = (2, 2)_{2/3} = \begin{pmatrix} t_L & \chi \\ b_L & T \end{pmatrix}$  [Agashe, Contino, DaRold, Pomarol '06]
  - $Zb_L\bar{b}_L$  protected by custodial  $SU(2)_{L+R} \otimes P_{LR}$  invariance  
 $Wt_L b_L, Zt_L t_L$  not protected, so shifts

Two  $t_R$  possibilities:

① Singlet  $t_R$  (ST Model) :  $(1, 1)_{2/3} = t_R$       New  $\psi_{VL} : \chi, T$

② Triplet  $t_R$  (TT Model) :

$$(1, 3)_{2/3} \oplus (3, 1)_{2/3} = \psi'_{t_R} \oplus \psi''_{t_R} = \begin{pmatrix} \frac{t_R}{\sqrt{2}} & \chi' \\ -\frac{t_R}{\sqrt{2}} & b' \end{pmatrix} \oplus \begin{pmatrix} \frac{t''}{\sqrt{2}} & \chi'' \\ b'' & -\frac{t''}{\sqrt{2}} \end{pmatrix}$$

New  $\psi_{VL} : \chi, T, \chi', b', \chi'', t'', b''$

# Yukawa Couplings

## Yukawa Couplings






- No  $Zb\bar{b}$  protection





$$\mathcal{L}_{\text{Yuk}} \supset \lambda_t \bar{Q}_L \Sigma \psi_{tR} + \lambda_b \bar{Q}_L \Sigma \psi_{bR} + h.c.$$


- With  $Zb\bar{b}$  protection

- ST Model  $\mathcal{L}_{\text{Yuk}} \supset \lambda_t \text{Tr}[\bar{Q}_L \Sigma] t_R + h.c.$


- TT Model  $\mathcal{L}_{\text{Yuk}} \supset \lambda_t \text{Tr}[\bar{Q}_L \Sigma \psi'_{tR}] + \lambda'_t \text{Tr}[\bar{Q}_L \Sigma \psi''_{tR}] + h.c.$

Warped model  $b'$  parameters

Warped model  $\chi$  parameters






Warped model  $\Gamma_\chi$ 




ST Model



TT Model



Warped model  $t'$  parameters

Warped model  $t'$  BR


# $b'$ Pair Production Details

$$pp \rightarrow b' \bar{b}' \rightarrow bZ\bar{b}Z \rightarrow bj\bar{j}\bar{b}ll$$

$M_{b_2}$ (GeV)	Signal $\sigma_s$ (in fb)		Background $\sigma_b$ (in fb)				$\mathcal{L}$ ( $\text{fb}^{-1}$ )
	$bZbZ$		$bZbZ$		$(bj\bar{j}bZ)_{\text{tot}}$		
	$y, p_T$ cuts	All cuts	$y, p_T$ cuts	All cuts	$y, p_T$ cuts	All cuts	
250	25253	25082	21.804	0.3797	16938	29.52	0.021
500	171.34	148.69	21.804	0.047	16938	3.74	3.514
750	14.508	12.221	21.804	0.0097	16938	0.997	42.752
1000	2.314	1.9214	21.804	0.0027	16938	0.259	271.92
1250	0.484	0.399	21.804	0.0011	16938	0.048	1310





$M_{b_2}$ (GeV)	QCD background (in fb)					
	$bjjbZ$		$bbjbZ$		$bbbbZ$	
	$y, p_T$ cuts	All cuts	$y, p_T$ cuts	All cuts	$y, p_T$ cuts	All cuts
250	16790	27.304	255.41	2.7	81.01	1.92
500	16790	3.513	255.41	0.256	81.01	0.194
750	16790	0.958	255.41	0.031	81.01	0.057
1000	16790	0.2514	255.41	0.0052	81.01	0.008

# $b'$ Signature (Model Independent)



Benchmark Points (Model I):

$M_{b_2}$ (GeV)	250	500	750	1000	1250	1500
$\kappa_{b_2 b Z}^L$	0.185	0.121	0.084	0.064	0.051	0.043
$\kappa_{b_2 t W}$	0.322	0.161	0.107	0.080	0.064	0.054
$\kappa_{h b_L b_2 R}$	0.714	0.937	0.972	0.985	0.990	0.993
$M_{b_2}$ (GeV)	1750	2000	2250	2500	2750	3000
$\kappa_{b_2 b Z}^L$	0.037	0.032	0.029	0.026	0.024	0.022
$\kappa_{b_2 t W}$	0.046	0.040	0.036	0.032	0.029	0.027
$\kappa_{h b_L b_2 R}$	0.995	0.996	0.997	0.998	0.998	0.998

Warped model  $b'$  :  $\Gamma$  and BR

# $b'$ Single Resonant II Details

$$pp \rightarrow b'Z \rightarrow bZZ \rightarrow bjj\ell^+\ell^-$$

$M_{b'}$ (GeV)	signal $\sigma_s$ (in fb)		background $\sigma_b$ (in fb)				$\mathcal{L}^{\text{SemiLep}}$ ( $\text{fb}^{-1}$ )
	$bjjZ$		$(bjjZ)_{EW}$		$(bjjZ)_{QCD}$		
	Primary cuts	all cuts	Primary cuts	all cuts	Primary cuts	all cuts	
250	1017.66	995.86	77.03	10.33	7853.02	867.82	0.66
500	16.84	15.50	8.81	0.68	419.75	14.11	45.94
750	1.26	1.14	1.85	0.10	56.26	0.86	551.26
1000	0.14	0.12	0.47	0.01	12.38	0.05	3399.67

$M_{b'}$ (GeV)	QCD background (in fb)		
	$bjjZ$	$bjbZ$	$bbbZ$
250	546.36	634.32	17.19
500	10.14	7.76	0.35
750	0.52	0.66	0.03
1000	0.02	0.06	0.002