EWSB after the Higgs

M. Schmaltz · Pheno 2014



Fermilab 95-759


t
d
c
s
u
b
γ
g
H
W
Z
τ
ν_e
μ
ν_μ
e
ν_τ
Fermions
Matter
Quarks
Lepton
Bosons
Force Carriers
Gauge Bosons
Higgs Boson
PARTICLEFEVER

STANDARD MODEL T-SHIRT

Show your love of particle physics by wearing this comfy, American Apparel 100% combed and ring-spun cotton black t-shirt featuring all of the known particles in the Standard Model – the current theory that describes all of particle physics. One of the sleeves is branded with Particle Fever.

$24.99

Buy Now ▸

SM Higgs

$$V_{SM} = -\mu^2 |H|^2 + \lambda |H|^4$$

$(89\ \text{GeV})^2$

~ 0.13



√s = 7 TeV, L ≤ 5.1 fb-1 √s = 8 TeV, L ≤ 19.7 fb-1
CMS Preliminary
Individual Results
V H → bb arXiv:1310.3687
μ(mH=125.0 GeV) = 1.0 ± 0.5
H → ττ arXiv:1401.5041
μ(mH=125.0 GeV) = 0.78 ± 0.27
H → γγ HIG-13-001
μ(mH=125.0 GeV) = 0.78 ± 0.27
H → WW arXiv:1312.1129
μ(mH=125.6 GeV) = 0.72 ± 0.19
H → ZZ arXiv:1312.5353
μ(mH=125.6 GeV) = 0.93 ± 0.27
0
0.5
1
1.5
2
Best fit σ/σSM


ATLAS Prelim.
m_H = 125.5 GeV
σ(stat.)
σ(sys inc. theory)
σ(theory)
Total uncertainty
± 1σ
± 2σ
H → γγ
$\frac{\mu_{VBF+VH}}{\mu_{ggF+ttH}} = 1.2^{+0.8}_{-0.6}$
+ 0.7
− 0.5
+ 0.4
− 0.2
+ 0.2
− 0.2
H → ZZ* → 4l
$\frac{\mu_{VBF+VH}}{\mu_{ggF+ttH}} = 0.6^{+2.4}_{-0.9}$
+ 2.3
− 0.9
+ 0.6
− 0.2
+ 0.3
− 0.2
H → WW* → lνlν
$\frac{\mu_{VBF+VH}}{\mu_{ggF+ttH}} = 1.8^{+1.9}_{-1.0}$
+ 1.4
− 0.9
+ 1.3
− 0.4
+ 0.5
− 0.2
H → ττ
$\frac{\mu_{VBF+VH}}{\mu_{ggF+ttH}} = 1.7^{+\infty}_{-1.2}$
+ 5.3
− 1.0
+ ∞
− 0.6
+ ∞
− 0.3
Combined
$\frac{\mu_{VBF+VH}}{\mu_{ggF+ttH}} = 1.4^{+0.7}_{-0.5}$
+ 0.5
− 0.4
+ 0.4
− 0.2
+ 0.2
− 0.1
2 σ
1 σ
0
1
2
3
4
5
$\sqrt{s}$ = 7 TeV $\int Ldt$ = 4.6-4.8 fb^{-1}
$\sqrt{s}$ = 8 TeV $\int Ldt$ = 20.3 fb^{-1}
$\mu_{VBF+VH} / \mu_{ggF+ttH}$

alternatives ?

2 Higgs doublets (type II)

H_u H_d



2HDM Type II
ATLAS Preliminary
Obs. 95% CL
Best fit
Exp. 95% CL
SM
√s = 7 TeV: ∫Ldt = 4.6-4.8 fb-1
√s = 8 TeV: ∫Ldt = 20.3 fb-1
Combined h → γγ,ZZ*,WW*
h → ττ,bb̄
tan β
10
4
3
2
1
0.4
0.3
0.2
0.1
-1
-0.8
-0.6
-0.4
-0.2
0
0.2
0.4
0.6
0.8
1
cos(β-α)

Higgs Effective Theory

$$\mathcal{L} \sim \mu^2 H^2 - \lambda H^4 + \sum_i \frac{O_i(H)}{\Lambda_i^2}$$

dim-6

Dimension 6 operators

- $(H^\dagger D_\mu H)^2 \xrightarrow{\text{precision EW}} \Lambda_T \gtrsim 5\text{ TeV}$
- $W^a_{\mu\nu} D^\mu H^\dagger \sigma^a D^\nu H \xrightarrow{\text{LHC Higgs}} \Lambda_{HW} \gtrsim 2\text{ TeV}$
- $\frac{g'^2}{16\pi^2} H^\dagger H B_{\mu\nu} B^{\mu\nu} \xrightarrow{h \to \gamma\gamma} \Lambda_\gamma \gtrsim \text{TeV}$

$\Big]$ complementary

no sign of new physics

Higgs Naturalness & New Physics at scale Λ

$$\mathcal{L} \sim \mu^2 H^2 - \lambda H^4 + \sum_{\dim 6} \frac{O_i}{\Lambda_i^2}$$

$$\mu^2 \sim \frac{g^2}{16\pi^2} \Lambda^2$$

dimensional analysis



$$\delta m^2_{Higgs} \sim \frac{g^2}{16\pi^2}\Lambda^2$$

from all New Physics which couples to the Higgs.

dimensional analysis fails dramatically

↳ missing something important!

What are we missing?

A. symmetries $\longrightarrow$ partners

B. there are no high scales

C. anthropics

D. other

A. symmetries



extend SM above scale M to include symmetry which protects $\mu^2 H^2$

- SUSY
- Shift symmetry

$\Big\}$ partners at $M \lesssim \frac{4\pi}{g} m_{Higgs}$

Where are the partners !!!?

SUSY

$\tilde{t}, \tilde{g}, \tilde{W}, \tilde{H}$

Little Higgs

T', W', B'

ATLAS SUSY Searches* - 95% CL Lower Limits

Status: Moriond 2014

ATLAS Preliminary

$\int \mathcal{L}\,dt = (4.6 - 22.9)\ \mathrm{fb}^{-1}$ $\sqrt{s}$ = 7, 8 TeV

	Model	e, μ, τ, γ	Jets	E_T^{miss}	$\int \mathcal{L}\,dt[\mathrm{fb}^{-1}]$	Mass limit			Reference
Inclusive Searches	MSUGRA/CMSSM	0	2-6 jets	Yes	20.3	$\tilde{q}, \tilde{g}$	1.7 TeV	m($\tilde{q}$)=m($\tilde{g}$)	ATLAS-CONF-2013-047
	MSUGRA/CMSSM	1 e, μ	3-6 jets	Yes	20.3	$\tilde{g}$	1.2 TeV	any m($\tilde{q}$)	ATLAS-CONF-2013-062
	MSUGRA/CMSSM	0	7-10 jets	Yes	20.3	$\tilde{g}$	1.1 TeV	any m($\tilde{q}$)	1308.1841
	$\tilde{q}\tilde{q}$, $\tilde{q}\to q\tilde{\chi}_1^0$	0	2-6 jets	Yes	20.3	$\tilde{q}$	740 GeV	m($\tilde{\chi}_1^0$)=0 GeV	ATLAS-CONF-2013-047
	$\tilde{g}\tilde{g}$, $\tilde{g}\to q\bar{q}\tilde{\chi}_1^0$	0	2-6 jets	Yes	20.3	$\tilde{g}$	1.3 TeV	m($\tilde{\chi}_1^0$)=0 GeV	ATLAS-CONF-2013-047
	$\tilde{g}\tilde{g}$, $\tilde{g}\to qq\tilde{\chi}_1^{\pm}\to qqW^{\pm}\tilde{\chi}_1^0$	1 e, μ	3-6 jets	Yes	20.3	$\tilde{g}$	1.18 TeV	m($\tilde{\chi}_1^0$)<200 GeV, m($\tilde{\chi}^{\pm}$)=0.5(m($\tilde{\chi}_1^0$)+m($\tilde{g}$))	ATLAS-CONF-2013-062
	$\tilde{g}\tilde{g}$, $\tilde{g}\to qq(\ell\ell/\ell\nu/\nu\nu)\tilde{\chi}_1^0$	2 e, μ	0-3 jets	-	20.3	$\tilde{g}$	1.12 TeV	m($\tilde{\chi}_1^0$)=0 GeV	ATLAS-CONF-2013-089
	GMSB ($\tilde{\ell}$ NLSP)	2 e, μ	2-4 jets	Yes	4.7	$\tilde{g}$	1.24 TeV	tanβ<15	1208.4688
	GMSB ($\tilde{\ell}$ NLSP)	1-2 τ	0-2 jets	Yes	20.7	$\tilde{g}$	1.4 TeV	tanβ >18	ATLAS-CONF-2013-026
	GGM (bino NLSP)	2 γ	-	Yes	20.3	$\tilde{g}$	1.28 TeV	m($\tilde{\chi}_1^0$)>50 GeV	ATLAS-CONF-2014-001
	GGM (wino NLSP)	1 $e, \mu + \gamma$	-	Yes	4.8	$\tilde{g}$	619 GeV	m($\tilde{\chi}_1^0$)>50 GeV	ATLAS-CONF-2012-144
	GGM (higgsino-bino NLSP)	γ	1 b	Yes	4.8	$\tilde{g}$	900 GeV	m($\tilde{\chi}_1^0$)>220 GeV	1211.1167
	GGM (higgsino NLSP)	2 e, μ (Z)	0-3 jets	Yes	5.8	$\tilde{g}$	690 GeV	m($\tilde{H}$)>200 GeV	ATLAS-CONF-2012-152
	Gravitino LSP	0	mono-jet	Yes	10.5	$F^{1/2}$ scale	645 GeV	m($\tilde{g}$)>10^{-4} eV	ATLAS-CONF-2012-147
3^{rd} gen. $\tilde{g}$ med.	$\tilde{g}\to b\bar{b}\tilde{\chi}_1^0$	0	3 b	Yes	20.1	$\tilde{g}$	1.2 TeV	m($\tilde{\chi}_1^0$)<600 GeV	ATLAS-CONF-2013-061
	$\tilde{g}\to t\bar{t}\tilde{\chi}_1^0$	0	7-10 jets	Yes	20.3	$\tilde{g}$	1.1 TeV	m($\tilde{\chi}_1^0$) <350 GeV	1308.1841
	$\tilde{g}\to t\bar{t}\tilde{\chi}_1^0$	0-1 e, μ	3 b	Yes	20.1	$\tilde{g}$	1.34 TeV	m($\tilde{\chi}_1^0$)<400 GeV	ATLAS-CONF-2013-061
	$\tilde{g}\to b\bar{t}\tilde{\chi}_1^+$	0-1 e, μ	3 b	Yes	20.1	$\tilde{g}$	1.3 TeV	m($\tilde{\chi}_1^0$)<300 GeV	ATLAS-CONF-2013-061
3^{rd} gen. squarks direct production	$\tilde{b}_1\tilde{b}_1$, $\tilde{b}_1\to b\tilde{\chi}_1^0$	0	2 b	Yes	20.1	$\tilde{b}_1$	100-620 GeV	m($\tilde{\chi}_1^0$)<90 GeV	1308.2631
	$\tilde{b}_1\tilde{b}_1$, $\tilde{b}_1\to t\tilde{\chi}_1^{\pm}$	2 e, μ (SS)	0-3 b	Yes	20.7	$\tilde{b}_1$	275-430 GeV	m($\tilde{\chi}_1^{\pm}$)=2 m($\tilde{\chi}_1^0$)	ATLAS-CONF-2013-007
	$\tilde{t}_1\tilde{t}_1$(light), $\tilde{t}_1\to b\tilde{\chi}_1^{\pm}$	1-2 e, μ	1-2 b	Yes	4.7	$\tilde{t}_1$	110-167 GeV	m($\tilde{\chi}_1^0$)=55 GeV	1208.4305, 1209.2102
	$\tilde{t}_1\tilde{t}_1$(light), $\tilde{t}_1\to Wb\tilde{\chi}_1^0$	2 e, μ	0-2 jets	Yes	20.3	$\tilde{t}_1$	130-210 GeV	m($\tilde{\chi}_1^0$) =m($\tilde{t}_1$)-m(W)-50 GeV, m($\tilde{t}_1$)<<m($\tilde{\chi}_1^{\pm}$)	1403.4853
	$\tilde{t}_1\tilde{t}_1$(medium), $\tilde{t}_1\to t\tilde{\chi}_1^0$	2 e, μ	2 jets	Yes	20.3	$\tilde{t}_1$	215-530 GeV	m($\tilde{\chi}_1^0$)=1 GeV	1403.4853
	$\tilde{t}_1\tilde{t}_1$(medium), $\tilde{t}_1\to b\tilde{\chi}_1^{\pm}$	0	2 b	Yes	20.1	$\tilde{t}_1$	150-580 GeV	m($\tilde{\chi}_1^0$)<200 GeV, m($\tilde{\chi}_1^{\pm}$)-m($\tilde{\chi}_1^0$)=5 GeV	1308.2631
	$\tilde{t}_1\tilde{t}_1$(heavy), $\tilde{t}_1\to t\tilde{\chi}_1^0$	1 e, μ	1 b	Yes	20.7	$\tilde{t}_1$	200-610 GeV	m($\tilde{\chi}_1^0$)=0 GeV	ATLAS-CONF-2013-037
	$\tilde{t}_1\tilde{t}_1$(heavy), $\tilde{t}_1\to t\tilde{\chi}_1^0$	0	2 b	Yes	20.5	$\tilde{t}_1$	320-660 GeV	m($\tilde{\chi}_1^0$)=0 GeV	ATLAS-CONF-2013-024
	$\tilde{t}_1\tilde{t}_1$, $\tilde{t}_1\to c\tilde{\chi}_1^0$	0	mono-jet/c-tag	Yes	20.3	$\tilde{t}_1$	90-200 GeV	m($\tilde{t}_1$)-m($\tilde{\chi}_1^0$)<85 GeV	ATLAS-CONF-2013-068
	$\tilde{t}_1\tilde{t}_1$(natural GMSB)	2 e, μ (Z)	1 b	Yes	20.3	$\tilde{t}_1$	150-580 GeV	m($\tilde{\chi}_1^0$)>150 GeV	1403.5222
	$\tilde{t}_2\tilde{t}_2$, $\tilde{t}_2\to\tilde{t}_1 + Z$	3 e, μ (Z)	1 b	Yes	20.3	$\tilde{t}_2$	290-600 GeV	m($\tilde{\chi}_1^0$)<200 GeV	1403.5222
EW direct	$\tilde{\ell}_{L,R}\tilde{\ell}_{L,R}$, $\tilde{\ell}\to\ell\tilde{\chi}_1^0$	2 e, μ	0	Yes	20.3	$\tilde{\ell}$	90-325 GeV	m($\tilde{\chi}_1^0$)=0 GeV	1403.5294
	$\tilde{\chi}_1^+\tilde{\chi}_1^-$, $\tilde{\chi}_1^+\to\tilde{\ell}\nu(\ell\tilde{\nu})$	2 e, μ	0	Yes	20.3	$\tilde{\chi}_1^{\pm}$	140-465 GeV	m($\tilde{\chi}_1^0$)=0 GeV, m($\tilde{\ell}, \tilde{\nu}$)=0.5(m($\tilde{\chi}_1^{\pm}$)+m($\tilde{\chi}_1^0$))	1403.5294
	$\tilde{\chi}_1^+\tilde{\chi}_1^-$, $\tilde{\chi}_1^+\to\tilde{\tau}\nu(\tau\tilde{\nu})$	2 τ	-	Yes	20.7	$\tilde{\chi}_1^{\pm}$	180-330 GeV	m($\tilde{\chi}_1^0$)=0 GeV, m($\tilde{\tau}, \tilde{\nu}$)=0.5(m($\tilde{\chi}_1^{\pm}$)+m($\tilde{\chi}_1^0$))	ATLAS-CONF-2013-028
	$\tilde{\chi}_1^{\pm}\tilde{\chi}_2^0\to\tilde{\ell}_L\nu\tilde{\ell}_L\ell(\tilde{\nu}\nu)$, $\ell\tilde{\nu}\tilde{\ell}_L\ell(\tilde{\nu}\nu)$	3 e, μ	0	Yes	20.3	$\tilde{\chi}_1^{\pm}, \tilde{\chi}_2^0$	700 GeV	m($\tilde{\chi}_1^{\pm}$)=m($\tilde{\chi}_2^0$), m($\tilde{\chi}_1^0$)=0, m($\tilde{\ell}, \tilde{\nu}$)=0.5(m($\tilde{\chi}_1^{\pm}$)+m($\tilde{\chi}_1^0$))	1402.7029
	$\tilde{\chi}_1^{\pm}\tilde{\chi}_2^0\to W\tilde{\chi}_1^0 Z\tilde{\chi}_1^0$	2-3 e, μ	0	Yes	20.3	$\tilde{\chi}_1^{\pm}, \tilde{\chi}_2^0$	420 GeV	m($\tilde{\chi}_1^{\pm}$)=m($\tilde{\chi}_2^0$), m($\tilde{\chi}_1^0$)=0, sleptons decoupled	1403.5294, 1402.7029
	$\tilde{\chi}_1^{\pm}\tilde{\chi}_2^0\to W\tilde{\chi}_1^0 h\tilde{\chi}_1^0$	1 e, μ	2 b	Yes	20.3	$\tilde{\chi}_1^{\pm}, \tilde{\chi}_2^0$	285 GeV	m($\tilde{\chi}_1^{\pm}$)=m($\tilde{\chi}_2^0$), m($\tilde{\chi}_1^0$)=0, sleptons decoupled	ATLAS-CONF-2013-093
Long-lived particles	Direct $\tilde{\chi}_1^+\tilde{\chi}_1^-$ prod., long-lived $\tilde{\chi}_1^{\pm}$	Disapp. trk	1 jet	Yes	20.3	$\tilde{\chi}_1^{\pm}$	270 GeV	m($\tilde{\chi}_1^{\pm}$)-m($\tilde{\chi}_1^0$)=160 MeV, τ($\tilde{\chi}_1^{\pm}$)=0.2 ns	ATLAS-CONF-2013-069
	Stable, stopped $\tilde{g}$ R-hadron	0	1-5 jets	Yes	22.9	$\tilde{g}$	832 GeV	m($\tilde{\chi}_1^0$)=100 GeV, 10 μs<τ($\tilde{g}$)<1000 s	ATLAS-CONF-2013-057
	GMSB, stable $\tilde{\tau}$, $\tilde{\chi}_1^0\to\tilde{\tau}(\tilde{e}, \tilde{\mu})+\tau(e, \mu)$	1-2 μ	-	-	15.9	$\tilde{\chi}_1^0$	475 GeV	10<tanβ<50	ATLAS-CONF-2013-058
	GMSB, $\tilde{\chi}_1^0\to\gamma\tilde{G}$, long-lived $\tilde{\chi}_1^0$	2 γ	-	Yes	4.7	$\tilde{\chi}_1^0$	230 GeV	0.4<τ($\tilde{\chi}_1^0$)<2 ns	1304.6310
	$\tilde{q}\tilde{q}$, $\tilde{\chi}_1^0\to qq\mu$ (RPV)	1 μ, displ. vtx	-	-	20.3	$\tilde{q}$	1.0 TeV	1.5 <$c\tau$<156 mm, BR(μ)=1, m($\tilde{\chi}_1^0$)=108 GeV	ATLAS-CONF-2013-092
RPV	LFV $pp\to\tilde{\nu}_\tau + X, \tilde{\nu}_\tau\to e + \mu$	2 e, μ	-	-	4.6	$\tilde{\nu}_\tau$	1.61 TeV	λ'_{311}=0.10, λ_{132}=0.05	1212.1272
	LFV $pp\to\tilde{\nu}_\tau + X, \tilde{\nu}_\tau\to e(\mu) + \tau$	1 $e, \mu + \tau$	-	-	4.6	$\tilde{\nu}_\tau$	1.1 TeV	λ'_{311}=0.10, $\lambda_{1(2)33}$=0.05	1212.1272
	Bilinear RPV CMSSM	1 e, μ	7 jets	Yes	4.7	$\tilde{q}, \tilde{g}$	1.2 TeV	m($\tilde{q}$)=m($\tilde{g}$), $c\tau_{LSP}$<1 mm	ATLAS-CONF-2012-140
	$\tilde{\chi}_1^+\tilde{\chi}_1^-$, $\tilde{\chi}_1^+\to W\tilde{\chi}_1^0, \tilde{\chi}_1^0\to ee\tilde{\nu}_\mu, e\mu\tilde{\nu}_e$	4 e, μ	-	Yes	20.7	$\tilde{\chi}_1^{\pm}$	760 GeV	m($\tilde{\chi}_1^0$)>300 GeV, λ_{121}>0	ATLAS-CONF-2013-036
	$\tilde{\chi}_1^+\tilde{\chi}_1^-$, $\tilde{\chi}_1^+\to W\tilde{\chi}_1^0, \tilde{\chi}_1^0\to\tau\tau\tilde{\nu}_e, e\tau\tilde{\nu}_\tau$	3 $e, \mu + \tau$	-	Yes	20.7	$\tilde{\chi}_1^{\pm}$	350 GeV	m($\tilde{\chi}_1^0$)>80 GeV, λ_{133}>0	ATLAS-CONF-2013-036
	$\tilde{g}\to qqq$	0	6-7 jets	-	20.3	$\tilde{g}$	916 GeV	BR(t)=BR(b)=BR(c)=0%	ATLAS-CONF-2013-091
	$\tilde{g}\to\tilde{t}_1 t$, $\tilde{t}_1\to bs$	2 e, μ (SS)	0-3 b	Yes	20.7	$\tilde{g}$	880 GeV		ATLAS-CONF-2013-007
Other	Scalar gluon pair, sgluon$\to q\bar{q}$	0	4 jets	-	4.6	sgluon	100-287 GeV	incl. limit from 1110.2693	1210.4826
	Scalar gluon pair, sgluon$\to t\bar{t}$	2 e, μ (SS)	2 b	Yes	14.3	sgluon	350-800 GeV		ATLAS-CONF-2013-051
	WIMP interaction (D5, Dirac χ)	0	mono-jet	Yes	10.5	M* scale	704 GeV	m(χ)<80 GeV, limit of<687 GeV for D8	ATLAS-CONF-2012-147

10^{-1} 1 Mass scale [TeV]

$\sqrt{s}$ = 7 TeV full data | $\sqrt{s}$ = 8 TeV partial data | $\sqrt{s}$ = 8 TeV full data

*Only a selection of the available mass limits on new states or phenomena is shown. All limits quoted are observed minus 1σ theoretical signal cross section uncertainty.


Leptoquarks
LQ1(ej) x2
LQ1(ej)+LQ1(νj)
LQ2(μj) x2
LQ2(μj)+LQ2(νj)
LQ3(νb) x2
LQ3(τb) x2
LQ3(τt) x2
0 1 2 3 4
Long-Lived Particles
stopped gluino (cloud)
stopped stop (cloud)
HSCP gluino (cloud)
HSCP stop (cloud)
q=2/3e HSCP
q=3e HSCP
neutralino, ctau=25cm, ECAL time
0 1 2 3 4
RS Gravitons
RS1(γγ), k=0.1
RS1(ee,uu), k=0.1
RS1(jj), k=0.1
RS1(WW→4j), k=0.1
RS1(ZZ→4j), k=0.1
bulk RS(ZZ→lljj), k=0.5
0 1 2 3 4
Dark Matter
j+MET, SI DM=100 GeV, Λ
j+MET, SD DM=100 GeV, Λ
γ+MET, SI DM=100 GeV, Λ
γ+MET, SD DM=100 GeV, Λ
l+MET, ξ=+1, SI DM=100 GeV, Λ
l+MET, ξ=+1, SD DM=100 GeV, Λ
l+MET, ξ=-1, SI DM=100 GeV, Λ
l+MET, ξ=-1, SD DM=100 GeV, Λ
0 1 2 3 4
Heavy Gauge Bosons
SSM Z'(ττ)
SSM Z'(jj)
SSM Z'(bb)
SSM Z'(ee)+Z'(μμ)
SSM W'(jj)
SSM W'(lν)
SSM W'(WZ→lνll)
SSM W'(WZ→4j)
0 1 2 3 4
Compositeness
dijets, Λ+ LL/RR
dijets, Λ- LL/RR
dimuons, Λ+ LLIM
dimuons, Λ- LLIM
single e, Λ HnCM
single μ, Λ HnCM
inclusive jets, Λ+
inclusive jets, Λ-
0 3 6 9 12 15
Excited Fermions
e* (M=Λ)
μ* (M=Λ)
q* (qg)
q* (qγ)
b*
0 1 2 3 4
Large Extra Dimensions
ADD (γγ), nED=4, MS
ADD (ee,μμ), nED=4, MS
ADD (j+MET), nED=4, MD
ADD (γ+MET), nED=4, MD
QBH, nED=4, MD=4 TeV
NR BH, nED=4, MD=4 TeV
Jet Extinction Scale
String Scale (jj)
0 3 6 9 12 15
Multijet Resonances
coloron(jj) x2
coloron(4j) x2
gluino(3j) x2
gluino(jjb) x2
0 1 2 3 4
CMS Preliminary

What are we missing?

A. symmetries

B. there are no high scales

$\longleftrightarrow$ UV physics is scale invariant

C. anthropics

D. other

B. scale invariant UV

if $m_{Higgs} \sim m_{top}$ is the largest scale in Nature/the SM then m^2_{Higgs} is automatically natural !

B. scale invariant UV

$$\frac{\lambda_t^2}{16\pi^2}\left(\cancel{\Lambda^2} + m_{top}^2 \qquad \right)$$

by scale invariance

B. scale non-invariance in the UV



"maybe gravity is different"

→ ignore gravity, M_{pl}.

- 't Hooft 1980
- Dubovsky, Gorbenko, Mirbabayi 2013
- ...

B. Scale invariant UV



B. scale non-invariant UV



B. Scale non-invariant UV



B. scale non-invariant UV



$$\frac{g'^4}{16\pi^2} M^2_{\text{transition}}$$

- even if there are no particles at the transition!
  (Banks-Zaks)
- $\overline{MS}$ and dim. reg. doesn't help

Marques Tavares, Schmaltz, Skiba
1308.0025

B. scale invariant UV



- naturalness requires $M_{trans} \lesssim \frac{4\pi}{g'^2} M_{Higgs}$
- experiment? charged vector bosons @ few TeV
  W', Z'

DGM 2013, MTSS 2013

What are we missing?

A. } symmetries ⟶ particles at LHC
B. }

last call for naturalness!

C. anthropics (Particle Fever, Nima)

D. other

D. other

Phillip v. Jolly (1874)

"almost everything is already discovered, and all that remains is to fill a few holes"

Michelson (1894)

"future discoveries must be looked for in the 6th decimal"

Higgs factory?

D. other

Rayleigh-Jeans law for black body radiation (1900)

$E(\omega) \propto \omega^2 d\omega$ "ultraviolet catastrophe"

$\Longrightarrow$ QM revolution

D. other

$\delta m^2_{Higgs} \propto \Lambda^2$ is our ultraviolet catastrophe

what is our Revolution?