

Physics with b -baryons at LHCb

J. McCarthy on behalf of the LHCb collaboration

University of Birmingham

06/05/2015


UNIVERSITY OF
BIRMINGHAM



The 2014 Phenomenology Symposium
University of Pittsburgh, 5-7 May 2014

Introduction

- Physics with b -baryons relatively unexplored.
- Baryons have non-zero spin
 - Probe helicity structure of HQE Hamiltonian
- Need precision measurements of mass, lifetime and branching fractions.
- CP measurements interesting.
 - Due interference between tree and loop diagrams.
 - Self tagging decays.
 - No CP asymmetries observed.
- Focus on results of baryon decays and Ξ_b^0 measurements.




Contents


- 1 Observation of the $\Lambda_b^0 \rightarrow J/\psi p\pi^-$ decay.
- 2 $\Lambda_b^0 \rightarrow \Lambda_c^+ D^-$ and $\Lambda_b^0 \rightarrow \Lambda_c^+ D_s^-$ decays.
- 3 $\Lambda_b^0(\Xi_b^0) \rightarrow D^0 ph^-$ and $\Lambda_b^0(\Xi_b^0) \rightarrow \Lambda_c^+ h^-$ decays.
- 4 Mass and lifetime measurements of Ξ_b^0
- 5 Λ_b^0 production and $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ branching fraction measurements.

$\Lambda_b^0 \rightarrow J/\psi p\pi^-$ observation. LHCb-PAPER-2014-020

- Search for $\Lambda_b^0 \rightarrow J/\psi p\pi^-$ in 3 fb^{-1} data.
- $\Lambda_b^0 \rightarrow J/\psi pK^-$ decay as control channel.
- Loose selection followed by PID and Neural Network.



$$\begin{aligned} \frac{\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi p\pi^-)}{\mathcal{B}(\Lambda_b^0 \rightarrow J/\psi pK^-)} &= \frac{N(\Lambda_b^0 \rightarrow J/\psi p\pi^-)}{N(\Lambda_b^0 \rightarrow J/\psi pK^-)} \times \frac{\eta(\Lambda_b^0 \rightarrow J/\psi pK^-)}{\eta(\Lambda_b^0 \rightarrow J/\psi p\pi^-)} \\ &= 0.0824 \pm 0.0025 \text{ (stat)} \pm 0.0042 \text{ (syst)} \end{aligned}$$



\mathcal{A}^{CP} measurements.

LHCb-PAPER-2014-020

- Measure raw asymmetries, $\mathcal{A}^{\text{raw}} = \frac{N(\Lambda_b^0) - N(\bar{\Lambda}_b^0)}{N(\Lambda_b^0) + N(\bar{\Lambda}_b^0)}$

$$\mathcal{A}^{\text{raw}}(\Lambda_b^0 \rightarrow J/\psi p\pi^-) = (+7.9 \pm 2.2)\%$$

$$\mathcal{A}^{\text{raw}}(\Lambda_b^0 \rightarrow J/\psi pK^-) = (+1.1 \pm 0.7)\%$$

- Related to CP asymmetry: $\mathcal{A}^{\text{raw}} = \mathcal{A}^{CP} + \mathcal{A}^{\text{prod}}(\Lambda_b^0) - \mathcal{A}^{\text{det}}(\pi/K) + \mathcal{A}^{\text{det}}(p)$
- Calculate difference in decay modes:

$$\begin{aligned}\Delta\mathcal{A}^{CP} &= \mathcal{A}^{CP}(J/\psi p\pi^-) - \mathcal{A}^{CP}(J/\psi pK^-) \\ &= \mathcal{A}^{\text{raw}}(J/\psi p\pi^-) - \mathcal{A}^{\text{raw}}(J/\psi pK^-) + \mathcal{A}^{\text{det}}(\pi) - \mathcal{A}^{\text{det}}(K)\end{aligned}$$

- Find $\mathcal{A}^{\text{det}}(\pi) - \mathcal{A}^{\text{det}}(K)$ from $\bar{B}^0 \rightarrow J/\psi \bar{K}^{*0}$ decay.

$$\begin{aligned}\mathcal{A}^{\text{raw}}(\bar{B}^0 \rightarrow J/\psi \bar{K}^{*0}) &= \mathcal{A}^{CP}(\bar{B}^0 \rightarrow J/\psi \bar{K}^{*0}) + \mathcal{A}^{\text{prod}}(B^0) + \mathcal{A}^{\text{det}}(\pi) - \mathcal{A}^{\text{det}}(K) \\ &\approx \mathcal{A}^{\text{det}}(\pi) - \mathcal{A}^{\text{det}}(K)\end{aligned}$$

Preliminary

$$\Delta\mathcal{A}^{CP} = (+5.7 \pm 2.3(\text{stat}) \pm 1.2(\text{syst}))\%$$



Contents

- 1 Observation of the $\Lambda_b^0 \rightarrow J/\psi p\pi^-$ decay.
- 2 $\Lambda_b^0 \rightarrow \Lambda_c^+ D^-$ and $\Lambda_b^0 \rightarrow \Lambda_c^+ D_s^-$ decays.
- 3 $\Lambda_b^0(\Xi_b^0) \rightarrow D^0 ph^-$ and $\Lambda_b^0(\Xi_b^0) \rightarrow \Lambda_c^+ h^-$ decays.
- 4 Mass and lifetime measurements of Ξ_b^0
- 5 Λ_b^0 production and $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ branching fraction measurements.

Beauty hadron decays into pairs of charm hadrons

arXiv:1403.3606

- Search for $\Lambda_b^0 \rightarrow \Lambda_c^+ D_s^-$ and $\Lambda_b^0 \rightarrow \Lambda_c^+ D^-$ decays.
- Reconstruct resonances: $D^+ \rightarrow K^-\pi^+\pi^+$, $D_s^+ \rightarrow K^-K^+\pi^+$, $\Lambda_c^+ \rightarrow pK^-\pi^+$
- First observation of these decays:



$$\frac{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ D^-)}{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ D_s^-)} = 0.042 \pm 0.003 \text{ (stat)} \pm 0.003 \text{ (syst)}$$

Mass measurement [arXiv:1403.3606]

- Measure relative to high statistics channel $\bar{B}^0 \rightarrow D^+ D_s^-$
- Double ratio removes any dependence on production fractions.

$$\left[\frac{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ D_s^-)}{\mathcal{B}(\bar{B}^0 \rightarrow D^+ D_s^-)} \right] / \left[\frac{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)}{\mathcal{B}(\bar{B}^0 \rightarrow D^+ \pi^-)} \right] = 0.96 \pm 0.02 \text{ (stat)} \pm 0.06 \text{ (syst)}$$

$$\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ D_s^-) = (1.1 \pm 0.1) \times 10^{-2}$$

$$\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ D^-) = (4.7 \pm 0.6) \times 10^{-4}$$

- Kinematics similar enough to measure mass difference
 - $[m(\Lambda_b^0) - m(\Lambda_c^+) - m(D_s^-)] - [m(B^0) - m(D^+) - m(D_s^-)]$ is small.
 - Small uncertainty on momentum scale ($\approx 0.03\%$).

$$m(\Lambda_b^0) - m(B^0) = 339.72 \pm 0.24 \text{ (stat)} \pm 0.18 \text{ (syst)} \text{ MeV}/c^2$$

$$m(\Lambda_b^0) = 5619.36 \pm 0.26 \text{ MeV}/c^2$$

- Averaged with LHCb $\Lambda_b^0 \rightarrow J/\psi \Lambda$ result [Phys. Rev. Lett. 110 (2013) 182001]

Contents


- 1 Observation of the $\Lambda_b^0 \rightarrow J/\psi p\pi^-$ decay.
- 2 $\Lambda_b^0 \rightarrow \Lambda_c^+ D^-$ and $\Lambda_b^0 \rightarrow \Lambda_c^+ D_s^-$ decays.
- 3 $\Lambda_b^0(\Xi_b^0) \rightarrow D^0 ph^-$ and $\Lambda_b^0(\Xi_b^0) \rightarrow \Lambda_c^+ h^-$ decays.
- 4 Mass and lifetime measurements of Ξ_b^0
- 5 Λ_b^0 production and $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ branching fraction measurements.

Measurement of $\Lambda_b^0 \rightarrow D^0 p\pi^-$


Phys. Rev. D89 (2014) 032001

- Dalitz analysis would allow extraction of γ
- Reconstruct $D^0 \rightarrow K^-\pi^+$ decay.
- Normalize to $\Lambda_b^0 \rightarrow \Lambda_c^+\pi^-$ with $\Lambda_c^+ \rightarrow pK^-\pi^+$.
- Search in 1 fb^{-1} 2011 data
- Measure Cabibbo favoured $\Lambda_b^0 \rightarrow D^0 p\pi^-$ with loose selection only.

$$\Lambda_b^0 \rightarrow \Lambda_c^+\pi^-$$



$$\Lambda_b^0 \rightarrow D^0 p\pi^-$$




$$\frac{\mathcal{B}(\Lambda_b^0 \rightarrow D^0 p\pi^-)}{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)} \times \frac{\mathcal{B}(D^0 \rightarrow K^-\pi^+)}{\mathcal{B}(\Lambda_c^+ \rightarrow pK^-\pi^+)} = 0.0806 \pm 0.0023 \pm 0.0035$$

Observation of $\Lambda_b^0(\Xi_b^0) \rightarrow D^0 p K^-$

Phys. Rev. D89 (2014) 032001

- Search also for Cabibbo suppressed Λ_b^0 and Ξ_b^0 decays.
- Apply BDT selection to separate signal and background.




$$\frac{\mathcal{B}(\Lambda_b^0 \rightarrow D^0 p K^-)}{\mathcal{B}(\Lambda_b^0 \rightarrow D^0 p \pi^-)} = 0.073 \pm 0.008^{+0.005}_{-0.006}$$

$$\frac{f_{\Xi_b^0} \times \mathcal{B}(\Xi_b^0 \rightarrow D^0 p K^-)}{f_{\Lambda_b^0} \times \mathcal{B}(\Lambda_b^0 \rightarrow D^0 p K^-)} = 0.44 \pm 0.09 \pm 0.06$$


Search for $\Lambda_b^0(\Xi_b^0) \rightarrow \Lambda_c^+ h^-$

Phys. Rev. D89 (2014) 032001

$$\Lambda_b^0(\Xi_b^0) \rightarrow \Lambda_c^+ \pi^-$$





$$\Lambda_b^0(\Xi_b^0) \rightarrow \Lambda_c^+ K^-$$



- First observation of $\Lambda_b^0 \rightarrow \Lambda_c^+ K^- > 10\sigma$

$$\frac{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ K^-)}{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)} = 0.0731 \pm 0.0016 \pm 0.0016$$



- Evidence for $\Xi_b^0 \rightarrow \Lambda_c^+ K^-$ at 3.3σ

ZOOMED

Contents

- 1 Observation of the $\Lambda_b^0 \rightarrow J/\psi p\pi^-$ decay.
- 2 $\Lambda_b^0 \rightarrow \Lambda_c^+ D^-$ and $\Lambda_b^0 \rightarrow \Lambda_c^+ D_s^-$ decays.
- 3 $\Lambda_b^0(\Xi_b^0) \rightarrow D^0 ph^-$ and $\Lambda_b^0(\Xi_b^0) \rightarrow \Lambda_c^+ h^-$ decays.
- 4 Mass and lifetime measurements of Ξ_b^0
- 5 Λ_b^0 production and $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ branching fraction measurements.


Mass of Ξ_b^0

LHCb-PAPER-2014-021

- CDF measured $m(\Xi_b^0) = 5787.8 \pm 5.0 \pm 1.3 \text{ MeV}/c^2$
- Measured using full 3 fb^{-1} data.
- Using $\Xi_b^0 \rightarrow \Xi_c^+ \pi^-$ with $\Xi_c^+ \rightarrow p K^- \pi^+$
- Normalise with $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ with $\Lambda_c^+ \rightarrow p K^- \pi^+$

Phys. Rev. Lett. 107 (2011) 102001

$\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ Preliminary $\Xi_b^0 \rightarrow \Xi_c^+ \pi^-$




$$m(\Xi_b^0) - m(\Lambda_b^0) = 172.44 \pm 0.39 \text{ (stat)} \pm 0.17 \text{ (syst)} \text{ MeV}/c^2$$

Preliminary $m(\Xi_b^0) = 5791.80 \pm 0.39 \text{ (stat)} \pm 0.17 \text{ (syst)} \pm 0.26 \text{ MeV}/c^2$

Lifetime of Ξ_b^0

LHCb-PAPER-2014-021

- First measurement of Ξ_b^0 lifetime.
- Fit to ratio of yields and function of decay time.



- Exponential function, $e^{\beta t}$, where $\beta = \frac{1}{\tau(\Xi_b^0)} - \frac{1}{\tau(\Lambda_b^0)}$

$$\begin{aligned} \frac{\tau(\Xi_b^0)}{\tau(\Lambda_b^0)} &= \frac{1}{1 - \beta \tau(\Lambda_b^0)} \\ &= 1.006 \pm 0.018 \text{ (stat)} \pm 0.010 \text{ (syst)} \end{aligned}$$

Preliminary $\tau(\Xi_b^0) = 1.477 \pm 0.026 \text{ (stat)} \pm 0.014 \text{ (syst)} \pm 0.013(\Lambda_b^0) \text{ ps}$

Contents

- 1 Observation of the $\Lambda_b^0 \rightarrow J/\psi p\pi^-$ decay.
- 2 $\Lambda_b^0 \rightarrow \Lambda_c^+ D^-$ and $\Lambda_b^0 \rightarrow \Lambda_c^+ D_s^-$ decays.
- 3 $\Lambda_b^0(\Xi_b^0) \rightarrow D^0 ph^-$ and $\Lambda_b^0(\Xi_b^0) \rightarrow \Lambda_c^+ h^-$ decays.
- 4 Mass and lifetime measurements of Ξ_b^0
- 5 Λ_b^0 production and $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ branching fraction measurements.



Λ_b^0 production fractions

LHCb-PAPER-2014-004



- $f_{\Lambda_b^0}$ varies as a function of p_T and η
- Measured by LHCb in semi-leptonic decays Phys. Rev. D85 (2012) 032008
- Updated measurement using $\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-$ and $\bar{B}^0 \rightarrow D^+ \pi^-$ decays.
- Fit exponential for p_T dependence, and linear η dependence.

$$\frac{f_{\Lambda_b^0}}{f_d}(X) = \frac{\mathcal{B}(B^0 \rightarrow D^+ \pi^-)}{\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)} \frac{\mathcal{B}(D^+ \rightarrow K^- \pi^+ \pi^+)}{\mathcal{B}(\Lambda_c^+ \rightarrow p K^- \pi^+)} \times \frac{N_{\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-}(X) \epsilon_{B^0 \rightarrow D^+ \pi^-}(X)}{N_{B^0 \rightarrow D^+ \pi^-}(X) \epsilon_{\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-}(X)}$$
$$\equiv S \times R(X)$$

Preliminary



Preliminary



$$\frac{f_{\Lambda_b^0}}{f_d}(p_T) = a + \exp(b + c \times p_T [\text{GeV}/c])$$

 $\leftarrow 40 - 50\%$

$$a = 0.151 \pm 0.016^{+0.024}_{-0.025}$$

$$b = -0.573 \pm 0.040^{+0.101}_{-0.097}$$

$$c = -0.095 \pm 0.007 \pm 0.014$$

$$\frac{f_{\Lambda_b^0}}{f_d}(\eta) = a + b \times (\eta - 3.198)$$

$$a = 0.387 \pm 0.013^{+0.028}_{-0.030}$$

$$b = 0.067 \pm 0.005^{+0.012}_{-0.009}$$

- From scale factor, S , extract absolute branching fraction:

$$\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-) = (4.46 \pm 0.36) \times 10^{-3}$$


- Most precise measurement of a Λ_b^0 branching fraction.

Summary



- First observation of $\Lambda_b^0 \rightarrow J/\psi p\pi^-$
- $\Delta\mathcal{A}^{CP} = \mathcal{A}^{CP}(J/\psi p\pi^-) - \mathcal{A}^{CP}(J/\psi pK^-) = (+5.7 \pm 2.3 \text{ (stat)} \pm 1.2 \text{ (syst)})\%$
- First observation of $\Lambda_b^0 \rightarrow \Lambda_c^+ D^-$ and $\Lambda_b^0 \rightarrow \Lambda_c^+ D_s^-$ decays.
- Used to measure $m(\Lambda_b^0) = 5619.36 \pm 0.26 \text{ MeV}/c^2$
- First observation of $\Lambda_b^0 \rightarrow \Lambda_c^+ K^-$ and evidence for $\Xi_b^0 \rightarrow \Lambda_c^+ K^-$
- Most precise measurements of Ξ_b^0 baryon.
 - $m(\Xi_b^0) = 5791.80 \pm 0.39 \text{ (stat)} \pm 0.17 \text{ (syst)} \pm 0.26 \text{ MeV}/c^2$
 - $\tau(\Xi_b^0) = 1.477 \pm 0.026 \text{ (stat)} \pm 0.014 \text{ (syst)} \pm 0.013(\Lambda_b^0) ps$
- Measured Λ_b^0 production fractions as a function of p_T and η
- Many new observations of baryonic decays shown.
- Expect many more in the years to come!

Back-Up Slides

$\Lambda_b^0 \rightarrow J/\psi p\pi^-$ Dalitz distributions



$\bar{B}^0 \rightarrow D^+ D_s^-$ and $B_s^0 \rightarrow D^+ D_s^-$ invariant mass distributions




$$\frac{\mathcal{B}(B_s^0 \rightarrow D^+ D_s^-)}{\mathcal{B}(\bar{B}^0 \rightarrow D^+ D_s^-)} = 0.038 \pm 0.004 \pm 0.003$$

$$\frac{\mathcal{B}(\bar{B}^0 \rightarrow \Lambda_c^+ \bar{\Lambda}_c^-)}{\mathcal{B}(\bar{B}^0 \rightarrow D^+ D_s^-)} < 0.0022 \text{ at } 95\% \text{ C.L.}$$




$$\frac{\mathcal{B}(B_s^0 \rightarrow \Lambda_c^+ \bar{\Lambda}_c^-)}{\mathcal{B}(B_s^0 \rightarrow D^+ D_s^-)} < 0.30 \text{ at } 95\% \text{ C.L.}$$

Efficiency corrected ratio of yields as a function of p_T





$\Lambda_b^0 \rightarrow D^0 p \pi^-$ Dalitz distributions






$$A_h^0 \rightarrow D^0 p \pi^-$$





$$A_h^0 \rightarrow D^0 p \pi^-$$



$$\Lambda_b^0 \rightarrow D^0 p K^-$$



Ξ_c^+ mass.



$$m(\Xi_c^+) - m(\Lambda_c^+) = 181.51 \pm 0.14 \text{ (stat)} \pm 0.17 \text{ (syst)} \text{ MeV}/c^2$$

Preliminary $m(\Xi_c^+) = 2467.97 \pm 0.14 \text{ (stat)} \pm 0.09 \text{ (syst)} \pm 0.14 \text{ MeV}/c^2$