

Higgs interferometry in the $\gamma\gamma$ channel

Stefan H"och



SLAC National Accelerator Laboratory



in collaboration with Lance Dixon and Ye Li

Higgs Couplings 2014

Torino, 10/01/14

Introduction

- ▶ Using interference effects in $gg \rightarrow \gamma\gamma$, LHC may bound Higgs width much better than in direct measurement [Dixon,Li] arXiv:1305.3854
- ▶ Maybe possible to get close to SM value of 4 MeV
- ▶ Similar idea can be used for $gg \rightarrow ZZ/WW$ in off-shell region [Caola,Melnikov] arXiv:1307.4935, [Campbell,Ellis,Williams] arXiv:1311.3589 (↗ talk by C. Williams)
- ▶ $gg \rightarrow \gamma\gamma$ more direct, as it operates in neighborhood of resonance
- ▶ $gg \rightarrow ZZ/WW$ method could be invalidated by e.g. form factors [Englert,Spannowsky] arXiv:1405.0285

Interference between Higgs production and continuum

[Dixon,Siu] hep-ph/0302233

- ▶ Full amplitude

$$\mathcal{A}_{gg \rightarrow \gamma\gamma} = \frac{-\mathcal{A}_{gg \rightarrow H} \mathcal{A}_{H \rightarrow \gamma\gamma}}{\hat{s} - m_H^2 + im_H \Gamma_H} + \mathcal{A}_{\text{cont}}$$

- ▶ Change in cross section from interference

$$\delta \hat{\sigma}_{gg \rightarrow H \rightarrow \gamma\gamma} = -2(\hat{s} - m_H^2) \frac{\text{Re}(\mathcal{A}_{gg \rightarrow H} \mathcal{A}_{H \rightarrow \gamma\gamma} \mathcal{A}_{\text{cont}}^*)}{(\hat{s} - m_H^2)^2 + m_H^2 \Gamma_H^2} - 2m_H \Gamma_H \frac{\text{Im}(\mathcal{A}_{gg \rightarrow H} \mathcal{A}_{H \rightarrow \gamma\gamma} \mathcal{A}_{\text{cont}}^*)}{(\hat{s} - m_H^2)^2 + m_H^2 \Gamma_H^2}$$

$$= \left[\begin{array}{c} g \\ t, b \\ g \end{array} \right] \begin{array}{c} \text{---} H \text{---} \\ \bullet \\ \text{---} W, t \\ \text{---} b, c, \tau \\ \text{---} \gamma \end{array} + \begin{array}{c} \text{---} \gamma \text{---} \\ \bullet \\ \text{---} \text{---} \end{array} + \dots \Big]$$

$$\times \left[\begin{array}{c} \text{---} \text{---} \\ \text{---} b, c, \dots \\ \text{---} \text{---} \end{array} + \begin{array}{c} \text{---} \text{---} \\ \text{---} u, c, d, s, b \\ \text{---} \text{---} \end{array} + \dots \Big]^*$$

- ▶ Real part of interference asymmetric around peak
- ▶ Imaginary part symmetric

Parametrizing new physics effects

- ▶ Effective coupling of Higgs to gluons & photons

$$\mathcal{L} = - \left[\frac{\alpha_s}{8\pi} c_g b_g G_{\mu\nu}^a G_a^{\mu\nu} + \frac{\alpha}{8\pi} c_\gamma b_\gamma F_{\mu\nu} F^{\mu\nu} \right] \frac{h}{v} \quad b_g = \frac{2}{3}, \quad b_\gamma = \frac{47}{9} \quad \text{at LO}$$

$c_{g/\gamma}$ – new physics correction factors

- ▶ In Narrow width approximation

$$d\sigma_{gg \rightarrow H \rightarrow \gamma\gamma} = \frac{d\hat{s} |\mathcal{A}_{gg \rightarrow H} \mathcal{A}_{H \rightarrow \gamma\gamma}|^2}{(\hat{s} - m_H^2)^2 + m_H^2 \Gamma_H^2} \propto \frac{c_g^2 c_\gamma^2}{\Gamma_H}$$


- ▶ Non-interference measurements invariant under scaling
 $c_{g/\gamma} \rightarrow \xi c_{g/\gamma}$ as $\Gamma_H \rightarrow \xi^4 \Gamma_H$
- ▶ Interference breaks degeneracy
- ▶ Allows to bound or even measure Higgs width

Mass shift from real part


[Martin] arXiv:1208.1533, arXiv:1303.3342

[deFlorian et al.] arXiv:1303.1397

- ▶ Smear lineshape with Gaussian of width 1.7 GeV (\sim detector resolution)






→




- ▶ Re-fitting to Gaussian of mass $M + \delta M$ gives $\delta M \sim 100$ MeV

Contributions to mass shift at NLO


[Dixon, Li] arXiv:1305.3854



+





+



Mass shift at NLO

[Dixon,Li] arXiv:1305.3854


- ▶ Large K-factor of Higgs production, smaller K-factor in background
→ relative size of interference reduced compared to LO
- ▶ Additional contribution from interference with tree-level diagrams
further reduces mass shift [deFlorian et al.] arXiv:1303.1397



Mass shift at NLO

[Dixon,Li] arXiv:1305.3854

- Mass shift vs jet veto p_T - mostly insensitive



Control masses

Possible control masses

1. $h \rightarrow ZZ \rightarrow 4l$
2. $h \rightarrow \gamma\gamma$ itself [Dixon,Li] arXiv:1305.3854
3. VBF-enriched $h \rightarrow \gamma\gamma$ [Dixon,Li,SH] in progress


- ▶ Theoretically ideal reference mass for measuring mass shift
→ ZZ channel, as $\delta m_{ZZ} \ll \delta m_{\gamma\gamma}$ [Kauer,Passarino] arXiv:1206.4803
- ▶ But experiments differ significantly

$$m_{\gamma\gamma} - m_{ZZ} = \begin{cases} +1.5 \pm 0.7 \text{ GeV} & \text{ATLAS} \\ -0.9 \pm 0.6 \text{ GeV} & \text{CMS} \end{cases}$$

Control mass from $h \rightarrow \gamma\gamma$

[Dixon,Li] arXiv:1305.3854, [Martin] arXiv:1303.3342



- ▶ Cancellation between qg and gg channels leaves strong dependence on $p_{T,h}$
- ▶ Points towards a possible measurement of shift in $\gamma\gamma$ channel alone by using sample with $p_{T,h} \gtrsim 40$ GeV as “control” region
- ▶ Experimental uncertainties (γ energy scale) would largely cancel



Control mass from VBF

[Dixon,Fidanza,deFlorian,Ita,Li,Mazzitelli,SH] in progress

- ▶ VBF more robust theoretically than high- $p_{T,h}$ region in $pp \rightarrow \gamma\gamma$
→ good possible control sample for mass measurement
- ▶ Mass shift from interference with $pp \rightarrow \gamma\gamma + 2$ jets
- ▶ About 1/3 the effect of $pp \rightarrow \gamma\gamma$, so 2/3 of effect remains



$$p_{T,\gamma} > 20 \text{ GeV}, |\eta_{\gamma}| < 2.5$$

$$p_{T,j} > 20 \text{ GeV}$$
$$m_{jj} > 800 \text{ GeV}, |\Delta\eta_{jj}| > 4$$


Mass shift versus width

[Dixon,Li] arXiv:1305.3854

- Assuming constant event yield, $c_{g\gamma} = c_g c_\gamma$ determined by

$$\frac{c_{g\gamma}^2 S}{m_H \Gamma_H} + c_{g\gamma} I \stackrel{!}{=} \left(\frac{S}{m_H \Gamma_H^{SM}} + I \right) \mu_{\gamma\gamma}$$



- Ignoring I leaves $c_{g\gamma} = \sqrt{\mu_{\gamma\gamma} \Gamma_H / \Gamma_H^{SM}}$



What if “the boson” was spin 2?

- ▶ Rejection of spin 2 hypothesis relies on $\cos \theta^*$ distribution
 [Maltoni et al.] arXiv:1306.6464, [Boer et al.] arXiv:1304.2654

- ▶ Without interference $\begin{cases} 1 & \text{for spin 0} \\ 1 + 6 \cos^2 \theta^* + \cos^4 \theta^* & \text{for } 2^+ \end{cases}$



- ▶ Interference from different helicity amplitudes:
 $\mathcal{A}(+, +, \pm, \pm)$ (spin 0) vs $\mathcal{A}(+, -, \pm, \mp)$ (spin 2_m^+)
 Assuming graviton-like couplings in spin-2 case

Signal vs interference in spin 2 case

[Dixon,Li,SH] in progress

$$\begin{aligned} |\overline{\mathcal{A}^{gg}}|^2 = & \left[\frac{\hat{s}^4}{M_G^4} \frac{\kappa_g^2}{256} f_0^{gg}(c) + \frac{\hat{s}^2}{M_G^2} \pi \xi M_G \Gamma_G f_i^{gg}(c) \right] \frac{1}{(\hat{s} - M_G^2)^2 + M_G^2 \Gamma_G^2} \\ & + \frac{\hat{s}^2}{M_G^2} \xi f_r^{gg}(c) \frac{\hat{s} - M_G^2}{(\hat{s} - M_G^2)^2 + M_G^2 \Gamma_G^2} \end{aligned}$$

where $c = \cos \theta^*$ and $\xi = \frac{11}{72} \kappa_g \alpha \alpha_s$



$$f_0^{gg}(c) = 1 + 6c^2 + c^4$$

$$f_i^{gg}(c) = 2 \left[\left(1 + \frac{(1-c)^2}{4} \right) \ln \left(\frac{2}{1-c} \right) + \left(1 + \frac{(1+c)^2}{4} \right) \ln \left(\frac{2}{1+c} \right) \right] - 3 + c^2$$

$$\begin{aligned} f_r^{gg}(c) = & \left(1 + \frac{(1-c)^2}{4} \right) \ln^2 \left(\frac{2}{1-c} \right) - \frac{(1+c)(3-c)}{2} \ln \left(\frac{2}{1-c} \right) \\ & + \left(1 + \frac{(1+c)^2}{4} \right) \ln^2 \left(\frac{2}{1+c} \right) - \frac{(1-c)(3+c)}{2} \ln \left(\frac{2}{1+c} \right) + 1 + c^2 \end{aligned}$$


Line shapes and signal yields

- ▶ Large constructive/destructive interference at large width
- ▶ Affects coupling measurement in spin 2 case




Angular dependence of imaginary part

- ▶ Standard cut $p_{T\min} = 40 \text{ GeV} \leftrightarrow \cos\theta_{\max}^* \approx 0.77$
- ▶ Imaginary part nearly flat in observed region \rightarrow difference in spin 0 and spin 2 yields can be accommodated by κ_g & Γ_G



Angular dependence of imaginary part

- Include radiation and coupling to quark part of energy-momentum tensor
→ Less flat distribution, depending on size of κ_g vs κ_q



Summary

- ▶ Interference effects allow to bound Higgs width well below experimental resolution in a fairly model independent way
- ▶ $\gamma\gamma$ channel shows large effect while working close to resonance mass
- ▶ Several possible control masses, including ZZ, $\gamma\gamma$ at high- $p_{T,h}$, and VBF
- ▶ Interferences also important for testing hypothesis involving non-SM quantum numbers