

FINITE SIZE OF HADRONS AND BOSE-EINSTEIN CORRELATIONS

A.Bialas and K.Zalewski
(Cracow)

- 1. WARNING**
- 2. TWO-PARTICLE CORRELATIONS**
- 3. INDEPENDENT PRODUCTION IN SPACE:
POSITIVE CORRELATION FUNCTION**
- 4. CORRELATED EMISSION: "EXCLUDED
VOLUME" EFFECT**
- 5. DATA**
- 6. COMMENTS**

ArXiv 1309.6169 [PLB 727 (2013) 182]

WARNING

BOSE-EINSTEIN CORRELATION BETWEEN MOMENTA OF TWO IDENTICAL HADRONS

$$C(p_1, p_2) \equiv \frac{N(p_1, p_2)}{N(p_1)N(p_2)} - 1 \quad (1)$$

IS USUALLY ANALYZED USING THE FORMULA

$$C(p_1, p_2) = \frac{\tilde{w}(P_{12}; Q)\tilde{w}(P_{12}; -Q)}{w(p_1)w(p_2)} = \frac{|\tilde{w}(P_{12}, Q)|^2}{w(p_1)w(p_2)} \quad (2)$$

WHERE $w(p.x)$ IS THE SINGLE-PARTICLE DISTRIBUTION (WIGNER FUNCTION) AND

$$\tilde{w}(P_{12}; Q) = \int dx e^{iQx} w(P_{12}; x); \quad w(p) = \int dx w(p; x) \\ P_{12} = (p_1 + p_2)/2; \quad Q = p_1 - p_2,$$

THIS PROCEDURE ASSUMES THAT HADRONS ARE UNCORRELATED.

DATA L3




Figure: L3 data for two-jet and three-jet events.

DATA CMS 1




Figure: Two-pion correlation function from CMS (pp at 7 TeV)

GENERAL TWO PARTICLE CORRELATIONS

LET $W(p_1, p_2; x_1, x_2)$ BE THE MOMENTUM AND SPACE "DISTRIBUTION" OF TWO PARTICLES ("SOURCE FUNCTION"). IF PARTICLES ARE IDENTICAL, THE OBSERVED MOMENTUM DISTRIBUTION IS

$$\begin{aligned}\Omega(p_1, p_2) &= \int dx_1 dx_2 W(p_1, p_2; x_1, x_2) + \\ &+ \int dx_1 dx_2 e^{i(x_1 - x_2)Q} W(P_{12}, P_{12}; x_1, x_2) \equiv \\ &\equiv \Omega_0(p_1, p_2) [1 + C(p_1, p_2)]\end{aligned}\quad (3)$$

WHERE $P_{12} = (p_1 + p_2)/2$, $Q = p_1 - p_2$, AND

$$\Omega_0(p_1, p_2) = \int dx_1 dx_2 W(p_1, p_2; x_1, x_2) \quad (4)$$

ONE SEES THAT $C(p_1, p_2)$ CONTAINS INFORMATION ONLY ON THE DISTRIBUTION OF $x_1 - x_2$.

NO INTER-PARTICLE CORRELATIONS

IF THERE ARE NO CORRELATIONS BETWEEN PARTICLES,

$$W(p_1, p_2; x_1, x_2) = w(p_1, x_1)w(p_2, x_2)$$

THEN $\Omega(p_1, p_2) = w(p_1)w(p_2) + |\tilde{w}(P_{12}, Q)|^2,$

WHERE $\tilde{w}(P_{12}, Q) = \int dx w(P_{12}, x)e^{ixQ}.$

THUS THE CORRELATION FUNCTION IS

$$C_2(p_1, p_2) = \frac{|\tilde{w}(P_{12}, Q)|^2}{w(p_1)w(p_2)} \geq 0!!!! \quad (5)$$

THIS IS THE COMMONLY USED FORMULA.

FROM $\tilde{w}(P_{12}, Q)$ ONE CAN RECOVER $w(P_{12}, x)$.

BUT: *THIS IS VALID ONLY IF THERE ARE NO INTER-PARTICLE CORRELATIONS.*

CORRELATIONS IN SPACE




IDEA: WHEN PIONS ARE TOO CLOSE TO EACH OTHER THEY ARE *NOT* PIONS ANYMORE!!! (BECAUSE THEIR CONSTITUENTS ARE MIXING AND THEIR WAVE FUNCTIONS ARE NOT WELL-DETERMINED).

SINCE HBT EXPERIMENTS MEASURE QUANTUM INTERFERENCE BETWEEN THE WAVE FUNCTIONS OF PIONS, THEY CANNOT SEE PIONS WHICH ARE TOO CLOSE TO EACH OTHER.

THEREFORE $W(P_{12}, P_{12}; x_1, x_2)$ MUST VANISH AT SMALL $|x_1 - x_2|$, IMPLYING *CORRELATION* BETWEEN POSITIONS OF TWO PIONS.

PICTURE

MIXING OF QUARKS



CORRELATIONS IN SPACE (2)

Repeat: $W(P_{12}, P_{12}; x_1, x_2)$ **MUST VANISH AT** $|x_1 - x_2| \approx 0$,
MEANING CORRELATION BETWEEN POSITIONS OF
TWO PIONS. THIS IS THE NECESSARY CONSEQUENCE
OF THE FUNDAMENTAL PROPERTY OF HADRONS:
THEY ARE NOT POINT-LIKE.

THUS THE TWO-PION DISTRIBUTION IS OF THE
FORM

$$W(P_{12}, P_{12}; x_1, x_2) = w(P_{12}; x_1)w(P_{12}; x_2)[1 - \Delta(x_1 - x_2)]. \quad (6)$$

THE CORRELATION FUNCTION:

$$C(P_{12}, Q) = \frac{|\tilde{w}(P_{12}, Q)|^2}{w(p_1)w(p_2)} - C_{corr}(p_1, p_2);$$
$$C_{corr} = \frac{\int dx_1 dx_2 e^{i(x_1 - x_2)Q} w(P_{12}; x_1)w(P_{12}; x_2)\Delta(x_1 - x_2)}{w(p_1)w(p_2)} \quad (7)$$

EXAMPLE

FOR ILLUSTRATION, TAKE

$$\Delta(x_1 - x_2) = \Theta[r_{cut}^2 - |\vec{x}_1 - \vec{x}_2|^2 - (t_1 - t_2)^2];$$
$$w(P, x) = e^{-|\vec{x}|^2/R^2} e^{-t^2/\tau^2} f(P)$$




Figure: Oscillating two-pion correlation function. $R = r_{cut} = \tau = 1$ fm.

DATA CMS 2




Figure: Two-pion correlation function for various multiplicities from CMS (pp at 7 TeV)

COMMENTS

(i) The presented qualitative argument shows that the observed negative values of the HBT correlation function are not accidental but reflect the fundamental fact that hadrons are not point-like. Therefore this region of Q^2 deserves more attention in data analysis. It seems that the effect simply **MUST BE THERE and the real experimental challenge is to determine its position and its size.**

(ii) More serious calculations, as well as a detailed comparison with data are clearly needed and are in progress (together with W.Florkowski).

Derivation of the symmetrized formula:

Density matrix in momentum space: $\rho(p_1, p_2; p'_1, p'_2) = \int dx_1 dx_2 e^{i(p_1 x_1 + p_2 x_2)} \int dx'_1 dx'_2 e^{-i(p'_1 x'_1 + p'_2 x'_2)} \rho(x_1, x_2; x'_1, x'_2)$.

The particle distribution is $\Omega(p_1, p_2) = \rho(p_1, p_2; p_1, p_2)$.

The Wigner function:

$W(p_1, p_2; x_1^+, x_2^+) = \int dx_1^- dx_2^- e^{i(p_1 x_1^- + p_2 x_2^-)} \rho(x_1, x_2; x_1', x_2')$

$x^+ = (x + x')/2; x_- = x - x'$

Symmetrization:

$$\rho(p_1, p_2; p'_1, p'_2) \rightarrow \rho(p_1, p_2; p'_1, p'_2) + \rho(p_1 p_2; p'_2, p'_1)$$

$$p_1 x_1 + p_2 x_2 - p_1 x'_1 - p_2 x'_2 = p_1 x_1^- - p_2 x_2^-$$

$$p_1 x_1 + p_2 x_2 - p_2 x'_1 - p_1 x'_2 = P_{12} x_1^- + P_{12} x_2^- + Q(x_1^+ - x_2^+) \quad (8)$$

$$\begin{aligned} \Omega(p_1, p_2) &= \int dx_1^+ dx_2^+ W(p_1, p_2; x_1^+, x_2^+) + \\ &+ \int dx_1^+ dx_2^+ e^{iQ(x_1^+ - x_2^+)} W(P_{12}, P_{12}; x_1^+, x_2^+) \end{aligned} \quad (9)$$