

## Cold nuclear matter physics at low-x from d+Au collisions at PHENIX

*Thursday, May 26, 2011 3:40 PM (20 minutes)*

RHIC experiments have observed that inclusive hadron yields in the forward rapidity (deuteron) direction for  $\sqrt{s_{NN}} = 200$  GeV d+Au collisions are suppressed relative to p+p collisions. The mechanism for the suppression has not been firmly established; theoretical descriptions include nuclear shadowing, initial-state parton energy loss, and gluon saturation. We present measurements by the PHENIX experiment at RHIC of di-hadron pair production in d+Au collisions where the particles in the pair are varied across a wide range of rapidity out to  $|\eta| = 3.8$  using a new forward electromagnetic calorimeter, the MPC. These di-hadron measurements probe down to parton momentum fractions  $x \sim 10^{-3}$  in the gold nucleus, where the interesting possibility of observing gluon saturation effects at RHIC is the greatest. Our measurements show that the correlated yield of back-to-back pairs in d+Au collisions is suppressed by up to an order of magnitude relative to p+p collisions, and increases with greater nuclear path thickness and with decreasing parton x in the Au nucleus.

**Author:** Dr CHIU, Mickey (Brookhaven National Lab)

**Presenter:** Dr CHIU, Mickey (Brookhaven National Lab)

**Session Classification:** Pre-equilibrium and initial state

**Track Classification:** Pre-equilibrium and initial state physics