

## Measurement of inclusive neutral pion yields in in p+p and Pb+Pb collisions at $\sqrt{s_{NN}}=2.76$ TeV with the ALICE EMCal

We present the measurement of inclusive differential  $\pi^0$  yields in p+p and centrality-selected Pb+Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV, using the large-acceptance ALICE Electromagnetic Calorimeter (EMCal). The recently completed EMCal enhances ALICE capabilities for the study of jets, neutral mesons, electrons, and direct photons. The EMCal acceptance (110 degrees in azimuth,  $|\eta| < 0.7$ ) and high granularity enable the reconstruction of neutral mesons over a wide transverse momentum range, employing invariant mass, shower-shape, and conversion techniques. We present first measurement of  $\pi^0 R_{AA}$  for both peripheral and central Pb+Pb collisions, normalized by the  $\pi^0$  yield measured in p+p collisions at  $\sqrt{s_{NN}} = 2.76$  TeV.

**Primary author:** Dr BRAIDOT, Ermes (Lawrence Berkeley National Lab. (LBNL))

**Presenter:** Dr BRAIDOT, Ermes (Lawrence Berkeley National Lab. (LBNL))

**Track Classification:** Pre-equilibrium and initial state physics