

## Charged Particle Ratios for p+p Collisions in $\sqrt{s} = 62.4$ GeV at RHIC

The ratios of particle production in hadronic interactions are important indicators of the collision dynamics [1]. These can be used to probe the process of hadronization in high energy collisions. We present measurements of mid-rapidity anti-particle to particle ratios in p + p collisions at  $\sqrt{s} = 62.4$  GeV from the STAR experiment. The measurements of the anti-particle to particle ratios are studied as a function of transverse momentum ( $p_T$ ) and comparison is made with corresponding ISR results for the p + p collisions at  $\sqrt{s} = 63$  GeV [2]. Identification of charged hadrons ( $\pi^\pm$ ,  $k^\pm$ , p and pbar) was done primarily through time projection chamber measurements. Charged hadrons are identified by using specific ionization energy loss ( $dE/dx$ ) at the low momentum region [3].

### References

- [1] H. Satz, Rep. Prog. Phys. 63 (2000) 151.
- [2] B. Alper et al., Nuclear Physics B 100 (1975) 237-290.
- [3] B. I. Abelev et al., [STAR Collaboration], Phys. Rev. C 79 (2009) 34909.

**Primary author:** GUPTA, Shikshit (Department of Physics)

**Presenter:** GUPTA, Shikshit (Department of Physics)

**Track Classification:** Hadron thermodynamics and chemistry