

Lattice QCD based equation of state at finite baryon density


Pasi Huovinen
J. W. Goethe Universität

Quark Matter 2011
May 24, 2011, Annecy, France

in collaboration with Peter Petreczky @ BNL

funded by BMBF (06FY9092) and DOE (DE-AC02-98CH1086)

Nuclear phase diagram



Taylor expansion for pressure

$$\frac{P}{T^4} = \sum_{i,j} c_{ij}(T) \left(\frac{\mu_B}{T}\right)^i \left(\frac{\mu_S}{T}\right)^j,$$

where

$$c_{ij}(T) = \frac{1}{i!j!} \frac{\partial^i}{\partial(\mu_B/T)^i} \frac{\partial^j}{\partial(\mu_S/T)^j} \frac{P}{T^4},$$

i.e. moments of baryon number and strangeness fluctuations and correlations

But: Most extensive study using p4 action with $N_\tau = 4$

⇒ large discretization effects?

Hadrons on lattice



- 16 pseudoscalar mesons on lattice
- Hadron masses depend on lattice cutoff
⇒ i.e. on temperature:
E.g. for pseudoscalar mesons on asqtad calculations

$$m_{\text{ps}_i}^2 = m_{\text{ps}_0}^2 + \frac{1}{r_1^2} \frac{a_{\text{ps}}^i x + b_{\text{ps}}^i x^2}{(1 + c_{\text{ps}}^i x)^{\beta_i}}$$

$$x = (a/r_1)^2$$

$$a = \frac{1}{N_\tau T}$$

30 MeV shift



Parametrization

$$c_{ij}(T) = \frac{a_{1ij}}{T^{n_{1ij}}} + \frac{a_{2ij}}{T^{n_{2ij}}} + \frac{a_{3ij}}{T^{n_{3ij}}} + \frac{a_{4ij}}{T^{n_{4ij}}} + \frac{a_{5ij}}{T^{n_{5ij}}} + SB_{ij},$$



where n_{kij} are integers with $1 < n_{kij} < 42$.

Require that


$$\begin{aligned} c_{ij}(T_{\text{sw}}) &= c_{ij}^{\text{HRG}}(T_{\text{sw}}) \\ \frac{d}{dT} c_{ij}(T_{\text{sw}}) &= \frac{d}{dT} c_{ij}^{\text{HRG}}(T_{\text{sw}}) \\ \frac{d^2}{dT^2} c_{ij}(T_{\text{sw}}) &= \frac{d^2}{dT^2} c_{ij}^{\text{HRG}}(T_{\text{sw}}) \end{aligned}$$

For second order terms also


$$c_{ij}(T = 800 \text{ MeV}) = 0.95 \cdot SB_{ij}$$




$$P/T^4$$



$$P/T^4$$




Softness of EoS




- s95p-v1 parametrization by P. Petreczky and P.H.


Softness of EoS




Softness of EoS




Softness of EoS




Softness of EoS



Pressure vs. energy density




p_T -spectra at SPS




- harder EoS, more transverse flow, flatter spectra

p_T -spectra at SPS



- $T_{\text{fo}} \approx 120$ MeV (bag) \Rightarrow 130 MeV (lattice)

v_2 at SPS ($b = 7$ fm)



- $T_{\text{fo}} \approx 120$ MeV (bag) $\Rightarrow 130$ MeV (lattice)

Conclusions

- lattice spacing dependence of hadron masses explains the difference between HRG and lattice QCD
 - 30 MeV shift in temperature
- EoS at finite baryon densities based on lattice QCD calculations of baryon number and strangeness fluctuations and correlations
 - ~10% uncertainty in speed of sound around the transition
- effect on flow when compared to bag model EoS tiny at SPS and (some?) RHIC low energy scan energies