

Charged particle production at large transverse momentum in PbPb collisions at $\sqrt{s}=2.76$ TeV measured with ALICE at the LHC.

Thursday, May 26, 2011 3:00 PM (20 minutes)

The observed suppression of high- p_T particle production in heavy-ion collisions is generally attributed to energy loss of partons as they propagate through the hot and dense QCD medium. Inclusive transverse momentum spectra of primary charged particles in Pb-Pb collisions at $\sqrt{s}=2.76$ TeV have been measured by the ALICE Collaboration at the LHC. The data are presented in intervals of collision centrality. The charged particle spectra are compared to those measured in pp collisions at the same collision energy, scaled by the number of underlying nucleon-nucleon collisions. This comparison is expressed in terms of the nuclear modification factor R_{AA} . The results indicate only weak medium effects in peripheral collisions, while there is clear evidence for strong medium effects in central collisions. In this talk we will present the results of an analysis of the full Pb-Pb statistics obtained in the Nov. 2010 run. With these data the transverse momentum range covered is increased to 50 GeV/c as compared to previous studies of R_{AA} by ALICE. The evolution of R_{AA} with collision centrality and transverse momentum will be discussed.

Primary author: OTWINOWSKI, Jacek (Research Division and ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung, Darmstadt, Germany)

Presenter: OTWINOWSKI, Jacek (Research Division and ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung, Darmstadt, Germany)

Session Classification: Jets

Track Classification: Jets