

Interference effects in medium-induced gluon radiation

Monday, May 23, 2011 3:40 PM (20 minutes)

The recent results at the LHC on jet quenching in heavy ion collisions call for a fundamental understanding of the evolution of a relatively hard jet propagating through a QCD medium like the quark-gluon plasma. As a first step in that sense, we consider the interference pattern for the medium-induced gluon radiation produced by a color singlet quark-antiquark antenna embedded in a QCD medium with size L . This setup is indeed well suited for studies of coherence phenomena like angular ordering. We focus on the most favorable kinematics for medium-induced gluon radiation in the BDMPS-Z regime, that is transverse momenta $k_{\perp} \sim \sqrt{\hat{q}L}$, short formation times $\tau_f \ll L$, and relatively large emission angles θ *gtrsim* $\theta_f \equiv (\hat{q}/\omega)^{1/3}$ (\hat{q} is the medium 'jet quenching' parameter and ω is the gluon energy). We demonstrate that, for a dipole opening angle $\theta_{q\bar{q}}$ larger than θ_f , there is no interference between medium-induced gluon emissions by the quark and the antiquark. That is, the in-medium contribution to the antenna pattern is the incoherent superposition of two individual BDMPS-Z spectra. Physically, this is so since, unlike the direct emissions which can be delocalized anywhere throughout the medium, the interference terms are non-zero only for the very early emissions, such that the size of the $q\bar{q}$ dipole at the time of emission be smaller than the transverse wavelength of the emitted gluon.

Primary authors: Dr IANCU, Edmon (CEA Saclay); Dr CASALDERREY SOLANA, Jorge (CERN)

Presenter: Dr CASALDERREY SOLANA, Jorge (CERN)

Session Classification: Theory developments

Track Classification: New theoretical developments