

## Higher harmonic jet tomography as a probe of fluctuating initial condition geometries in A+A

*Friday, May 27, 2011 3:40 PM (20 minutes)*

While 2nd Fourier harmonics of jet quenching have been thoroughly explored in the literature and shown to be sensitive to (1) the underlying jet path length dependence of energy loss and (2) the differences between the mean eccentricity predicted by Glauber and CGC/KLN models of initial conditions, the sensitivity of higher harmonics,  $v_n(p_T, b)$ , to differences between the fluctuation spectrum of geometries has remained relatively unexplored.

We demonstrate that higher azimuthal jet harmonics ( $n > 2$ ) of  $R_{AA}(p_T, \phi)$  and  $I_{AA}(p_T, \phi)$  are remarkably insensitive to the differences of geometrical density fluctuations comparing between Glauber and CGC/KLN models of the initial conditions. Therefore, the differential elliptic  $v_2(p_T)$  vs.  $v_2(I_{AA})(p_T)$  moment correlation between the 2nd moment of monojet  $R_{AA}$  and dijet  $I_{AA}$  nuclear modifications factors remains the most sensitive probe to differentiate between Glauber and CGC/KLN initial state sQGP geometries.

**Primary author:** BETZ, Barbara (Columbia University)

**Co-authors:** TORRIERI, Giorgio (Frankfurt Institute for Advanced Studies); GYULASSY, Miklos (Columbia University)

**Presenter:** BETZ, Barbara (Columbia University)

**Session Classification:** Jets

**Track Classification:** Jets