



$\Lambda_c^+(2940)$ in $p\bar{p} \rightarrow pD^0\bar{\Lambda}_c(2286)$
annihilation reaction with a
hadronic molecule scenario

Yubing Dong (董宇兵)

Institute of High Energy physics,
Chinese Academy of Sciences, Beijing, China

Collaborators: Amand Faessler, Thomas Gutsche,
V. E. Lyubovitskij (Tuebingen, Germany)


Content

- a Introduction: Hadronic molecules approach
- b Our approach, example
- c New resonance of $\Lambda_c(2940)$
- d Productions
- e Summary

a, Introduction

Charmonium

$c\bar{c}$



$\psi(4040)$

$Y(3940)$ $X(3940)$

$Z(3930)$

$X(3872)$

$\psi(3770)$

$D\bar{D}(3730)$

$\psi(2^3S_1)$

3690

$\eta'_c(2^1S_0)$

3640

$h_c(1^1P_1)$

3520

$\chi(1^3P_0)$

$\chi(1^3P_1)$

$\chi(1^3P_2)$

$\eta_c(1^1S_0)$

2980

$J/\psi(1^3S_0)$

3100

J^{PC}

0^{-+}

1^{+-}

1^{--}


0^{++}

1^{++}

2^{++}

Potential models worked well for charmonium spectroscopy

Charmonium '10



Charged charmonium spectrum
-- A completely new scenario of strong QCD!

States close to open thresholds
-- The role played by open D meson channels?

Close to DD^* threshold

2015/3/3

QNP2015 (March 2-6), Valparaiso, Chile

XYZ resonances [Hidden-charm(bottom)]

- 1, Conventional quark model ×
- 2, Narrow width
- 3, Near threshold of two mesons...

Interpretations:

- Molecule, baryonium
- tetraquark
- Hybrids
- Coupling channel...

Molecular scenario
Quark and hadron level

Frame Works:

QCD sum rule

Non relativistic QCD

Heavy quark effective theory

Heavy hadron chiral
perturbation theory

Potential models

Lattice calculations

Hadronic molecules

- Weekly bound state of two or three hadrons
- Obvious examples: Nuclei and hyper-nuclei
- Baryon-baryon bound state: $M_H < M_1 + M_2$
- Dynamical Generation of molecular bound states/ resonances

Long-range one-pion exchange (Tornqvist, 1991)

Meson-exchange models (Lohse, et al., 1990)

Unitarized coupled channel models with chiral Lagrangians (Olier, et al., 1997;

Jido et al., 2005,

Gammermann et al., 08)


b, Our approach(4-Dim.):

Bound state description of hadronic molecule on QFT based on the Compositeness condition: $Z_M=0$

(See Weinberg: PR130 (1966),776; Salam, Nuov. Cim. 25 (1962),224...)

$$L_{XDD} = X_\mu J^\mu$$

$$= \frac{g_x}{\sqrt{2}} X_\mu \int d^4 y \Phi(y^2) [D(x+y/2) \bar{D}^{*\mu}(x-y/2) + \bar{D}(x+y/2) D^{*\mu}(x-y/2)]$$



Mass operator

$$Z_x = 1 - g^2 \Pi'(M^2) = 0 \leftarrow$$

□

Vertex function/Correlation function

Vertex function $\Phi(y^2) \rightarrow$ finite size effects
/distributions of the constituents in bound state

*****Local limit:**

$$\Phi(y^2) \rightarrow \delta^4(y^2)$$

*****momentum space**

$$\tilde{\Phi}(k^2) \rightarrow \exp(-k_E^2 / \Lambda^2)$$

Λ is model parameter

k_E is relative Euclidean Jacobi momentum (4-D)

□

Example: New resonances: X(3872) (meson)

Basics about X(3872)

first seen in

$X(3872) \rightarrow J/\psi \pi^+ \pi^-$ by BELLE (2003),
also seen by CDF, D0 (2004) and BABAR (2005).

$\Gamma_X \approx 3 \text{ MeV}$

quantum numbers:

$C=+$ from $X(3872) \rightarrow \gamma J/\psi$, $I=0$ no signal in $X \rightarrow \pi \pi^0 J/\psi$

$J^{PC} = 1^{++}$ or $J^{PC} = 2^{-+}$ from $X(3872) \rightarrow J/\psi \pi^+ \pi^-$ helicity amplitude

$X(3872.2 \pm 0.8)$ close to $D^0 \bar{D}^{*0}$ threshold with $m_{thr} = 3871.81 \pm 0.36 \text{ Me}$

S-wave $D^0 \bar{D}^{*0}$ hadron molecule favors $J^{PC} = 1^{++}$



charmonium interpretation disfavored, $1^{++}(2^3P_1)$ too low in mass compare

$m(2^3P_2) \approx m(Z(3930))$

2015/3/3

$X(3872) \rightarrow \pi^+ \pi^- J/\psi$

arXiv:0809.1224 605 fb⁻¹



recent results



$M(X(3872)) = (3871.46 \pm 0.37 \pm 0.07) \text{ MeV}$
by combining two modes together

$$B(B^0 \rightarrow X(3872)(K^+ \pi^-)_{NR}) \times B(X(3872) \rightarrow J/\psi \pi^+ \pi^-) = (8.1 \pm 2.0^{+1.1}_{-1.4}) \times 10^{-6}$$

Calculation: Molecular scenario

Ansatz: $X(3872)$ is S-wave molecule with $J^{PC} = 1^{++}$

$$|X(3872)\rangle = \cos\theta \left[\frac{Z_{D^0\bar{D}^{*0}}^{1/2}}{\sqrt{2}} (|D^0\bar{D}^{*0}\rangle + |D^{*0}\bar{D}^0\rangle) + \frac{Z_{D^\pm D^{*\mp}}^{1/2}}{\sqrt{2}} (|D^+D^{*-}\rangle + |D^-D^{*+}\rangle) + Z_{J_\psi\omega}^{1/2} |J_\psi\omega\rangle + Z_{J_\psi\rho}^{1/2} |J_\psi\rho\rangle \right] + \sin\theta |c\bar{c}\rangle$$

$$(m_{D^0} = 1864.85 \text{ MeV}, m_{D^{*0}} = 2006.7 \text{ MeV}, m_X = m_{D^0} + m_{D^{*0}} - \epsilon)$$

- dominant $|D^0\bar{D}^{*0}\rangle + |D^{*0}\bar{D}^0\rangle$ component
- quantitatively see Swanson (2004): for $\epsilon = 0.3 \text{ MeV}$,
 $Z_{D^0\bar{D}^{*0}} = 0.92$, $Z_{D^\pm D^{*\mp}} = 0.033$, $Z_{J_\psi\omega} = 0.041$, $Z_{J_\psi\rho} = 0.006$
- small admixture of $1^{++} c\bar{c}$ component: $\propto \sin\theta$
- Compositeness condition: $Z_X = 1 - (\Sigma_X^M(m_X^2))' - (\Sigma_X^C(m_X^2))' = 0$ fixes coupling of X to its components

Effective Lagrangians

$$\begin{aligned}\mathcal{L}_X^L(x) &= g_{XD^0D^0} X_\mu(x) J_{D^0D^0}^\mu(x) \\ &+ g_{XD^zD^z} X_\mu(x) J_{D^zD^z}^\mu(x) \\ &+ \frac{g_{XJ_\psi^\omega}}{m_X} \epsilon_{\mu\nu\alpha\beta} \partial^\nu X^\alpha(x) J_{J_\psi^\omega}^{\mu\beta}(x) \\ &+ \frac{g_{XJ_\psi^\rho}}{m_X} \epsilon_{\mu\nu\alpha\beta} \partial^\nu X^\alpha(x) J_{J_\psi^\rho}^{\mu\beta}(x),\end{aligned}$$

$$J_{D\bar{D}^*}^\mu(x) = \frac{1}{\sqrt{2}}(D(x)\bar{D}^{*\mu}(x) + \bar{D}(x)D^{*\mu}(x)),$$

$$J_{J_\psi V}^{\mu\beta} = J_\psi^\mu V^\beta,$$

Non-local ones


$$\begin{aligned}J_{D\bar{D}^*}^\mu(x) \rightarrow \mathcal{J}_{D\bar{D}^*}^\mu(x) &= \frac{1}{\sqrt{2}} \int d^4y \Phi_{D\bar{D}^*}(y^2) (D(x+y/2) \\ &\times \bar{D}^{*\mu}(x-y/2) \\ &+ \bar{D}(x+y/2)D^{*\mu}(x-y/2)),\end{aligned}$$

$$J_{J_\psi V}^\mu(x) \rightarrow \mathcal{J}_{J_\psi V}^{\mu\beta}(x) = J_\psi^\beta(x) \int d^4y \Phi_V(y^2) V^\mu(x+y)$$

Radiative decays

$$X(3872) \rightarrow J/\psi, \psi(2S) + \gamma$$

Decay width (keV)



Approach	$\Gamma(X(3872) \rightarrow \gamma J/\psi)$
$[c\bar{c}]$, Ref. [9]	11
$[c\bar{c}]$, Ref. [33]	71
$[c\bar{c}]$, Ref. [33]	139
[molecule], Ref. [33]	8
Our results	124.8 - 231.3 ($\epsilon = 0.7$ MeV)
	129.8 - 239.1 ($\epsilon = 1$ MeV)
	138.0 - 251.4 ($\epsilon = 1.5$ MeV)

PRD77, 094013

PRD79, 094013

Strong decay(two-body, three-body)

New measurement

• $\Gamma(X \rightarrow \psi(2S)\gamma)/\Gamma(X \rightarrow J/\psi\gamma) = 3.5 \pm 1.4$

BABAR, PRL 102, (2009)

possible evidence for charmonium component ?

Exotic charmonium-like spectroscopy at LHCb:
a study of the $X(3872)$ and of the $Z(4430)^-$ states

1409.6472

Radiative Decay $X(3872) \rightarrow J/\psi \gamma, \psi' \gamma$

To study this further, LHCb has recently measured [7] the ratio of branching fractions

$$R_{\psi\gamma} = \frac{B(X(3872) \rightarrow \psi(2S)\gamma)}{B(X(3872) \rightarrow J/\psi\gamma)},$$

as a constraint on the charmonium content of the $X(3872)$. The branching fraction $B(X(3872) \rightarrow \psi(2S)\gamma)$ is in fact expected to be very small for a pure molecule ($O(10^{-3})$) [8-10], but it could be enhanced for an admixture of a $D^{*0}\bar{D}^0$ molecule and charmonium. The BaBar collaboration has measured a relative large branching fraction for the $X(3872)$ into $\psi(2S)\gamma$, with $R_{\psi\gamma} = 3.4 \pm 1.4$ [11], a result generally inconsistent with a pure molecular interpretation; in contrast, no significant signal was found by Belle [12].

$$\bar{R}_{\psi\gamma} = 2.46 \pm 0.64 \pm 0.29,$$

Results


KS

Results for $X(3872) \rightarrow \gamma J/\psi$ and $\psi(2s)$


JPG38, 015001

Configuration	$\Gamma(X(3872) \rightarrow \gamma J/\psi, \gamma \psi(2S))$ keV
molecular DD^* component	60 - 120(J/ψ) 0.3 ($\psi(2S)$)
pure $J/\psi V$ component	6(J/ψ) 0 ($\psi(2S)$)
interfering DD^* and $J/\psi V$ components	30 - 65 (J/ψ) 0.3 ($\psi(2S)$)

additional charmonium contribution with $Z_{c\bar{c}}^{1/2} = \sin\theta \approx -0.2$ required



dotted - J/ψ , solid - $\psi(2s)$ mode



$$R_{2s} = \frac{\Gamma(X \rightarrow \psi(2S) + \gamma)}{\Gamma(X \rightarrow J/\psi + \gamma)} = 3.5 \pm 1.4$$

(BABAR, 2009)

2015/3/3

C, New baryon resonance of $\Lambda_c(2940)$


PRL 98, 012001 (2007)

PHYSICAL REVIEW LETTERS

week ending
5 JANUARY 2007

Observation of a Charmed Baryon Decaying to $D^0 p$ at a Mass Near $2.94 \text{ GeV}/c^2$

(*BABAR* Collaboration)



The results for the $\Lambda_c(2940)^+$ baryon are

$$m = [2939.8 \pm 1.3(\text{stat}) \pm 1.0(\text{syst})] \text{ MeV}/c^2,$$

$$\Gamma = [17.5 \pm 5.2(\text{stat}) \pm 5.9(\text{syst})] \text{ MeV}.$$

For the $\Lambda_c(2880)^+$ baryon the results are

$$m = [2881.9 \pm 0.1(\text{stat}) \pm 0.5(\text{syst})] \text{ MeV}/c^2,$$

$$\Gamma = [5.8 \pm 1.5(\text{stat}) \pm 1.1(\text{syst})] \text{ MeV}.$$

Recently a new baryon resonance $\Lambda_c(2940)^+$ has been discovered in the decay channel $D^0 p$ by the *BABAR* Collaboration [1] and confirmed as a resonant structure in the final state $\Sigma_c(2455)^{0,++} \pi^\pm \rightarrow \Lambda_c^+ \pi^+ \pi^-$ by Belle

Experimental Constraints on the Spin and Parity of the $\Lambda_c(2880)^+$

(Belle Collaboration)

We report the results of several studies of the $\Lambda_c^+ \pi^+ \pi^- X$ final state in continuum e^+e^- annihilation data collected by the Belle detector. An analysis of angular distributions in $\Lambda_c(2880)^+ \rightarrow \Sigma_c(2455)^{0,++} \pi^{+,-}$ decays strongly favors a $\Lambda_c(2880)^+$ spin assignment of $\frac{5}{2}$ over $\frac{3}{2}$ or $\frac{1}{2}$. We find evidence for $\Lambda_c(2880)^+ \rightarrow \Sigma_c(2520)^{0,++} \pi^{+,-}$ decay and measure the ratio of $\Lambda_c(2880)^+$ partial widths $\Gamma(\Sigma_c(2520)\pi)/\Gamma(\Sigma_c(2455)\pi) = 0.225 \pm 0.062 \pm 0.025$. This value favors the $\Lambda_c(2880)^+$ spin-parity assignment of $\frac{5}{2}^+$ over $\frac{5}{2}^-$. We also report the first observation of $\Lambda_c(2940)^+ \rightarrow \Sigma_c(2455)^{0,++} \pi^{+,-}$ decay and measure $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ mass and width parameters. These studies are based on a 553 fb^{-1} data sample collected at or near the $Y(4S)$ resonance at the KEKB collider.

TABLE I. Signal yield, mass, and width for the $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$. The first uncertainty is statistical, the second one systematic.

State	Yield	M (MeV/ c^2)	Γ (MeV)
$\Lambda_c(2880)^+$	690 ± 50	$2881.2 \pm 0.2 \pm 0.4$	$5.8 \pm 0.7 \pm 1.1$
$\Lambda_c(2940)^+$	220^{+80}_{-60}	$2938.0 \pm 1.3^{+20}_{-40}$	13^{+8+27}_{-5-7}

CHARMED BARYONS ($C = +1$)

$$\Lambda_c^+ = udc, \quad \Sigma_c^{++} = uuc, \quad \Sigma_c^+ = udc, \quad \Sigma_c^0 = ddc, \\ \Xi_c^+ = usc, \quad \Xi_c^0 = dsc, \quad \Omega_c^0 = ssc$$

Λ_c^+

$$I(J^P) = 0(\frac{1}{2}^+)$$

J is not well measured; $\frac{1}{2}$ is the quark-model prediction.

$$\text{Mass } m = 2286.46 \pm 0.14 \text{ MeV}$$

$\Lambda_c(2940)^+$

$$I(J^P) = 0(?^?)$$

$$\text{Mass } m = 2939.3^{+1.4}_{-1.5} \text{ MeV}$$

$$\text{Full width } \Gamma = 17^{+8}_{-6} \text{ MeV}$$

$\Lambda_c(2940)^+$ DECAY MODES

Decay Mode	Fraction (Γ_i/Γ)	p (MeV/ c)
pD^0	seen	420
$\Sigma_c(2455)^{0,++} \pi^\pm$	seen	—

Different interpretations

1), quark model:

Isgur-Karl ($3/2^+$, $5/2^-$)

Heavy-light diquark model
(radial excitation of Λ_c)

2), Chiral quark model:

D-Wave

3), Molecular Model

near threshold of pD^*

the assignment of the resonance

$$\mathcal{L}_{\Lambda_c}(x) = g_{\Lambda_c} \bar{\Lambda}_c^+(x) \Gamma^\mu \int d^4y \Phi(y^2) (\cos\theta D_\mu^{*0}(x) p(x+y) + \sin\theta D_\mu^{*+}(x) n(x+y)) + \text{H.c.},$$

$$Z_{\Lambda_c} = 1 - \Sigma'_{\Lambda_c}(m_{\Lambda_c}) = 0.$$



$$\Gamma^\mu = \gamma^\mu \text{ for } J^P = \frac{1}{2}^+ \text{ and } \Gamma^\mu = \gamma_5 \gamma^\mu \text{ for } J^P = \frac{1}{2}^-$$

$$|\Lambda_c(2940)^+\rangle = \cos\theta |pD^{*0}\rangle + \sin\theta |nD^{*+}\rangle.$$





FIG. 1 (color online). Diagram describing the $\Lambda_c(2940)^+$ mass operator.




FIG. 2 (color online). Diagrams contributing to the decays $\Lambda_c(2940)^+ \rightarrow pD^0, \Sigma_c^{++}\pi^-, \Sigma_c^0\pi^+$.

Effective Lagrangians

$$\begin{aligned}
 \mathcal{L}_{VPBB} = & ig_1 \bar{B}^{kmn} \gamma^\mu \gamma^5 [V_\mu, P]_k^l B_{lmn} \\
 & + ig_2 \bar{B}^{kmn} \gamma^\mu \gamma^5 [V_\mu, P]_k^l B_{lnm} \\
 & + ig_3 \bar{B}^{kmn} \gamma^\mu \gamma^5 ((V_\mu)_k^l P_m^s - P_k^l (V_\mu)_m^s) B_{lns} \\
 & - ig_3 \bar{B}^{knm} \gamma^\mu \gamma^5 ((V_\mu)_k^l P_m^s - P_k^l (V_\mu)_m^s) B_{lsm}
 \end{aligned}$$

PRD81, 014006

The strong two-body decay widths of the $\Lambda_c(2940)^+$ baryon are calculated according to the expressions

$$\begin{aligned}
 \Gamma(\Lambda_c[1/2^+] \rightarrow B + M) = & \frac{g_{\Lambda_c BM}^2}{16\pi m_{\Lambda_c}^3} \lambda^{1/2}(m_{\Lambda_c}^2, m_B^2, m_M^2) \\
 & \times ((m_{\Lambda_c} - m_B)^2 - m_M^2) \quad (9)
 \end{aligned}$$

for the positive parity $\Lambda_c(2940)^+$ state and accordingly

$$\begin{aligned}
 \Gamma(\Lambda_c[1/2^-] \rightarrow B + M) = & \frac{f_{\Lambda_c BM}^2}{16\pi m_{\Lambda_c}^3} \lambda^{1/2}(m_{\Lambda_c}^2, m_B^2, m_M^2) \\
 & \times ((m_{\Lambda_c} + m_B)^2 - m_M^2) \quad (10)
 \end{aligned}$$

for the negative parity choice for $\Lambda_c(2940)^+$.

Calculated results

TABLE I. Partial decay widths of $\Lambda_c(2940)^+$ in MeV.

$\cos\theta$	$\frac{1}{2}^+$ modes			$\frac{1}{2}^-$ modes		
	$\Lambda_c^+ \rightarrow pD^0$	$\Lambda_c^+ \rightarrow \Sigma_c^{++}\pi^-$	$\Lambda_c^+ \rightarrow \Sigma_c^0\pi^+$	$\Lambda_c^+ \rightarrow pD^0$	$\Lambda_c^+ \rightarrow \Sigma_c^{++}\pi^-$	$\Lambda_c^+ \rightarrow \Sigma_c^0\pi^+$
1	0.11	0.58	0.72	19.15	612.68	756.72
0.95	0.17	0.85	0.98	29.75	907.64	1040.36
0.9	0.20	0.96	1.08	34.40	1033.00	1153.95
0.8	0.23	1.11	1.20	41.09	1208.89	1305.10
0.7	0.25	1.20	1.27	46.17	1338.06	1407.80
0.6	0.27	1.27	1.30	50.24	1437.58	1478.96
0.5	0.28	1.31	1.32	53.47	1511.85	1522.78
0.4	0.29	1.32	1.30	55.83	1560.10	1538.24
0.3	0.29	1.32	1.30	55.83	1560.10	1538.24
0.2	0.29	1.30	1.26	57.15	1577.04	1519.78
0.1	0.26	1.14	1.03	54.20	1447.05	1309.75
0.05	0.24	1.04	0.91	50.68	1334.05	1174.51
0	0.18	0.74	0.60	38.15	964.41	781.52

Radiative decay(1 + /2)

PRD81, 034035

Gauge Invariance

$$\mathcal{M}^\mu(p, p') = \bar{u}_{\Lambda'_c}(p') \Gamma^\mu(p, p') u_{\Lambda_c}(p),$$

$$\Gamma^\mu(p, p') = F_1(q^2) \gamma^\mu + F_2(q^2) i \sigma^{\mu\nu} q_\nu + F_3(q^2) q^\mu.$$


$$F_1(q^2) = F_3(q^2) \frac{q^2}{m_{\Lambda_c} - m_{\Lambda'_c}}.$$

$$\mathcal{M}^\mu(p, p') = \frac{F_{\Lambda_c \Lambda'_c \gamma}}{2m_{\Lambda_c}} \bar{u}_{\Lambda'_c}(p') i \sigma^{\mu\nu} q_\nu u_{\Lambda_c}(p).$$

$$\Gamma(\Lambda_c(2940)^+ \rightarrow \Lambda_c(2286)^+ + \gamma) = \frac{\alpha P^{*3}}{m_{\Lambda_c}^2} F_{\Lambda_c \Lambda'_c \gamma}^2$$

TABLE III. Radiative decay width of $\Lambda_c(2940)^+$ in keV.


θ (in grad)	Λ (GeV)					
	0.25	0.4	0.5	0.75	1	1.25
0	11.1	35.4	61.7	113.1	142.7	156.8
5	9.2	29.2	51.0	91.5	112.2	119.4
10	7.4	23.2	40.6	71.0	83.9	85.5
15	5.7	17.6	30.8	52.1	58.6	56.2
20	4.1	12.5	22.0	35.5	37.1	32.4
25	2.7	8.1	14.4	21.7	20.1	14.7



Diagrams contributing to the radiative decay process
 $\Lambda_c(2940)^+ \rightarrow \Lambda_c(2286)^+ + \gamma$.


$$q_\mu \mathcal{M}^\mu(p, p') = 0$$

Three-body decay



$$\begin{aligned} \mathcal{L}_{\pi^- D^*0 p \Sigma_c^{++}} &= \left[\frac{1}{4}(g_1 + g_2) - \frac{3}{2}g_3 \right] \bar{\Sigma}_c^{++} \pi^- i\gamma^\mu \gamma_5 p D_\mu^{*0} + \text{H.c.}, & \mathcal{L}_{\pi^- D^{*+} n \Sigma_c^{++}} &= -\frac{3}{2}g_3 \bar{\Sigma}_c^{++} \pi^- i\gamma^\mu \gamma_5 n D_\mu^{*+} + \text{H.c.}, \\ \mathcal{L}_{\pi^0 D^*0 p \Sigma_c^+} &= \frac{1}{2} \left[\frac{1}{4}(g_1 + g_2) - 3g_3 \right] \bar{\Sigma}_c^+ \pi^0 i\gamma^\mu \gamma_5 p D_\mu^{*0} + \text{H.c.}, & \mathcal{L}_{\pi^0 D^{*+} n \Sigma_c^+} &= \frac{1}{2} \left[\frac{1}{4}(g_1 + g_2) - 3g_3 \right] \bar{\Sigma}_c^+ \pi^0 i\gamma^\mu \gamma_5 n D_\mu^{*+} + \text{H.c.}, \\ \mathcal{L}_{\pi^+ D^*0 p \Sigma_c^0} &= -\frac{3}{2}g_3 \bar{\Sigma}_c^0 \pi^+ i\gamma^\mu \gamma_5 p D_\mu^{*0} + \text{H.c.}, & \mathcal{L}_{\pi^+ D^{*+} n \Sigma_c^0} &= \left[\frac{1}{4}(g_1 + g_2) - \frac{3}{2}g_3 \right] \bar{\Sigma}_c^0 \pi^+ i\gamma^\mu \gamma_5 n D_\mu^{*+} + \text{H.c.}, \\ \mathcal{L}_{\pi \Sigma_c \Lambda_c'} &= -\frac{1}{2} \sqrt{\frac{3}{2}} \left(g_2' - \frac{1}{2}g_1' \right) \bar{\Lambda}_c' i\gamma^5 \pi \Sigma_c + \text{H.c.}, & \mathcal{L}_{D^* N \Lambda_c'} &= -g_{D^* N \Lambda_c'} \kappa_{D^* N \Lambda_c'} \bar{N} \sigma^{\mu\nu} \partial_\nu D_\mu^* \Lambda_c' + \text{H.c.}, \\ \mathcal{L}_{\rho \pi \pi} &= g_{\rho \pi \pi} \rho_k^\mu \pi_i \partial_\mu \pi_j \epsilon_{ijk}, & \Gamma &= \frac{\beta}{512\pi^3 M_{\Lambda_c}^3} \int_{4M_\pi^2}^{(M_{\Lambda_c} - M_{\Lambda_c'})^2} ds_2 \int_{s_1^-}^{s_1^+} ds_1 \sum_{\text{pol}} |M_{\text{inv}}|^2, \\ \mathcal{L}_{\rho D^* N \Lambda_c'} &= \frac{g_{\rho D^* N \Lambda_c'}}{2M_N} \bar{N} D_\mu^{*+} i\sigma^{\mu\nu} \rho_\nu \Lambda_c' + \text{H.c.}, \end{aligned}$$

d, Production @ PANDA



- Home
- News
- Meetings
- Collaboration
- Computing
- Detector
- Physics
- Documents
- Communications

Welcome to the PANDA Experiment Website

The PANDA Experiment will be one of the key experiments at the Facility for Antiproton and Ion Research (FAIR) which is under construction and currently being built on the area of the GSI Helmholtzzentrum für Schwerionenforschung in Darmstadt, Germany.

The central part of FAIR is a synchrotron complex providing intense pulsed ion beams (from p to U). Antiprotons produced by a primary proton beam will then be filled into the High Energy Storage Ring (HESR) which collide with the fixed target inside the PANDA Detector.


The PANDA Collaboration with more than 450 scientist from 17 countries intends to do basic research on various topics around the weak and strong forces, exotic states of matter and the structure of hadrons.

In order to gather all the necessary information from the antiproton-proton collisions being able to provide precise trajectory reconstruction, energy and momentum identification of charged particles.

What do you want to know more about?



The Physics Program
Information about the various physics topics going to be investigated by PANDA.



The Detector
Detailed description and technical information about the different detection systems.



The Accelerator Facility
Information about host laboratory GSI, the Facility for Antiproton and Ion Research and the accelerator.



The PANDA Collaboration
Contact information, structure and working groups within the Collaboration.

Forthcoming experiments at PANDA, with the \bar{p} momentum in the range from 1 to 15 GeV/c, which corresponds to total center-of-mass energies in the antiproton-proton system between 2.25 and 5.5 GeV, can give rich contributions to these investigations [1]. For example, $p\bar{p}$ annihilation reactions are expected to provide substantial information on the charm baryon $\Lambda_c(2286)$ as well as the baryon resonance $\Lambda_c(2940)$ recently observed by the BABAR Collaboration [2] and confirmed by the Belle Collaboration [3].

BEPC, BABAR, BELLE, JLab. PANDA



2015/3/3

QNP2015 (March 2-6), Valparaiso, Chile

Production

$$|\Lambda'_c(2940)\rangle = |pD^{*0}\rangle$$

PRD90, 094001

$$\mathcal{L}_{\Lambda'_c p D}^{2^{1+}} = g_{\Lambda'_c p D} \bar{\Lambda}'_c i \gamma_5 p D^0 + \text{H.c.},$$


$$\mathcal{L}_{\Lambda'_c p D^*}^{2^{1+}} = g_{\Lambda'_c p D^*} \bar{\Lambda}'_c \gamma^\mu p D_\mu^{*0} + \text{H.c.} \quad 1+ / 2$$

$$\frac{d\sigma}{dM_{pD}} = \frac{1}{1024\pi^4 s} \frac{1}{\sqrt{s - 4M_N^2}} \times \int d\cos\theta_3 d\Omega_1^* |\vec{q}_1^*| |\vec{q}_2| |\mathcal{M}_{\text{inv}}|^2$$



$$\mathcal{L}_{\Lambda'_c p D}^{1^-} = f_{\Lambda'_c p D} \bar{\Lambda}'_c p D^0 + \text{H.c.},$$

1 - / 2

$$\mathcal{L}_{\Lambda'_c p D^*}^{1^-} = f_{\Lambda'_c p D^*} \bar{\Lambda}'_c \gamma^\mu \gamma^5 p D_\mu^{*0} + \text{H.c.}$$



Initial state interaction



$$T(\vec{q}', \vec{q}; E) = V(\vec{q}', \vec{q}; E) + \int \frac{d^3 p V(\vec{q}', \vec{p}) T(\vec{p}, \vec{q}; E)}{E(q) - E(p) + i\epsilon}$$

$$V_{N\bar{N}}(\vec{q}', \vec{q}) = V_{N\bar{N}}^\pi(\vec{q}', \vec{q}) + V_{N\bar{N}}^{\text{opt}}(\vec{q}', \vec{q}).$$

The π -exchange potential is given by [23,24]

$$V_{N\bar{N}}^\pi(\vec{q}', \vec{q}) = \frac{g_{\pi NN}^2}{12M_N^2 M_\pi^2 + k_\pi^2} \times (\vec{\sigma}_1 \cdot \vec{\sigma}_2 + \hat{S}_{12}(\vec{k}_\pi)) (\vec{\tau}_1 \cdot \vec{\tau}_2) F_\pi^2(k_\pi^2),$$

The optical potential for the $N\bar{N}$ scattering state is :

$$V_{N\bar{N}}^{\text{opt}}(r) = (u_0 + iw_0)e^{-r^2/2r_0^2}$$







FIG. 3. Initial state interaction factor $|J_0|^2$ in dependence on $s^{1/2}$.




(a)



Differential cross section $d\sigma/dM_{pD}$ for $J^P = \frac{1}{2}^+$ of the $\Lambda_c(2940)$.



(b)





Differential cross section $d\sigma/dM_{pD}$ for $J^P = \frac{1}{2}^-$ of the $\Lambda_c(2940)$.



(a)

Differential cross section $d\sigma/dM_{pD}$ for $J^P = \frac{1}{2}^+$ of the $\Lambda_c(2940)$.



(b)

Differential cross section $d\sigma/dM_{pD}$ for 5.5 GeV for $J^P = \frac{1}{2}^-$ of the $\Lambda_c(2940)$.

e, Summary

- 1), Hadronic molecules: old expectations-
renewed interest in heavy mesons
- 2), Effective approach is applied
to the states (Compositeness, 4-dim.)
- 3), Hadronic loop is considered
- 4), Production

5), Some resonances

X(3872), $\Lambda_c(2940)$, $Z_c(3900)$, deuteron, Y(4260)
 $Z_b(10610)$, $Z'_b(10650)$

Thanks! Organizers for QNP2015