

Tetraquarks

AD Polosa

Sapienza Università di Roma

anti-triplet as anti-quarks \bar{q} . Baryons can now be constructed from quarks by using the combinations (qqq) , $(qqqq\bar{q})$, etc., while mesons are made out of $(q\bar{q})$, $(qq\bar{q}\bar{q})$, etc. It is assuming that the lowest



of $(\bar{d}\bar{d})^2$, $(\bar{d}\bar{d}\bar{d}\bar{d})^2$ etc. It is assuming that the lowest

Taken from Gell-Mann paper
on quark model


Large N_c and Tetraquarks

Following E. Witten's paper on $1/N$ and Baryons (see also S. Coleman's lectures), tetraquarks should instantly fall apart into mesons.

However, as pointed by S. Weinberg in a recent paper (PRL 110, 2013), this applies only at the leading order N^2 *disconnected* diagram. The leading order *connected* diagram has only one color loop and excludes the leading order of free two meson propagation.



Large N_c and Tetraquarks





which implies that the $4q$ decay amplitude into two ordinary mesons can be $1/N^{1/2}$

This discussion has been enlarged by M. Knecht and S. Peris (arXiv:1307.1273) and discussed in three papers by T. Cohen and R. Lebed et al. (arXiv:1401.1815, arXiv:1403.8090). According to them, tetraquarks are not narrow because of $1/N$ counting but due to other effects.

On the other hand tetraquarks appear in the spectrum of QCD in the Corrigan-Ramond large N limit ('larks' in the antifundamental) as *narrow* hadrons.

Z(4430)⁻ at LHCb | April 2014



Signal: 13.9 σ

Other assignments ruled out at 9.7 σ

First observed by BELLE in 2007 and not confirmed by BaBar



About CERN

Students & Educators

Scientists

CERN people

Accelerators

Experiments

Physics

Computing

Engineering

Updates

LHCb confirms existence of exotic hadrons

ABOUT CERN


Cian O'Luanaigh

The [Large Hadron Collider beauty \(LHCb\)](#) collaboration today announced results that confirm the existence of exotic hadrons – a type of matter that cannot be classified within the traditional quark model.

NEWS AND COMMENTARY IN PHYSICS

[Four-Quark State Confirmed](#)

June 4, 2014



The LHCb experiment provides conclusive evidence for the existence of the four-quark particle called $Z(4430)$.

Synopsis on R. Aaij *et al.* (LHCb Collaboration)

[Phys. Rev. Lett. 112, 222002 \(2014\)](#)

Charged $Z_c(3900)$

Found in $Y(4260) \rightarrow Z_c^\pm(3900) \pi^\mp \rightarrow J/\psi \pi^\pm \pi^\mp$

Exotic charged charmonium-like state!

$$G = G_\pi C_{J/\psi} =$$

$$= -1(-1) = +1$$

$$P = +1 \text{ (S-wave)}$$

$$\Rightarrow Z_c^0 \text{ has } J^{PC} = 1^{+-}$$

$$I^G J^{PC} = 1^+ 1^{+-}$$

BESIII, arXiv:1303.5949



$$M = 3899.0 \pm 3.6 \pm 4.9 \text{ MeV}$$

$$\Gamma = 46 \pm 10 \pm 20 \text{ MeV}$$

Belle, arXiv:1304.0121

$$M = 3894.5 \pm 6.6 \pm 4.5 \text{ MeV}$$

$$\Gamma = 63 \pm 24 \pm 26 \text{ MeV}$$



$$Z_c(3900)^{\pm 0}$$

Notes from the Editors: Highlights of the Year

Published December 30, 2013 | *Physics* 6, 139 (2013) | DOI: 10.1103/Physics.6.139

Physics looks back at the standout stories of 2013.

As 2013 draws to a close, we look back on the research covered in *Physics* that really made waves in and beyond the physics community. In thinking about which stories to highlight, we considered a combination of factors: popularity on the website, a clear element of surprise or discovery, or signs that the work could lead to better technology. On behalf of the *Physics* staff, we wish everyone an excellent New Year.


– Matteo Rini and Jessica Thomas

Four-Quark Matter

Quarks come in twos and threes—or so nearly every experiment has told us. This summer, the BESIII Collaboration in China and the Belle Collaboration in Japan reported they had sorted through the debris of high-energy electron-positron collisions and seen a **mysterious particle** that appeared to contain four quarks. Though other explanations for the nature of the particle, dubbed $Z_c(3900)$, are possible, the “tetraquark” interpretation may be gaining traction: BESIII has since **seen** a series of other particles that appear to contain four quarks.

Strangers from Beyond our Solar System

Detector experiments hunting for rare events can go years and never see anything out of the ordinary. So it was cause for excitement when IceCube, a giant neutrino telescope at the South Pole, reported the **detection of two neutrinos with energies of around 1000 tera-electron-volts (TeV)**, roughly a billion times more energetic than those arriving from the Sun. Scientists at IceCube have since further analyzed their data and **reported** 26 more neutrinos with energies above 30 TeV. Researchers will need to observe many more of the neutrinos before they can be certain of their source, and that may require a larger detector. But they believe the particles were produced outside of the Solar System (experiments haven't detected neutrinos from so far away since 1987) and may be carrying information about astrophysical events, like gamma-ray bursts, in distant galaxies.



Images from popular *Physics* stories in 2013.

Article Options

- Printable Version
- Share/Email This
- Export Citation (BibTeX)
- Export Citation (RIS)

New in Physics

- Tuning Ductility**
Synopsis | Mar 19, 2014
- Quantum Superrotor**
Viewpoint | Mar 19, 2014
- The Smallest Shock Wave**
Focus | Mar 18, 2014
- An Untested Window into Nuclear Structure**
Synopsis | Mar 18, 2014
- Tuning in to the Smallest (Man-Made) Mechanical Resonator**
Viewpoint | Mar 17, 2014
- Snapshots from the March Meeting—No-Photon Lasers, Hyperbolic Metamaterials, and More**
Editorial | Mar 17, 2014
- Earth as Giant Gravitational Wave Detector**
Synopsis | Mar 13, 2014

particles in distant galaxies.

of the solar system (experiments haven't detected neutrinos from so far away since 1987) and may be carrying information about astrophysical events, like gamma-ray bursts, in distant galaxies. But they believe the particles were produced outside of the Solar System (experiments haven't detected neutrinos from so far away since 1987) and may be carrying information about astrophysical events, like gamma-ray bursts, in distant galaxies. But they believe the particles were produced outside of the Solar System (experiments haven't detected neutrinos from so far away since 1987) and may be carrying information about astrophysical events, like gamma-ray bursts, in distant galaxies.


Earth as Giant Gravitational Wave Detector
Editorial | Mar 17, 2014
Snapshots from the March Meeting—No-Photon Lasers, Hyperbolic Metamaterials, and More

One or two Z'?

A Pilloni

$Z'_c(4020), Z'_c(4025)$

BESIII, PRL112, 022001



$Z'_c(4025) \rightarrow D^* D^*$ ←

$I^G J^{PC} = 1^+ 1^{+-}$

$M = 4026.3 \pm 4.5 \text{ MeV}$

$\Gamma = 24.8 \pm 9.5 \text{ MeV}$


→ $Z'_c(4020) \rightarrow h_c \pi$

$I^G J^{PC} = 1^+ 1^{\bar{-}}$

$M = 4022.9 \pm 2.8 \text{ MeV}$

$\Gamma = 7.9 \pm 3.7 \text{ MeV}$

BESIII, PRL111, 242001



Radial Excitations?

$$Z(4430) \rightarrow \psi(2S) \pi^-$$

$$Z(3900) \rightarrow J/\psi \pi^-$$

See also the calculation by S. Brodsky, D. Hwang & R. Lebed in a *diquark-antidiquark* model arXiv:1406.7281

$$m(\psi(2S)) - m(J/\psi) \simeq m(Z(4430)) - m(Z(3900))$$

'A crucial consequence of a $Z(4430)$ charged particle is that a charged state decaying into $J/\psi + \pi^\pm$ (or $\eta_c + \rho^\pm$) should be found around 3880 MeV'

Taken from L. Maiani, A. D. Polosa and V. Riquer, arXiv:0708.3997 [hep-ph] At that time there was no hint of $Z_c(3900)$ in data.

$$Z_c(4025) \rightarrow h_c(S_{c\bar{c}} = 0) \pi^-$$

Does this pattern fit in an any extended quark model?

Since 2003/4 new Charmonium-Like States

State	M (MeV)	Γ (MeV)	J^{PC}	Process (decay mode)	Experiment ($\#\sigma$)	1 st observation
$X(3823)$	3823.1 ± 1.9	< 24	$?^{? -}$	$B \rightarrow K + (\chi_{c1} \gamma)$	Belle [4] (3.8)	Belle 2013
$X(3872)$	3871.68 ± 0.17	< 1.2	1^{++}	$B \rightarrow K + (J/\psi \pi^+ \pi^-)$ $p\bar{p} \rightarrow (J/\psi \pi^+ \pi^-) + \dots$ $B \rightarrow K + (J/\psi \pi^+ \pi^- \pi^0)$ $B \rightarrow K + (D^0 \bar{D}^0 \pi^0)$ $B \rightarrow K + (J/\psi \gamma)$ $B \rightarrow K + (\psi(2S) \gamma)$ $pp \rightarrow (J/\psi \pi^+ \pi^-) + \dots$	Belle [5, 6] (12.8), BABAR [7] (8.6) CDF [8–10] (np), DØ [11] (5.2) Belle [12] ^a (4.3), BABAR [13] ^a (4.0) Belle [14, 15] ^a (6.4), BABAR [16] ^a (4.9) Belle [17] ^a (4.0), BABAR [18, 19] ^a (3.6) BABAR [19] ^a (3.5), Belle [17] ^a (0.4) LHCb [20] (np)	Belle 2003
$X(3915)$	3917.5 ± 1.9	20 ± 5	0^{++}	$B \rightarrow K + (J/\psi \omega)$ $e^+ e^- \rightarrow e^+ e^- + (J/\psi \omega)$	Belle [21] (8.1), BABAR [22] (19) Belle [23] (7.7), BABAR [13, 24] (7.6)	Belle 2004
$\chi_{c2}(2P)$	3927.2 ± 2.6	24 ± 6	2^{++}	$e^+ e^- \rightarrow e^+ e^- + (D\bar{D})$	Belle [25] (5.3), BABAR [26] (5.8)	Belle 2005
$X(3940)$	3942_{-8}^{+9}	37_{-17}^{+27}	$?^{? +}$	$e^+ e^- \rightarrow J/\psi + (D^* \bar{D})$ $e^+ e^- \rightarrow J/\psi + (\dots)$	Belle [27] (6.0) Belle [28] (5.0)	Belle 2007
$G(3900)$	3943 ± 21	52 ± 11	1^{--}	$e^+ e^- \rightarrow \gamma + (D\bar{D})$	BABAR [29] (np), Belle [30] (np)	BABAR 2007
$Y(4008)$	4008_{-49}^{+121}	226 ± 97	1^{--}	$e^+ e^- \rightarrow \gamma + (J/\psi \pi^+ \pi^-)$	Belle [31] (7.4)	Belle 2007
$Y(4140)$	4144.5 ± 2.6	15_{-7}^{+11}	$?^{? +}$	$B \rightarrow K + (J/\psi \phi)$	CDF [32, 33] (5.0), CMS [34] (>5)	CDF 2009
$X(4160)$	4156_{-25}^{+29}	139_{-65}^{+113}	$?^{? +}$	$e^+ e^- \rightarrow J/\psi + (D^* \bar{D}^*)$	Belle [27] (5.5)	Belle 2007


^a Not included in the averages for M and Γ .

New Charmonium & Bottomonium Like States

State	M (MeV)	Γ (MeV)	J^{PC}	Process (decay mode)	Experiment ($\# \sigma$)	1 st observation
Y(4260)	4263_{-9}^{+8}	95 ± 14	1^{--}	$e^+e^- \rightarrow \gamma + (J/\psi \pi^+ \pi^-)$ $e^+e^- \rightarrow (J/\psi \pi^+ \pi^-)$ $e^+e^- \rightarrow (J/\psi \pi^0 \pi^0)$	BABAR [35, 36] (8.0), CLEO [37] (5.4) Belle [31] (15) CLEO [38] (11) CLEO [38] (5.1)	BABAR 2005
Y(4274)	$4274.4_{-6.7}^{+8.4}$	32_{-15}^{+22}	$?^?+$	$B \rightarrow K + (J/\psi \phi)$	CDF [33] (3.1)	CDF 2010
X(4350)	$4350.6_{-5.1}^{+4.6}$	$13.3_{-10.0}^{+18.4}$	$0/2^{++}$	$e^+e^- \rightarrow e^+e^- (J/\psi \phi)$	Belle [39] (3.2)	Belle 2009
Y(4360)	4361 ± 13	74 ± 18	1^{--}	$e^+e^- \rightarrow \gamma + (\psi(2S) \pi^+ \pi^-)$	BABAR [40] (np), Belle [41] (8.0)	BABAR 2007
X(4630)	4634_{-11}^{+9}	92_{-32}^{+41}	1^{--}	$e^+e^- \rightarrow \gamma (\Lambda_c^+ \Lambda_c^-)$	Belle [42] (8.2)	Belle 2007
Y(4660)	4664 ± 12	48 ± 15	1^{--}	$e^+e^- \rightarrow \gamma + (\psi(2S) \pi^+ \pi^-)$	Belle [41] (5.8)	Belle 2007
Z_c^+(3900)	3898 ± 5	51 ± 19	1^{7-}	$Y(4260) \rightarrow \pi^- + (J/\psi \pi^+)$ $e^+e^- \rightarrow \pi^- + (J/\psi \pi^+)$	BESIII [43] (np), Belle [44] (5.2) Xiao <i>et al.</i> [45] ^a (6.1)	BESIII 2013
Z_1^+ (4050)	4051_{-43}^{+24}	82_{-55}^{+51}	$?$	$B \rightarrow K + (\chi_{c1}(1P) \pi^+)$	Belle [46] (5.0), BABAR [47] (1.1)	Belle 2008
Z_2^+ (4250)	4248_{-45}^{+185}	177_{-72}^{+321}	$?$	$B \rightarrow K + (\chi_{c1}(1P) \pi^+)$	Belle [46] (5.0), BABAR [47] (2.0)	Belle 2008
Z^+ (4430)	4443_{-18}^{+24}	107_{-71}^{+113}	$?$	$B \rightarrow K + (\psi(2S) \pi^+)$	Belle [48, 49] (6.4), BABAR [50] (2.4)	Belle 2007
$Y_b(10888)$	10888.4 ± 3.0	$30.7_{-7.7}^{+8.9}$	1^{--}	$e^+e^- \rightarrow (\Upsilon(nS) \pi^+ \pi^-)$	Belle [51, 52] (2.0)	Belle 2010
$Z_b^+(10610)$	10607.2 ± 2.0	18.4 ± 2.4	1^{+-}	$\Upsilon(5S) \rightarrow \pi^- + (\Upsilon(nS) \pi^+), n = 1, 2, 3$ $\Upsilon(5S) \rightarrow \pi^- + (h_b(nP) \pi^+), n = 1, 2$	Belle [53, 54] (16) Belle [53, 54] (16)	Belle 2011
$Z_b^+(10650)$	10652.2 ± 1.5	11.5 ± 2.2	1^{+-}	$\Upsilon(5S) \rightarrow \pi^- + (\Upsilon(nS) \pi^+), n = 1, 2, 3$ $\Upsilon(5S) \rightarrow \pi^- + (h_b(nP) \pi^+), n = 1, 2$	Belle [53, 54] (16) Belle [53, 54] (16)	Belle 2011

^aNot included in the averages for M and Γ .

Mass Matchings




$$\Upsilon(5S) \rightarrow \pi^\pm Z_b^\mp(10610) \rightarrow \pi^\pm \pi^\mp \Upsilon(nS) \quad n = 1, 2, 3$$

$$\Upsilon(5S) \rightarrow \pi^\pm Z_b^\mp(10650) \rightarrow \pi^\pm \pi^\mp h_b(kP) \quad k = 1, 2$$

$$m_B + m_{B^*} \simeq 10604 \text{ MeV}$$

$$2m_{B^*} \simeq 10650 \text{ MeV}$$



$$Z_c(3900) \rightarrow \pi^\pm J/\psi$$

$$Z'_c(4025) \rightarrow h_c \pi^\pm$$

$$m_{D^0} + m_{D^{*+}} = 3875 \text{ MeV}$$

$$m_{D^{*0}} + m_{D^{*+}} = 4017 \text{ MeV}$$

$$+24 \text{ MeV}$$

$$+8 \text{ MeV}$$

Moreover: no molecular matchings for the Z(4430)

X(3872), $J^{PC} = 1^{++}$

Discovered by **Belle**, and soon confirmed by CDF, BaBar, D0. Later observed at CMS and ATLAS. Both produced in B meson decays and *prompt*, in hadron collisions. Four-Quark matter as well? Its isospin seems to be $I=0$ — *whereas the diquark model would naively predict $I=1$ (just change the flavors of light quarks q & q').*

4 CMS Collaboration arXiv:1302.3968

4 Measurement of the cross section ratio




Figure 1: The $J/\psi\pi^+\pi^-$ invariant-mass spectrum for $10 < p_T < 50$ GeV and $|y| < 1.2$. The lines represent the signal-plus-background fits (solid), the background-only (dashed), and the signal-only (dotted) components. The inset shows an enlargement of the X(3872) mass region.

Figure 1: The $J/\psi\pi^+\pi^-$ invariant-mass spectrum for $10 < p_T < 50$ GeV and $|y| < 1.2$. The lines represent the signal-plus-background fits (solid), the background-only (dashed), and the signal-only (dotted) components. The inset shows an enlargement of the X(3872) mass region.

X: Mass and Isospin Problems

Similarly to what happens to Z_c , Z_c' and Z_b , also the $X(3872)$ appears to be very close to the the closeby (and quantum numbers allowed) open charm threshold

$$m_{D^0} + m_{D^{*0}} = 3872 \text{ MeV}$$

This time the coincidence is really striking because the value is just exactly there. Actually in terms of mass there is another surprising 'coincidence' in the X case

$$m_{J/\psi} + m_{\rho^0} = 3872 \text{ MeV}$$


The X decays in both channels, preferring the first one, and **also decays into $J/\psi \omega$**

$$\frac{\mathcal{B}(X \rightarrow J/\psi \rho)}{\mathcal{B}(X \rightarrow J/\psi \omega)} \approx 1$$

which is an interesting hint of *isospin violation*

Extreme Interpretations of X

and of all other XYZ states



A compact 'tetraquark'

Jaffe & Wilczek

Maiani, Piccinini, Polosa, Riquer '2005

Terasaki, Ali, Ebert...

Approach based on SU(3) symmetry



A loosely bound molecule


Tornqvist; Braaten, Kusinoki; Voloshin; Barnes;

Swanson; Close; Mehen, Fleming; Hanhart, Guo, Meissner
very long list...'2005 —

Based on strong hadron dynamics

Diquarks

Diquarks



$$H = \sum_i m_i + \sum_{i < j} 2\kappa_{ij} \mathbf{S}_i \cdot \mathbf{S}_j$$

The spin interactions inside the diquark are assumed to dominate over all other possible pairings

$$H \approx 2\kappa_{q\bar{q}}(s_q \cdot s_{\bar{q}}) \text{ type I } \text{Maiani, Piccinini, Polosa, Riquer, PRD71 (2005)}$$

$$H \approx 2\kappa_{q\bar{c}}(s_q \cdot s_c + s_{\bar{q}} \cdot s_{\bar{c}}) \text{ type II } \text{Maiani, Piccinini, Polosa, Riquer, PRD89 (2014)}$$

SU(3) One-Gluon-Exchange

$$T_{R_1}^a \otimes T_{R_2}^a = \bigoplus_i \frac{1}{2} (C_{S_i} - C_{R_1} - C_{R_2}) \mathbb{1}_{S_i}$$

$$\begin{aligned} T_{\mathbf{3}}^a \otimes T_{\mathbf{3}}^a &= \frac{1}{2} (4/3 - 4/3 - 4/3) \mathbb{1}_{\bar{\mathbf{3}}} \oplus \\ &\quad \oplus \frac{1}{2} (10/3 - 4/3 - 4/3) \mathbb{1}_{\mathbf{6}} = \\ &= -\frac{2}{3} \mathbb{1}_{\bar{\mathbf{3}}} \oplus \frac{1}{3} \mathbb{1}_{\mathbf{6}} \end{aligned}$$

$$T_{\mathbf{3}}^a \otimes T_{\bar{\mathbf{3}}}^a = -\frac{4}{3} \mathbb{1}_{\mathbf{1}} \oplus \frac{1}{6} \mathbb{1}_{\mathbf{8}}$$

Diquark spin basis

$|s, \bar{s}\rangle_J$ notation

$$J^P = 0^+ \quad C = + \quad X_0 = |0, 0\rangle_0, \quad X'_0 = |1, 1\rangle_0 \quad X(3915), X(3940) ?$$

$$J^P = 1^+ \quad C = + \quad X_1 = \frac{1}{\sqrt{2}} (|1, 0\rangle_1 + |0, 1\rangle_1) \quad X(3872)$$

$$J^P = 1^+ \quad G = + \quad Z = \frac{1}{\sqrt{2}} (|1, 0\rangle_1 - |0, 1\rangle_1), \quad Z' = |1, 1\rangle_1$$


$$J^P = 2^+ \quad C = + \quad X_2 = |1, 1\rangle_2 \quad \text{Linear combi } Z, Z' \text{ to make } Z_c\text{'s}$$

The Hamiltonian, of the 'type I' diquark model, turns out to be diagonal on 1^{++} and 2^{++} states but *not* on 1^{+-} and 0^{++} .

In the 'type II' model the mass eigenvectors coincide with the listed states and **$M(X_1) \sim M(Z), M(Z') - M(Z) \sim 134 \text{ MeV}$** .

Spectrum

Maiani, Piccinini, Polosa, Riquer, PRD89 (2014)



1_{BC}

0_{++}


1_{++}

1_{+-}

2_{++}





3000

Since 2003/4 new Charmonium-Like States

State	M (MeV)	Γ (MeV)	J^{PC}	Process (decay mode)	Experiment ($\#\sigma$)	1 st observation
$X(3823)$	3823.1 ± 1.9	< 24	$?^{? -}$	$B \rightarrow K + (\chi_{c1} \gamma)$	Belle [4] (3.8)	Belle 2013
$X(3872)$	3871.68 ± 0.17	< 1.2	1^{++}	$B \rightarrow K + (J/\psi \pi^+ \pi^-)$ $p\bar{p} \rightarrow (J/\psi \pi^+ \pi^-) + \dots$ $B \rightarrow K + (J/\psi \pi^+ \pi^- \pi^0)$ $B \rightarrow K + (D^0 \bar{D}^0 \pi^0)$ $B \rightarrow K + (J/\psi \gamma)$ $B \rightarrow K + (\psi(2S) \gamma)$ $pp \rightarrow (J/\psi \pi^+ \pi^-) + \dots$	Belle [5, 6] (12.8), BABAR [7] (8.6) CDF [8–10] (np), DØ [11] (5.2) Belle [12] ^a (4.3), BABAR [13] ^a (4.0) Belle [14, 15] ^a (6.4), BABAR [16] ^a (4.9) Belle [17] ^a (4.0), BABAR [18, 19] ^a (3.6) BABAR [19] ^a (3.5), Belle [17] ^a (0.4) LHCb [20] (np)	Belle 2003
$X(3915)$	3917.5 ± 1.9	20 ± 5	0^{++}	$B \rightarrow K + (J/\psi \omega)$ $e^+ e^- \rightarrow e^+ e^- + (J/\psi \omega)$	Belle [21] (8.1), BABAR [22] (19) Belle [23] (7.7), BABAR [13, 24] (7.6)	Belle 2004
$\chi_{c2}(2P)$	3927.2 ± 2.6	24 ± 6	2^{++}	$e^+ e^- \rightarrow e^+ e^- + (D\bar{D})$	Belle [25] (5.3), BABAR [26] (5.8)	Belle 2005
$X(3940)$	3942_{-8}^{+9}	37_{-17}^{+27}	$?^{? +}$	$e^+ e^- \rightarrow J/\psi + (D^* \bar{D})$ $e^+ e^- \rightarrow J/\psi + (\dots)$	Belle [27] (6.0) Belle [28] (5.0)	Belle 2007
$G(3900)$	3943 ± 21	52 ± 11	1^{--}	$e^+ e^- \rightarrow \gamma + (D\bar{D})$	BABAR [29] (np), Belle [30] (np)	BABAR 2007
 $Y(4008)$	4008_{-49}^{+121}	226 ± 97	1^{--}	$e^+ e^- \rightarrow \gamma + (J/\psi \pi^+ \pi^-)$	Belle [31] (7.4)	Belle 2007
$Y(4140)$	4144.5 ± 2.6	15_{-7}^{+11}	$?^{? +}$	$B \rightarrow K + (J/\psi \phi)$	CDF [32, 33] (5.0), CMS [34] (>5)	CDF 2009
$X(4160)$	4156_{-25}^{+29}	139_{-65}^{+113}	$?^{? +}$	$e^+ e^- \rightarrow J/\psi + (D^* \bar{D}^*)$	Belle [27] (5.5)	Belle 2007

^a Not included in the averages for M and Γ .

New Charmonium & Bottomonium Like States

State	M (MeV)	Γ (MeV)	J^{PC}	Process (decay mode)	Experiment ($\# \sigma$)	1 st observation
 Y(4260)	4263_{-9}^{+8}	95 ± 14	1^{--}	$e^+e^- \rightarrow \gamma + (J/\psi \pi^+ \pi^-)$ $e^+e^- \rightarrow (J/\psi \pi^+ \pi^-)$ $e^+e^- \rightarrow (J/\psi \pi^0 \pi^0)$	BABAR [35, 36] (8.0), CLEO [37] (5.4) Belle [31] (15) CLEO [38] (11) CLEO [38] (5.1)	BABAR 2005
Y(4274)	$4274.4_{-6.7}^{+8.4}$	32_{-15}^{+22}	$?^{?+}$	$B \rightarrow K + (J/\psi \phi)$	CDF [33] (3.1)	CDF 2010
X(4350)	$4350.6_{-5.1}^{+4.6}$	$13.3_{-10.0}^{+18.4}$	$0/2^{++}$	$e^+e^- \rightarrow e^+e^- (J/\psi \phi)$	Belle [39] (3.2)	Belle 2009
 Y(4360)	4361 ± 13	74 ± 18	1^{--}	$e^+e^- \rightarrow \gamma + (\psi(2S) \pi^+ \pi^-)$	BABAR [40] (np), Belle [41] (8.0)	BABAR 2007
 X(4630)	4634_{-11}^{+9}	92_{-32}^{+41}	1^{--}	$e^+e^- \rightarrow \gamma (\Lambda_c^+ \Lambda_c^-)$	Belle [42] (8.2)	Belle 2007
 Y(4660)	4664 ± 12	48 ± 15	1^{--}	$e^+e^- \rightarrow \gamma + (\psi(2S) \pi^+ \pi^-)$	Belle [41] (5.8)	Belle 2007
Z_c⁺(3900)	3898 ± 5	51 ± 19	1^{7-}	$Y(4260) \rightarrow \pi^- + (J/\psi \pi^+)$ $e^+e^- \rightarrow \pi^- + (J/\psi \pi^+)$	BESIII [43] (np), Belle [44] (5.2) Xiao <i>et al.</i> [45] ^a (6.1)	BESIII 2013
Z ₁ ⁺ (4050)	4051_{-43}^{+24}	82_{-55}^{+51}	?	$B \rightarrow K + (\chi_{c1}(1P) \pi^+)$	Belle [46] (5.0), BABAR [47] (1.1)	Belle 2008
Z ₂ ⁺ (4250)	4248_{-45}^{+185}	177_{-72}^{+321}	?	$B \rightarrow K + (\chi_{c1}(1P) \pi^+)$	Belle [46] (5.0), BABAR [47] (2.0)	Belle 2008
Z ⁺ (4430)	4443_{-18}^{+24}	107_{-71}^{+113}	?	$B \rightarrow K + (\psi(2S) \pi^+)$	Belle [48, 49] (6.4), BABAR [50] (2.4)	Belle 2007
Y _b (10888)	10888.4 ± 3.0	$30.7_{-7.7}^{+8.9}$	1^{--}	$e^+e^- \rightarrow (\Upsilon(nS) \pi^+ \pi^-)$	Belle [51, 52] (2.0)	Belle 2010
Z _b ⁺ (10610)	10607.2 ± 2.0	18.4 ± 2.4	1^{+-}	$\Upsilon(5S) \rightarrow \pi^- + (\Upsilon(nS) \pi^+), n = 1, 2, 3$ $\Upsilon(5S) \rightarrow \pi^- + (h_b(nP) \pi^+), n = 1, 2$	Belle [53, 54] (16) Belle [53, 54] (16)	Belle 2011
Z _b ⁺ (10650)	10652.2 ± 1.5	11.5 ± 2.2	1^{+-}	$\Upsilon(5S) \rightarrow \pi^- + (\Upsilon(nS) \pi^+), n = 1, 2, 3$ $\Upsilon(5S) \rightarrow \pi^- + (h_b(nP) \pi^+), n = 1, 2$	Belle [53, 54] (16) Belle [53, 54] (16)	Belle 2011

^aNot included in the averages for M and Γ .

Y States: One Unit of L

$$|s, \bar{s}; S, L\rangle_{J=1}$$

$$Y_1 = |0, 0; 0, 1\rangle_1$$

$$Y(4260) \text{ like } X(3872) \longrightarrow Y_2 = \frac{1}{\sqrt{2}} (|1, 0; 1, 1\rangle_1 + |0, 1; 1, 1\rangle_1)$$

$$Y_3 = |1, 1; 0, 1\rangle_1$$

$$Y_4 = |1, 1; 2, 1\rangle_1$$

				1^{--}
		$Y_1 : P(s_{c\bar{c}} = 1) : P(s_{c\bar{c}} = 0) = 3 : 1$		$Y(4008)$
$Y(4260) \rightarrow \gamma + X(3872)$	\longrightarrow	$Y_2 : P(s_{c\bar{c}} = 1) = 1$		$Y(4260)$
$h_c(1P)$	\longrightarrow	$Y_3 : P(s_{c\bar{c}} = 1) : P(s_{c\bar{c}} = 0) = 1 : 3$		$Y(4290) \text{ or } Y(4220)$
		$Y_4 : P(s_{c\bar{c}} = 1) = 1$		$Y(4630)$

Tentatively we identify $Y(4360)$ and $Y(4660)$ decaying into $\psi(2S)$ as radial excitations of $Y(4008)$ and $Y(4260)$.




Charmed Baryonium

G. Cotugno, R. Faccini, ADP, C. Sabelli *Phys. Rev. Lett.* **104**, 132005 (2010)

One may easily suspect that $Y(4660)$ and $Y(4630)$ are one and the same particle (call it Y_B) showing how this hypothesis improves the fit to Belle data.

Under this hypothesis we found the remarkable ratio

$$\frac{\mathcal{B}(Y_B \rightarrow \Lambda_c \bar{\Lambda}_c)}{\mathcal{B}(Y_B \rightarrow \psi(2S) \pi^+ \pi^-)} = 24.6 \pm 6.6$$



Molecules

Enter a citation to look up or terms to search. ✕

Ex: "PRL 112 068103", "Phys. Rev. Lett. 112, 068103",
"10.1103/PhysRevLett.112.068103"

Close

Molecular Charmonium: A New Spectroscopy?

Phys. Rev. Lett. **38**, 317 – Published 14 February 1977

A. De Rújula, Howard Georgi, and S. L. Glashow

PDF

Export Citation

Citing Articles (149)

 Like 0

 Tweet 0

ABSTRACT


AUTHORS

REFERENCES

ABSTRACT

Recent data compel us to interpret several peaks in the cross section of e^-e^+ annihilation into hadrons as being due to the production of four-quark molecules, i.e., resonances between two charmed mesons. A rich spectroscopy of such states is predicted and may be studied in e^-e^+ annihilation.


IS THE X(3872) SOME SORT OF DD* DEUTERON?



$$k_{\text{rel}} = \sqrt{2\mu\langle T \rangle_{\psi}^2} \approx \begin{cases} 80 \text{ MeV} & \text{for deuterium} \\ 50 \text{ MeV} & \text{for } X; \quad U_0 \approx -7 \text{ MeV} \quad \mathcal{E}_b \approx -0.14 \text{ MeV} \end{cases}$$


$$\frac{\hbar^2}{2\mu r_0^2} - \frac{g^2}{4\pi} \frac{e^{-\frac{m_{\pi}c}{\hbar} r_0}}{r_0} = \mathcal{E}_b = 0.14 \text{ MeV} \Rightarrow r_0 \approx 12 \text{ fm}$$

Barely Bound States in TeV Hadron Collisions?



$$p_{\perp}^{\text{mol}} > 5.5 \text{ GeV}$$

$$|y^{\text{mol}}| < 1$$



Production xsect 300 times smaller than the observed one

C. Bignamini, B. Grinstein, F. Piccinini, ADP, C. Sabelli, *Phys Rev Lett*, 103, 162001 (2009)


P. Artoisenet and E. Braaten, *Phys Rev D*81, 114018 (2010)

C. Bignamini, B. Grinstein, F. Piccinini, ADP, C. Sabelli, *Phys Lett*, B684, 228 (2010)

A. Esposito, F. Piccinini, A. Pilloni, A.D. Polosa, *J. Mod. Phys.* 4, 1569, (2013)

F-K. Guo, U. Meissner and Wang, arXiv: 1308.0193, 1402.6236


Production: MC Tuning



red: cc^* HERWIG/PYTHIA
 green: $cc^*g(\text{recoiling})$ ALPGEN + HERWIG/PYTHIA
 blue: full qcd HERWIG/PYTHIA


Final state πD scatterings?

Hadronization produces π mesons




How many π mesons in a spherical sector of 3 fm of base radius around D ?
Possible multiple interactions to correct the k_0 ? Not many! Estimate ~ 3

Complanar Pions




$$\tan \theta = \frac{v_0 \sin \theta_0 \sqrt{1 - V^2}}{V + v_0 \cos \theta_0}$$




Any Improvements?

The mechanism works: *feed down* from higher bins — but it does not help in the *bins of interest* (up to 100 MeV for the com relative momentum in the wold-be-molecule, k_0)




Deuteron at LHC?

Usual reply: A lot!!
Indeed Alice has 30K antideuterons.
— In which p_T range though?







Recall that X has been observed with a $p_T > 5.5$ GeV!



$$\left(\frac{k_0}{2}\right)_{\max} \approx (|\mathbf{p}| \sin \varphi/2)_{\max} \approx p_{\perp}$$



MC Extrapolation



Reverse MC Extrapolation



Selection Rules

Hadronization must be 4q

$$|\psi\rangle = \alpha|[Qq]_{\bar{\mathbf{3}}_c}[\bar{Q}\bar{q}]_{\mathbf{3}_c}\rangle_c + \beta|(Q\bar{Q})_{\mathbf{1}_c}(q\bar{q})_{\mathbf{1}_c}\rangle_o + \gamma|(Q\bar{q})_{\mathbf{1}_c}(\bar{Q}q)_{\mathbf{1}_c}\rangle_o$$

- All 'woud-be' loosely bound molecules do not form any bound state.
- Sometimes a compact 4quark state is formed, but it could be that $|\alpha| < |\beta|, |\gamma|$
- An amplification mechanism might be at work when the closed channel level matches the onset of the continuum spectrum of two mesons with the same quantum numbers.

BTW observe that from an estimate by Ali & Wang


$$\frac{\sigma(pp \rightarrow Y(4260))}{\sigma(pp \rightarrow X(3872))} \gtrsim 10^{-2}$$

It would be very significant to directly measure this ratio.

Do we know 'amplification' mechanisms between open/closed channels?

Another Mechanism

Borrow some ideas from cold atom physics. The Fano-Feshbach mechanism.




*e.g. DD^**

$\epsilon \cdot \hat{d} \cdot DD_$*

Another Mechanism


Borrow some ideas from cold atom physics. The Fano-Feshbach mechanism.



$$a \sim |C| \sum_n \frac{c \langle [Qq]_{\mathbf{3}_c} [\bar{Q}\bar{q}]_{\mathbf{3}_c}, n | H_{cO} | (Q\bar{q})_{\mathbf{1}_c} (\bar{Q}q)_{\mathbf{1}_c} \rangle_0}{E_0 - E_n}$$

Another Mechanism

Consider also that the $J/\psi \rho^+$ is sensibly lower than the related open charm charged molecule. This could be why there is no charged X and I -violat.



Lattice

4-quarks from lattice?

Esposito, Papinutto, Piloni, ADP, Tantalo Phys.Rev. D88 (2013) 054029

On simulating a proton on the lattice, the interpolating operators

$$O = \epsilon^{abc} u^a u^b d^c, \quad \epsilon^{abc} u^a u^b d^c \bar{s}^d s^d \dots$$

are equally good. One might wonder if there is any chance of studying genuine tetraquark configurations on the lattice as they might turn out not to be distinguishable from standard charmonia.

On the other hand *states with two charm quarks* cannot mix with standard charmonia.

$$\mathcal{O}_{T^0}^1 = \epsilon^{ABC} c^B c^C \epsilon^{ADE} \bar{u}^D \bar{u}^E$$

$$\mathcal{O}_{T^0}^2 = \bar{u}^A c^A \bar{u}^B c^B$$

$$\mathcal{O}_{T^0}^3 = \sum_a \bar{u}^A T_i^{AB} c^B \bar{u}^C T_a^{CD} c^D$$

$$\mathcal{O}_{T^0}^4 = \bar{u}^A c^A \bar{u}^B c^B \bar{s}^C s^C$$

More Exotic States

$$|T^0\rangle = |Q_u = -2, Q_c = +2\rangle$$

$$|T^+\rangle = |Q_u = -1, Q_d = -1, Q_c = +2\rangle$$

$$|T_s^+\rangle = |Q_u = -1, Q_s = -1, Q_c = +2\rangle$$

$$|T^{++}\rangle = |Q_d = -2, Q_c = +2\rangle$$


$$|T_s^{++}\rangle = |Q_s = -2, Q_c = +2\rangle$$

Production from heavy baryons

	Bottom quark decays	
Starting baryon	$b \rightarrow c\bar{u}d$ ($O(\lambda^2)$)	$b \rightarrow c\bar{u}s$ ($O(\lambda^3)$)
Ξ_{bc}^+ [bcu]	$pT^0 \rightarrow pD^0D^0$ $nT^+ \rightarrow nD^0D^+$ $\Lambda^0(\Sigma^0)T_s^+ \rightarrow \Lambda^0(\Sigma^0)D^0D_s^+$	$\Sigma^+T^0 \rightarrow \Sigma^+D^0D^0$ $\Lambda^0(\Sigma^0)T^+ \rightarrow \Lambda^0(\Sigma^0)D^0D^+$ $\Xi^0T_s^+ \rightarrow \Xi^0D_s^+D^0$
Ξ_{bc}^0 [bcd]	$nT^0 \rightarrow nD^0D^0$ $\Delta^-T^+ \rightarrow \Delta^-D^+D^0$ $\Sigma^-T_s^+ \rightarrow \Sigma^-D_s^+D^0$	$\Lambda^0(\Sigma^0)T^0 \rightarrow \Lambda^0(\Sigma^0)D^0D^0$ $\Sigma^-T^+ \rightarrow \Sigma^-D^+D^0$ $\Xi^-T_s^+ \rightarrow \Xi^-D_s^+D^0$
Ξ_{bcs}^0 [bcs]	Same final states as [bcd] with $b \rightarrow c\bar{u}s$ (they differ by just $d \leftrightarrow s$)	$\Xi^0T^0 \rightarrow \Xi^0D^0D^0$ $\Xi^-T^+ \rightarrow \Xi^-D^+D^0$ $\Omega^-T_s^+ \rightarrow \Omega^-D_s^+D^0$

Lattice Spectrum Studies

$$m_{\text{eff}} = -\ln \frac{C(x_0 + 1)}{C(x_0)}$$




Conclusions

- **Multiquarks hadrons have been discovered!** We need a coherent description which aims to explain the largest number/totality of collected data.
- **Loosely bound molecules** promptly produced at high energy hadron colliders require a miracle in final state interactions.
- **Diquark-antidiquark** states give an easy understanding of prompt production but require a richer flavor structure induced by SU(3) symmetry.
- Is the **quantum mechanics** of Feshbach resonances a possible solution? Not with standard charmonia vs open charm molecules. Most likely the closed channel has to be diquark-antidiquark.

Backup Slides

Can Alice add knowledge on X(3872)?

Let's consider the recombination mechanism (Muller et al.)



The phase space is densely populated of partons. The **soft** part of the spectrum of produced hadrons is assumed to be formed by the decay of a deconfined state of constituent quarks which '**recombine**'. The hard part is instead described by pQCD with an implementation of the parton energy loss in medium. This simple picture turns out to be effective at explaining the observation of

$$\frac{\#p}{\#\pi^+} \geq 1 \text{ for } 1.5 \text{ GeV} \leq p_{\perp} \leq 4 \text{ GeV @ RHIC}$$

this cannot be explained with fragmentation functions (**fitted** at e+e-) where the number of expected pions is much higher.

- I - The transverse momentum of partons is steeply falling with p_T
- II - Fragmentation functions favor the situation where the energy of the fragmenting parton is *not* concentrated in a *single hadron* but distributed amid all *radiated partons*.

Thus fragmentation is inefficient at producing high p_T hadrons in general. In particular pions are favored with respect to baryons .

But if th phase space is highly populated with partons, then it can simply happen that

$$P_\pi = p_u + p_{\bar{d}}$$

$$P_p = p_u + p_u + p_d$$

Assume


$$p_u \simeq p_d \simeq p_{\bar{d}} \simeq p$$

$$\text{Prob}(N\text{-quark hadron}(p)) \propto (\text{Prob}(\text{quark}(q)))^N$$

and assume also exponentially falling parton spectra: same yields at the same hadron transverse momenta

$$\#(\text{protons}) \propto (\exp(-p_T/3))^3 \simeq \#(\text{pions}) \propto (\exp(-p_T/2))^2$$

As for the recombination mechanism, we might expect the X should be produced with similar yields as other charmonia/bottomonia (either 4q or molecule interpretations are assumed). Charm enhancement at ALICE? Then X enhancement as well!




Maiani, Polosa, Riquer, Salgado arXiv:0707.4578 [hep-ph]

Fragmentation functions should be modified for a tetraquark X as diquarks are produced whereas the hard part of the spectrum should be left unchanged for the molecule.

Exotic hadrons and HIC

- Model the fragmentation functions of diquark-antidiquark mesons. Confronted with data from ALICE.




The numerators are dominated by the 'coalescence' mechanism (B. Muller et al). Molecule and tetraquark denominators should also be different

$$R_{CP} = \frac{N_{\text{coll}}(b)}{N_{\text{coll}}(b=0)} \left(\frac{dN_H/d^2p_{\perp}(b=0)}{dN_H/d^2p_{\perp}(b)} \right)$$

$$R_{AA} = \frac{1}{N_{\text{coll}}(b=0)} \left(\frac{dN_H/d^2p_{\perp}(b=0)}{dN_H/d^2p_{\perp}|pp} \right)$$

CHARMED DIQUARKS

The octet with diquarks -
the 'azimuthal approach'



$$\mathcal{Q}_{i\alpha} = \epsilon_{ijk} \epsilon_{\alpha\beta\gamma} \bar{q}_C^{j\beta} \gamma_5 q^{k\gamma} = [qq]_0$$

$$\mathcal{Q}_\alpha^{jk} = \epsilon_{\alpha\beta\gamma} \bar{q}_C^{\beta(j} \vec{\gamma} q^{k)\gamma} = [qq]_1$$

J^{PC}	$dq-dq^*$
0^{++}	$[cq]_0[\bar{c}\bar{q}]_0 \vee ([cq]_1[\bar{c}\bar{q}]_1)_0$
1^{++}	$\frac{[cq]_1[\bar{c}\bar{q}]_0 + [cq]_0[\bar{c}\bar{q}]_1}{\sqrt{2}}$
1^{+-}	$\frac{[cq]_1[\bar{c}\bar{q}]_0 - [cq]_0[\bar{c}\bar{q}]_1}{\sqrt{2}} \vee ([cq]_1[\bar{c}\bar{q}]_1)_1$
2^{++}	$([cq]_1[\bar{c}\bar{q}]_1)_2$

$$([\]_s [\]_s)_J$$

ISOSPIN VIOLATIONS

We set in the flavor basis X_u, X_d

$$M = \begin{pmatrix} 2m_u & 0 \\ 0 & 2m_d \end{pmatrix} + \delta \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$$

where the mixing matrix has a diagonal structure in the Isospin $I = 0, 1$ basis, its eigenvectors being

$$\frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

At the charmonium scale we expect the annihilations to be small and quark mass to dominate - observed $X \rightarrow \omega / \rho$ isospin breaking


Learn from Hadron Multiplicity

$$\rho(m) = \# \text{ of states in } (m, m+dm) = \sum_{m^*} \delta(m - m^*) \rightarrow$$

$$\rightarrow \sum_{m^* = m_f, m_\rho, m_\omega, \dots} \frac{g_{m^*}}{\sigma_{m^*} \sqrt{2\pi}} \exp\left(-\frac{(m - m^*)^2}{2\sigma_{m^*}^2}\right)$$

1996 (4627 states)

$$\sigma_{m^*} = \Gamma_{m^*} / 2 \sim O(100) \text{ MeV}$$



Hadron Gas

The partition function of a hot gas of hadrons could be written using the degeneracy factor as extrapolated by the exponential density of hadrons

$$\begin{aligned} Z &= \sum_E \underbrace{g(E)}_{\text{degeneracy of } E} e^{-E/T} = \sum_m \rho(m) \sum_p e^{-\frac{\sqrt{p^2+m^2}}{T}} \\ &\simeq \frac{V}{(2\pi)^3} \int dm \rho(m) e^{-\frac{m}{T}} \int dp p^2 e^{-\frac{p^2}{2mT}} \\ &= V \left(\frac{T}{2\pi} \right)^{3/2} \int dm \frac{a}{(m^2 + m_0^2)^{3/2}} m^{3/2} e^{m \left(\frac{T-T_H}{TT_H} \right)} \\ &\approx A + B|T - T_H|^{1/2} \end{aligned}$$

Hadron matter cannot exist at temperature $T > T_H$ (the partition function would not be convergent). As T approaches T_H , *from below*, heavier and heavier resonances *enter in the sum* — while their momenta do not increase.



T_H must be the *ultimate temperature of hadronic matter*, where relevant dof must change (N. Cabibbo & G. Parisi 1975)

LOW EQUATION


When shallow bound states are allowed in low energy potential scattering it is possible a description of scattering lengths and phase shifts which does not require the precise knowledge of the scattering potential.

$$\sigma_{\text{tot}} = \frac{2\pi\hbar^2}{\mu|E_b|}$$

DD* Luminosity; $k_0 < \Lambda$



vs. a ~ 12 fm from Low Eq.



The coupling is extracted from the X decay width into DD^* components; yet the binding energy is too low.

X at LHCb

LHCb arXiv:1404.0275

$$\frac{\mathcal{B}(X(3872) \rightarrow \psi(2S) \gamma)}{\mathcal{B}(X(3872) \rightarrow J/\psi \gamma)} = 3.4 \pm 1.4$$

“...The measured value agrees with expectations for a pure charmonium interpretation of the X(3872) state and a mixture of charmonium and molecular interpretations. However, it does not support a pure DD* molecular interpretation of the X(3872) state ...”

yet I would continue, citing Eichten, Lane & Quigg hep-ph/0511179

“ ...but the mass of X (3872) is too low to be gracefully identified with the 2^3P_1 charmonium state, especially if Z(3931) is to be identified as the 2^3P_2 level...”

btw charmonium violating isospin?...

Imperfect Matchings

The matchings shown above are valid only with *neutral* components though. We know that the *two* following *thresholds do not match*

$$D^\pm \bar{D}^{*0} \text{ (or } D^0 \bar{D}^{*\pm}) \neq J/\psi \rho^\pm$$

being that

$$m_{D^\pm} - m_{D^0} \approx 4.8 \pm 0.1 \text{ MeV}$$

$$m_{D^{*\pm}} - m_{D^{*0}} \approx 3.3 \pm 0.3 \text{ MeV}$$

Also the $D^+ D^{*-}$ threshold is pretty far away

$$m_{D^+ D^{*-}} - m_{D^0 \bar{D}^{*0}} \approx 8 \text{ MeV}$$

According to this last fact, we someone concluded e.g. that the dd^* component in the X has to be suppressed so that the X *cannot be a pure $l=0$ state* and this might be the seed of observed isospin violations. *On the other hand the Z_c should be a $pure\ l=1$ state and should not violate isospin as in $Z_c \rightarrow J/\psi \eta$ (not forbidden in 4q models).*

Quantum Numbers

Let's refer to $Z_c^0(3900)$. It is a 1^{+-} particle. A simple way of building it is the following. Take two colored *positive parity* bosons (*diquarks*) in S-wave ($L=0$).

$$|s_{Qq}, s_{\bar{Q}\bar{q}}, S\rangle = |1_{cq}, 1_{\bar{c}\bar{q}}, 1\rangle$$

Indeed exchanging coordinates, spins and charges, in this two boson system we can make $C=-1$

$$(-1)^L (-1)^{2s+S} C = +1 \Rightarrow C = -1$$

Another $C=-1$ state can be formed by taking simply

$$|1_{cq}, 0_{\bar{c}\bar{q}}, 1\rangle - |0_{cq}, 1_{\bar{c}\bar{q}}, 1\rangle$$


Thus we got two $J^{PC}=1^{+-}$ states, and automatically one $J^{PC}=1^{++}$ state

$$|1_{cq}, 0_{\bar{c}\bar{q}}, 1\rangle + |0_{cq}, 1_{\bar{c}\bar{q}}, 1\rangle$$


The latter could simply be the $X(3872)$ the first one; discovered back in 2003

Eightfold Way


50 years of quark model





diagonalize simultaneously T_8^3, T_8^8




diagonalize simultaneously T_{10}^3, T_{10}^8




Three quarks in the fundamental. The list of *elementary* particles gets much shorter...




Charmonium Predictions




Charmonium Levels



Bottomonium Predictions



Bottomonium Levels



Diquarks

$$[cq]_i = \epsilon_{ijk} \bar{c}_c^j \gamma_5 q^k \quad \text{spin 0}$$

$$[cq]_i = \epsilon_{ijk} \bar{c}_c^j \gamma q^k \quad \text{spin 1}$$

$|s, \bar{s}\rangle_J$ notation

$$|0, 1\rangle_1 := \frac{1}{2} \sigma^2 \otimes \sigma^2 \sigma^i$$

$$|1, 0\rangle_1 := \frac{1}{2} \sigma^2 \sigma^i \otimes \sigma^2$$


$$|1, 1\rangle_0 := \frac{1}{2\sqrt{3}} \sigma^2 \sigma^i \otimes \sigma^2 \sigma^i$$

$$|1, 1\rangle_1 := \frac{i}{2\sqrt{2}} \epsilon^{ijk} \sigma^2 \sigma^j \otimes \sigma^2 \sigma^k$$

...


Before $Z_c(4025)$ was found

Maiani, Piccinini, ADP, Riquer, PRD 2013 – type I model



*Fit with two resonances
with no further constraints.
The second structure is
a fluctuation $P=12\%$*

$$\chi^2/\text{dof} = 41/65$$



*Fit with two resonances
with $M(Z') > M(Z)$.
The second structure is
a fluctuation $P=53\%$*

$$\chi^2/\text{dof} = 47/65$$

Spin problem in type I

In the type I diquark model we have two 1^{+-} states the heavier, Z , at about 3880 MeV

$$|Z^{(\prime)}\rangle = \alpha^{(\prime)}(|10\rangle_u - |01\rangle_u) + \beta^{(\prime)}|11\rangle_u + (u \rightarrow d)$$

The expected spin of the cc^* pair being computed as

$$\ell_{c\bar{c}} = (3/2 + 2\langle Z^{(\prime)} | \mathbf{S}_c \cdot \mathbf{S}_{\bar{c}} | Z^{(\prime)} \rangle)^{1/2}$$

equal to $\sqrt{2}$ if $S_{cc^*}=1$, and 0 if $S_{cc^*}=0$. Contrary to the experimental fact that the Z is observed to decay predominantly in J/ψ , we found

$$\underbrace{\ell_{c\bar{c}}(Z')}_{\text{lighter}} \approx 3\ell_{c\bar{c}}(Z)$$

This problem is solved in the type II model in which the $S_q.S_{q^*}$ interaction is not the dominating one.

Quark-antiquark spin basis

$$|s_{q\bar{q}}, s_{c\bar{c}}\rangle_J$$

$$X_0 = \frac{1}{2}|0_{q\bar{q}}, 0_{c\bar{c}}\rangle_0 + \frac{\sqrt{3}}{2}|1_{q\bar{q}}, 1_{c\bar{c}}\rangle_0,$$

$$X'_0 = \frac{\sqrt{3}}{2}|0_{q\bar{q}}, 0_{c\bar{c}}\rangle_0 - \frac{1}{2}|1_{q\bar{q}}, 1_{c\bar{c}}\rangle_0$$

$$X_1 = |1_{q\bar{q}}, 1_{c\bar{c}}\rangle_1 \quad \text{decays into } J/\psi$$

$$Z = \frac{1}{\sqrt{2}} (|1_{q\bar{q}}, 0_{c\bar{c}}\rangle_1 - |0_{q\bar{q}}, 1_{c\bar{c}}\rangle_1)$$

$$Z' = \frac{1}{\sqrt{2}} (|1_{q\bar{q}}, 0_{c\bar{c}}\rangle_1 + |0_{q\bar{q}}, 1_{c\bar{c}}\rangle_1)$$



$$X_2 = |1_{q\bar{q}}, 1_{c\bar{c}}\rangle_2$$

Seince in type II the mass eigenvectors correspond to those listed before, we can 'Fierz' them in the quark-antiquark spin basis, and realize that both Z and Z' have also a **spin 0** component in agreement with the observed



$$Z(4020) \rightarrow \pi + h_c(1^1P_1), \text{ seen}$$

$$Z(3900) \rightarrow \pi + h_c(1^1P_1), \text{ seen (?)}$$

The 1π Correction



red: 1 pion
blue: 0 pions



X at LHC (CMS data)

