Comparison of MC and TP Dosimetry Simulations for Cancer Treatment



Raymond Mumme May 12, 2014 Supervisor: Thiago Lima (PhD student) Head Supervisor: Dr. Manjit Dosanjh



European Collaboration
CERN, CNAO (Italy), Med-Austron (Austria), HIT (Germany)

Types of Therapy

Internal:
 brachytherapy,
 injection

•External: X-rays, Protons, Light ions



Imaging

•Gamma camera •SPECT •CT •PET





Project Goals

Current Goals

- Treatment plan simulationsMonte Carlo simulations
- •FLUKA/flair
- External radiation

Simple case: 131-idodine treatment of thyroid cancer
Kinetics
Coding and calculating dose distributions (2D -> 3D ->Monte Carlo)
Voxels
Patient data



Applications	Places Syst	:em 👹 🥸 👱	<u> </u>				¢,		Mon May 12	2, 11:50 AM	owner
Į					CNAOup	d.flair - flair					_ • X
🖌 🛶 🍖 📔 🌺 Fla	air 🛛 🗿 Input	📑 Geometry 💡	🚴 Run 🛛 🔝 Plot							Calcula	tor 🔻 👔
A Cut aste Copy Clipboard	Show▼ ↓ Comment▼ ▼ CRefresh Edit	∲Move Up ♦Move Down Add▼	# Preprocessor	▼ X Delete ¥a Se Clone	11¥ arch Filter▼	Input					
Input General Primary Geometry Media Transport Biasing Goring Flair	COMPOUND STERNHE: MAT-PROI MATERIAL MATERIAL COMPOUND MATERIAL MATER	v f1: 0.755329 f3: 0.012827 I Mat: AIR ▼ P onization: 85.7 L Z: D f1: 0.625018 f3: 0.333021 L Z: D f1: 0.691148 f3: 0.073252 f5: L Z: 7.0 L Z: 29.0 L Z: 29.0 L Z: 9.0 L Z: 9.0 L Z: 0.240183 f3: 0.240183 f3: 0.8 f3: 0.8 f5:		n — M M Cba Cba Nom A Nam M M M M M M M M M M M M M M M M M M M		Gas		gen, 296K ▼ er, 296K ▼	$\begin{array}{c} \text{Lience} & 13 \\ \text{M2: OXYGEN} \\ \hline \text{M2: OXYGEN} \\ \hline \text{M2: OXYGEN} \\ \hline \text{M2: OXYGEN} \\ \hline \text{M2: AYDROGEN} \\ \hline \text{M2: HYDROGEN} \\ \hline \text{M2: AYDROGEN} \\ \hline \text{M2: AYDROGEN} \\ \hline \text{M2: AYDROGEN} \\ \hline \text{M2: CONSENSE } \\ \hline \text{M2: CARBON } \\ \hline \text{M2: CARBON } \\ \hline \text{M4: TITAHIUM} \\ \hline \\ \hline \text{M6: CONSENSE } \\ \hline \end{array}$		
	TO FIT EXP for the da look at th MAT-PROI I. HAT-PROI STOP	DATA - BRAGG RULE ta with BP somethin e data with BEAM2 w ed!! p onization: 63.5 .+2+3.	~ 67 e¥ g like 63 e¥ was t e need a bit more	the best fit in range 63.5 h Typ Ma 5+6+	e: V t: RW3 	Gas V	pressure: to Mat: "		RHOR: Step: A		

Project Goals

Current Goals

- Treatment plan simulationsMonte Carlo simulationsFLUKA/flair
- External radiation

Simple case: 131-idodine treatment of thyroid cancer
Kinetics
Coding and calculating dose distributions (2D -> 3D ->Monte Carlo)
Voxels
Patient data

Dosimetry Calculations

- MIRD Method
- •Cumulated activity over time in the source organ
- •Energy emitted by source organ
- •Fraction of energy absorbed by target organ
- •3D and Monte Carlo
- •Video

