




Introduction to Experiments or How to detect the Higgs boson

Cosmic rays are used to study the performance of the detector. Free of charge! 😊



Hess received the Nobel Prize in Physics in 1936 for his discovery (1912)



2013 NOBEL PRIZE IN PHYSICS

François Englert Peter W. Higgs



© The Nobel Foundation. Photo: Lovisa Engblom.




8 October 2013

The Royal Swedish Academy of Sciences has decided to award the Nobel Prize in Physics for 2013 to

François Englert and Peter Higgs

“for the theoretical discovery of a mechanism that contributes to our understanding of the origin of mass of subatomic particles, and which recently was confirmed through the discovery of the predicted fundamental particle, by the ATLAS and CMS experiments at CERN’s Large Hadron Collider”



Outline

- Introduction
- CERN and the Large Hadron Collider (LHC)
 - The accelerator
 - The detectors
- The Higgs discovery
 - Spin and parity
 - Recent results on Higgs decaying to fermions




The Standard Model

- Is a very successful theory and describes the world around us.
- The Standard Model is a discovery in itself
- However, it explains only a fraction of the universe (~5%)
 - 95% is dark energy and dark matter. What is made of? The search is ongoing...
 - What about super symmetry (SUSY)


The Standard Model (1970-90s)

- Matter particles: fermions (1/2 integer spin)
- ‘Force’ particles: bosons (integer spin)
- Higgs field causes electro weak symmetry breaking and gives particles their masses



→ Nucleon level (partons) : binding energy ~98% of the mass


→ Most of the (luminous) mass in the universe comes from QCD confinement energy




A bit of history

1964 The BEH mechanism

The Standard Model completed



LHC approved on
16 December 1994



In 1976:


A PHENOMENOLOGICAL PROFILE OF THE HIGGS BOSON

John Ellis, Mary K. Gaillard ^{*)} and D.V. Nanopoulos ⁺⁾
 CERN -- Geneva

The Roadmap:


We should perhaps finish with an apology and a caution. We apologize to experimentalists for having no idea what is the mass of the Higgs boson, unlike the case with charm ^{3),4)} and for not being sure of its couplings to other particles, except that they are probably all very small. For these reasons we do not want to encourage big experimental searches for the Higgs boson, but we do feel that people performing experiments vulnerable to the Higgs boson should know how it may turn up.

How ?




$E=3.5\text{TeV} \rightarrow$
 $V=99.999996\%$ of c

$$m = \frac{m_0}{\sqrt{1 - \frac{v^2}{c^2}}}$$



Energy = Matter
 $E^2 = (m_0 c^2)^2 + (pc)^2$



LHC (2012)


2x4 TeV How much energy?

- 1.9×10^{14} protons per beam at 4 TeV
- How much energy in the two beams

8 Lamborghini



speed=350 km/h

8 Lamborghini



Experimental High Energy Physics – detecting particles

Two Protons collide at high energy
Large Hadron Collider (LHC) at CERN



Center-of-Mass Energy (Nominal)
14 TeV ?

Center-of-Mass Energy (close to nominal)
13 TeV

Restart in 2015

LHCb

ATLAS

Center-of-Mass Energy (2012)


8 TeV

CMS


ALICE

Center-of-Mass Energy (2010-2011)

7 TeV




Large Hadron Collider (LHC)



• The Accelerator


- 100 - 150 m below surface at 1.9 Kelvin in a tunnel 27 km long.
- The protons circulate at a speed of ~ 11000 turns/sec
- There are 2808 bunches
- Collisions at 40 MHz (every 25 ns)
- **600 000 000 collisions per second !**




The experiments


*CMS: heavier than
the Eiffel Tower*

*ATLAS: as big as a
5 storey building*



Största och mest sofistikerade detektorer






Principles of Detection




The collision energy condenses into particles (e , p , π , μ , γ , K ...)

Detectors surrounding the collision point (or *after* in case of fixed target) are sensitive to the passage of energetic particles.



Partikeldetektorer



Interaction of Particles with Matter

In order to detect a particle it must interact with matter!

The most important interaction processes are electromagnetic:

Charged Particles:

- Energy loss due to ionization (e.g. charged track in straw detector) heavy particles (*not electrons/positrons!*)
- Energy loss due to photon emission (electrons, positrons) bremsstrahlung

Photons:

Interaction of photons with matter (e.g. EM calorimetry)

Photoelectric effect

Compton effect

Pair production

Other important electromagnetic processes:

Multiple Scattering (Coulomb scattering)

scintillation light (e.g. TOF systems)

Cherenkov radiation

Transition Radiation (e.g. particle id normally electrons)

*Can calculate the above effects with a combo of classical E&M and QED.
In most cases calculate approximate results, exact calculations very difficult.*

Bethe-Bloch Formula for Energy Loss

Average energy loss for **heavy charged particles**

Energy loss due to ionization and excitation

Valid for energies < 100 's GeV and $\beta \gg z\alpha$ ($\approx z/137$)

heavy = $m_{\text{incident}} \gg m_e$
proton, k , π , μ

$$-\frac{dE}{dx} = 2\pi N_a r_e^2 m_e c^2 \rho \frac{Z}{A} \frac{z^2}{\beta^2} \left[\ln\left(\frac{2m_e \gamma^2 v^2 W_{\text{max}}}{I^2}\right) - 2\beta^2 \right]$$

Fundamental constants

r_e = classical radius of electron

m_e = mass of electron

N_a = Avogadro's number

c = speed of light

$$= 0.1535 \text{ MeV-cm}^2/\text{g}$$

Absorber medium

I = mean ionization potential

Z = atomic number of absorber

A = atomic weight of absorber

ρ = density of absorber

δ = density correction

C = shell correction

Incident particle

z = charge of incident particle

β = v/c of incident particle

$$\gamma = (1 - \beta^2)^{-1/2}$$

W_{max} = max. energy transfer
in one collision

$$W_{\text{max}} = \frac{2m_e (c\beta\gamma)^2}{1 + m_e/M + \sqrt{1 + (\beta\gamma)^2 + (m_e/M)^2}} \approx 2m_e (c\beta\gamma)^2$$

Note: the classical dE/dx formula contains many of the same features as the QM version: $(z/\beta)^2$, & $\ln[\]$


$$-dE/dx = \frac{4\pi z^2 r_e^2 m_e c^2 N_e}{\beta^2} \ln \frac{b_{\text{max}}}{b_{\text{min}}}$$

Bethe-Bloch Energy Loss

$$-\frac{dE}{dx} = 2\rho N_a r_e^2 m_e c^2 r \frac{Z}{A} \frac{z^2}{b^2} \frac{1}{\beta^2} \ln\left(\frac{2m_e g^2 v^2 W_{\max}}{I^2}\right) - 2b^2 \frac{1}{\beta^2}$$

PDG plots:

<http://pdg.lbl.gov/index.html>




Calculated

$$g = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$


Bremsstrahlung (braking radiation)

- A fast moving particle is decelerated in the electrical field of the nuclei.
- Above a few tens MeV, bremsstrahlung is the most dominated process for **electrons and positrons**
- It becomes important to muons (and pions) at a few hundred GeV
- What about the atomic electrons? Yes, the electron cloud gives an *additional contribution* to the bremsstrahlung
- *Let's see how this is used in the detector layout later*





Photons - 3 interactions

Compton scattering




Photoelectric Effect



Photons interacting with matter

- Photoelectric effect
- Compton scattering
- Pair production
- *Mass Attenuation Coefficient* = Interaction probability/density



Multiple Scattering

A charged particle traversing a medium is deflected by many small angle scatterings. These scatterings are due to the **coulomb field of atoms** and are **assumed to be elastic**. In each scattering the energy of **the particle is constant but the particle direction changes**.


In the simplest model of multiple scattering we ignore large angle scatters.

In this approximation, the distribution of scattering angle θ_{plane} after traveling a distance x through a material with radiation length $=L_r$ is approximately gaussian:

$$\frac{dP(\theta_{plane})}{d\theta_{plane}} = \frac{1}{\theta_0 \sqrt{2\pi}} \exp\left[-\frac{\theta_{plane}^2}{2\theta_0^2}\right] \quad \text{with} \quad \theta_0 = \frac{13.6 \text{ MeV}}{\beta p c} z \sqrt{x/L_r} (1 + 0.038 \ln\{x/L_r\})$$


In the above equation $\beta=v/c$, and p =momentum of incident particle

This is not good for tracking!



Cherenkov light

- Named after the Russian scientist P. Cherenkov who was the first to study the effect in depth (he won the Nobel Prize for it in 1958)
- From Relativity, nothing can go faster than the speed of light c (in vacuum)
- However, due to the refractive index n of a material, a particle *can* go faster than the *local* speed of light in the medium $c_p = c/n$
- Fast electrons in a reactor emitting blue light (Cherenkov radiation)
- This is analogous to the bow wave of a boat travelling over water or the sonic boom of an aeroplane travelling faster than the speed of sound



Cherenkov radiation

The left corner of the triangle represents the location of the superluminal particle at some initial moment ($t=0$). The right corner of the triangle is the location of the particle at some later time t . In the given time t , the particle travels the distance


$$x_p = v_p t = \beta ct$$

whereas the emitted electromagnetic waves are constricted to travel the distance

$$x_{em} = v_{em} t = \frac{c}{n} t$$


So:

$$\cos \theta = \frac{1}{n\beta} = \frac{v_1}{v_p}$$




Cherenkov Detector

Mean Cherenkov Angle per track



LHCb



AMS



Transition Radiation (Particle ID)

- Transition Radiation: photon emitted by a charged particle when traversing the boundary between materials with different dielectrical constants ($\epsilon_1 \epsilon_2$)

$$g = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$


- $\gamma > 1000$

- Intensity: $I \sim \gamma = E/m$, $\theta \sim 1/\gamma$

→ Identification of transition radiation photons used for

particle identification (mostly electrons)

of particles with momenta between 1 and few 100 GeV



Transition radiation

(particle identification)

- Number of emitted photons per boundary $N_{ph} \approx \frac{W}{\hbar\omega_p} \propto \alpha$ is very small.
- Need many transitions to produce a sizable signal.

TR Radiators:

- stacks of thin foils made out of CH_2 (polyethylene), $\text{C}_5\text{H}_4\text{O}_2$ (Mylar)
- hydrocarbon foam and fiber materials. Low Z material preferred to keep re-absorption small ($\propto Z^5$)




alternating arrangement of radiator stacks and detectors
→ minimizes re-absorption




TR X-ray detectors:

- Detector should be sensitive for $3 \leq E_\gamma \leq 30$ keV.
- Mainly used: Gas detectors: MWPC, drift chamber, straw tubes...
- Detector gas: $\sigma_{\text{photo effect}} \propto Z^5$

→ gas with high Z required, e.g. Xenon ($Z=54$)




TRT (ATLAS): 3 straws and radiators



High threshold

Low threshold


Detector Challenges (Highlights)



- **Trigger Challenge** : How to select 400 out of 20×10^6 events per second while keeping the interesting (including unknown) physics
- **Computing Challenge** : How to reconstruct, store and distribute 400 increasingly complex events per second (over 100 Petabyte per experiment)

The first LHC run


Event rate = luminosity x cross-sections



2010

O(2) Pile-up events

150 ns inter-bunch spacing



2011

O(10) Pile-up events

50 ns inter-bunch spacing




Design value (expected to be reached at L=10³⁴ !)

2012

O(20) Pile-up events

50 ns inter-bunch spacing



The detection of the Higgs boson





Higgs production

Vector boson fusion
VBF

Top-antitop fusion
ttH




Gluon-gluon fusion
 $gg \rightarrow H$

Higgs strahlung
VH




Detect Higgs by decay products


- Variety of decay channels
- Massive particles more likely
- Difficult to detect from background
- Life time is 1.56×10^{-22} s (!)
(predicted in the Standard Model)
- $\gamma\gamma$ is clean, but rare





April-July 2012: 8 TeV , 5.8 fb^{-1}




Measure energy of photons emitted




Measure decay products of Z bosons




H → 4l





Higgs events $H \rightarrow 4l$ (muons)



$$H \rightarrow \gamma\gamma$$



From CMS Higgs $\rightarrow \gamma\gamma$





Higgs to Fermions

- Recent analysis shows evidence of Higgs boson decaying to fermions (leptons or quarks). Not previously observed!
- It is important to measure this decay but no surprise is expected
- Difficult due to high background

BR related to mass. High mass fermions preferred (τ, b)


- Both ATLAS and CMS has “evidence” in the channel $H \rightarrow \tau\tau$
....but not yet the famous “ 5σ ” needed to claim discovery

Higgs decays to fermions ($\tau\tau$) in ATLAS (26 Nov 2013)




The taus decay into an electron (blue line) and a muon (red line)

Higgs decays to fermions ($\tau\tau$) in CMS (3 Dec. 2013)



One tau decays to neutrinos and a muon (red lines on the right), while the other decays into a charged hadron (blue towers) and a neutrino



Thank You!

Literature

- CERN Academic Training


<http://indico.cern.ch/conferenceDisplay.py?confId=266737>

- CERN ATLAS


<http://www.atlas.ch/HiggsResources/>

Spares

Higgs “exclusion plots”




The ATLAS experiment



LHC

20 Years, projecting, constructing and Simulating...



1. Accelerators:

Powerful machines capable of accelerating particles to very high energies before being collided with other particles

2. Detectors:


Gigantic instruments that record particle collisions

3. Computers:

To collect, store, distribute and analyze enormous quantities of data generated by detectors


The biggest scientific instrument ever built

The Higgs search as of 18/11/2011 ☺



To be continued.....

After 10 min of LHC running: full history of SM



What LEP (CERN) and Tevatron (Fermilab)

Until year 2000




Figure 2.10: Higgs exclusion range from LEP experiments.

Until year 2011





Figure 2.11: Combined Run II Higgs limits from the Tevatron experiments.




$\mu^+(Z_1) p_T : 43 \text{ GeV}$


8 TeV DATA

4-lepton Mass : 126.9 GeV

$\mu^-(Z_1) p_T : 24 \text{ GeV}$



$e^-(Z_2) p_T : 10 \text{ GeV}$




$e^+(Z_2) p_T : 21 \text{ GeV}$

CMS Experiment at LHC, CERN
Data recorded: Mon May 28 01:35:47 2012 CEST
Run/Event: 195099 / 137440354
Lumi section: 115

On example: the discovery of the quarks at SLAC in 1968

$$l = \frac{h}{p}, P = 20 \text{ GeV} \Rightarrow l \gg 10^{-17} \text{ m}$$



- The quark model was independently proposed by physicists [Murray Gell-Mann](#) and [George Zweig](#) in 1964.
- Gell-Mann found the quarks in:

“Three quarks for Muster Mark!
Sure he has not got much of a bark
And sure any he has it's all beside the mark.”

—James Joyce, *Finnegans Wake*