

Exotic Higgs Boson Decays

Jessie Shelton



U. Illinois at Urbana-Champaign



SUSY 2015
August 25, 2015

The SM-like Higgs boson


- A light SM-like Higgs is **narrow**:



$$\Gamma_h(125 \text{ GeV}) = 4.1 \text{ MeV}$$

Exotic decays of the SM-like Higgs

- Presence of new light degrees of freedom can distort Higgs Brs by $O(1)$ even for small couplings



Simple example:
one new scalar


$$\Delta\mathcal{L} = \frac{\zeta}{2} s^2 |H|^2$$

Why exotic Higgs decays?

- Motivations for new physics at the weak scale:
 - co-responsible for **generating it**
 - **stabilize it**
 - **thermal dark matter**
 - ...why not?

Why exotic Higgs decays?

- Motivations for new physics at the weak scale:
 - co-responsible for **generating it**
 - **stabilize it**
 - **thermal dark matter**
 - ...why not?
- These motivations apply **horizontally** as well as **vertically**




Why exotic Higgs decays?

- **Extended Higgs sectors:** SM + s , MSSM + S , composite models, ...
 - simplest realization of Higgs portal coupling: $|S|^2|H|^2$
 - **NMSSM:** dynamically generate μ , relax phenomenological constraints on $V(H)$, neutralino dark matter
 - **electroweak phase transition:** baryogenesis, cosmological history of the SM

Why exotic Higgs decays?

- **Naturalness**


- Twin Higgs and related models:



- light weak-scale states needed for naturalness can be **SM singlets**
- Higgs portal interactions by construction; also possibly hypercharge

Why exotic Higgs decays?

- **Dark matter:**
 - First work on exotic Higgs decay: $h \rightarrow$ dark matter
 - “WIMP miracle”: a statement about cold dark matter freezing out via perturbative interactions
 - Hidden sector freezeout:




Why exotic Higgs decays?

- **Why not?**
 - Hidden sectors are a **generic** ingredient in UV theories: e.g., SUSY-breaking
 - Generic signatures of new physics may be **light, weakly coupled states** just as well as **heavier, SM-charged states**
 - Characterize signatures by **leading operators** mediating SM-HS interactions
 - **Higgs portal**: unique possibilities at LHC: direct Higgs production, small SM width

Example: $h \rightarrow 4b$

- Generic prediction of Higgs-portal (pseudo-)scalars:

$$h \rightarrow ss(aa) \rightarrow 4b$$




- characterized by two quantities: m_s , $\text{Br}(h \rightarrow ss)$

Example: $h \rightarrow 4b$

- Generic prediction of Higgs-portal (pseudo-)scalars:

$$h \rightarrow ss(aa) \rightarrow 4b$$




mass, couplings related to
phase transitions

- Electroweak baryogenesis:
 - couplings of new (complex) scalar singlet to SM Higgs can drive EWPT to be **strongly first-order**
 - If $m_s < 2 m_h$: $h \rightarrow ss$ unavoidable, same operator gives $s \rightarrow bb$
 - Higgs properties and in particular $\text{Br}(h \rightarrow \text{SM})$ leading constraints

Example: $h \rightarrow 4b$

- Generic prediction of Higgs-portal (pseudo-)scalars:

$$h \rightarrow ss(aa) \rightarrow 4b$$



dark mediator properties
may be related to other DM
signals



- Dark matter:

- Thermal WIMP: $XX \rightarrow a \rightarrow \text{SM}$,
 $XX \rightarrow ss (aa)$
- Leading signatures: indirect
detection*, direct detection*, exotic
Higgs decays*
- Branching ratio bounded by **BBN
constraints** on mediator lifetime:
effectively free parameter (and can
be very small)

Example: $h \rightarrow 4b$

- Generic prediction of Higgs-portal (pseudo-)scalars:

$$h \rightarrow ss(aa) \rightarrow 4b$$



composite glueballs
decay via Higgs mixing


- Neutral naturalness
 - s is composite: dark glueballs
 - Leading signatures: Higgs properties, especially $\text{Br}(h \rightarrow \text{SM})$
 - Composite: decays can be parametrically separated from production \Rightarrow displacement

Lessons for collider searches

- **Signature-based approach:** typically, many models yield the same final state
 - relatively few ways to couple SM singlet new physics to the SM
 - **helps inform searches:** Yukawa ordered, gauge-ordered, ...

Lessons for collider searches

- **Signature-based approach:** typically, many models yield the same final state
 - relatively few ways to couple SM singlet new physics to the SM
 - **helps inform searches:** Yukawa ordered, gauge-ordered, ...
- Minor changes or additions to the BSM physics can lead to **$O(1)$ changes in signatures**
 - nature does not guarantee minimality!
 - important to cast a wide net



Exotic Higgs decays at the LHC

A cartoon:




Exotic Higgs decays at the LHC

A cartoon:

Several searches
in Run I

Improve capability
for especially well-
motivated modes




Exotic Higgs decays at the LHC

A cartoon:

Several searches
in Run I

Institute broad search
program for signals
with good prospects

Improve capability
for especially well-
motivated modes



motivation



reach

Exotic Higgs decays at the LHC

- The LHC as an intensity frontier machine
 - Higgs production cross-section at 8 TeV: ~ 20 pb
 - Integrated luminosity, ~ 20 / fb
 - $\rightarrow \sim 400000$ Higgs bosons served
 - If: reasonable reconstruction efficiency, good S/B : statistics for branching fractions $\sim 10^{-4}$


Exotic Higgs decays at the LHC

- Indirect limits: observation of SM modes



Higgs decays to (pseudo-)scalars


- Our example process $h \rightarrow ss(aa) \rightarrow 4b$



Four soft b -jets:


$$p_T \lesssim 30 \text{ GeV}$$

use VH associated production



Higgs decays to (pseudo-)scalars

- Current status: **mass-dependent efficiency** for an $h \rightarrow 4b$ event to pass SM $h \rightarrow 2b$ search criteria
 - For light (~ 15 GeV) scalars: $Br(h \rightarrow 4b) \lesssim 0.7$
 - Heavier scalars: **no limit**
- Future prospects:
 - analyses with, without jet substructure
 - ultimate 95% CL sensitivity in both cases *estimated* to be
$$Br(h \rightarrow 4b) \approx 0.1$$




Higgs decays to (pseudo-)scalars

- Power of clean dimuon resonance: $h \rightarrow ss(aa) \rightarrow 2b2\mu$

$$\frac{Br(a \rightarrow 2\mu)}{Br(a \rightarrow 2b)} \sim \frac{m_\mu^2}{3m_b^2} \approx 2 \times 10^{-4}$$

resonant dimuon pair plus:




mass-drop + b -tag

$2j$ (m_{jj} , $m_{jj\mu\mu}$ consistent with resonant origin)


$2b$ (m_{bb} , $m_{bb\mu\mu}$ consistent with resonant origin)

Extending reach

- Extend sensitivity with other final states: **new trigger strategies**



X




2

- These are (mostly) **not easy searches**
 - active theory-experiment interaction important!

Higgs decays to (pseudo-)scalars


- **Displaced** decays are in some ways easier: S/B



- but triggering and reconstruction are highly nontrivial

Non-minimal dark sectors

- BSM states will often prefer to decay to **other dark states**, if such decays are available
- A weakly-coupled example: **Higgsed dark $U(1)$**




- Higgs mixing: $h \rightarrow ss$, but
now $s \rightarrow Z_D Z_D$,
 $h \rightarrow Z_D Z_D$

$$\Delta\mathcal{L} = V(S) + \frac{\kappa}{4} S^2 |H|^2 + \epsilon B_{\mu\nu} V^{\mu\nu}$$

Lessons from LHC Run I


- Leptophilic decays make for a much easier signal: statistics-limited




Easy: multiple resonant light leptons

$$\text{Br} \lesssim 4 \times 10^{-4}$$

Moderate: multiple electroweak objects, poor mass resolution



$$\text{Br} \lesssim 0.1 - 0.6$$




Hard: all-hadronic

$$\text{Br} \lesssim 0.9$$



Beyond the LHC

Other colliders, other possibilities



Fixed target


$e-p$: LHeC,
FCC-he



lepton colliders

A Higgs window

- Higgs portal coupling is a powerful window into dark sectors



Summary and conclusions

- The observed 125 GeV Higgs boson is **highly sensitive** to the potential existence of **new light degrees of freedom**
- **Higgs portal couplings + BSM at weak scale** are vital ingredients of many theories of cosmology, naturalness
- **LHC as a Higgs factory**: interesting results and prospects for many exotic decay modes
 - **Triggers** will be important for maintaining sensitivity
- Unprecedented opportunities to explore Higgs portal!