



Contribution ID: 58

Type: Poster

Structure of ^{68}Ni : new insights on the low-lying 0^+ and 2^+ states from two-neutron transfer on ^{66}Ni and beta-decay of ^{68}Co

The region around the nucleus ^{68}Ni , with a shell closure at $Z = 28$ and a sub-shell closure at $N = 40$, is the source of considerable interest in nuclear-structure studies. Despite a significant set of experimental and theoretical information available on ^{68}Ni [1-5], the origin of its structure is still being questioned. A recent clarification of the energy and spin assignment of several low-lying 0^+ and 2^+ states [6-9] and state-of-the-art shell model calculations [5,10] hinted to the possibility of triple shape coexistence and highlighted the need of additional experimental investigation.

To better understand the structure of ^{68}Ni , two complementary experiments: the two-neutron transfer reactions on ^{66}Ni at 2.85 MeV/u and the beta-decay of ^{68}Co were performed at ISOLDE.

On one hand, the $^{66}\text{Ni}(t,p)^{68}\text{Ni}$ reaction represents a unique tool to probe the nature of 0^+ states in ^{68}Ni . Coincidences between the outgoing light charged particles and gamma-rays were detected using the combined MINIBALL [11] gamma-ray spectrometer and the T-REX particle detection array [12]. Results of such coincidence analysis together with the reconstruction of angular distributions of the reaction products, revealing the most populated states, will be presented. An interpretation based on calculations within the Distorted-Wave Born Approximation (DWBA) and shell model two-nucleon amplitudes will be discussed.

On the other hand, the measurement of the beta-decay of the low spin isomer in ^{68}Co selectively produced in the decay chain of ^{68}Mn allowed us to build a revised decay scheme to ^{68}Ni based on the clear identification of beta-gamma- $E0$ delayed coincidences. A strong emphasis will be put on the connections between the three lowest lying 0^+ and 2^+ determined from observed transitions and upper limits.

- [1] M. Bernas et al., PLB 113, 279 (1982)

- [2] R. Broda et al., PRL 74, 868 (1995)

- [3] W. Mueller et al., PRC 61, 054308 (2000)

- [4] O. Sorlin et al., PRL 88, 092501 (2002)

- [5] S. Lenzi et al., PRC 82, 054301 (2010)

- [6] S. Suchyta et al., PRC 89, 021301(R) (2014)

- [7] F. Recchia et al., PRC 88, 041302 (2013)

- [8] C.J. Chiara et al., PRC 86, 041304 (2012)

- [9] R. Broda et al., PRC 86, 064312 (2012)

- [10] D. Tsunoda et al., PRC 89, 031301(R) (2014)

- [11] N. Warr et al., EPJA 49, 40 (2013)

- [12] V. Bildstein et al., EPJA 48, 85 (2012)

Primary author: Mr RAABE, Riccardo for the IS467 and IS504 collaborations (KU Leuven, Instituut voor Kern- en Stralingsfysica)

Presenter: Mr RAABE, Riccardo for the IS467 and IS504 collaborations (KU Leuven, Instituut voor Kern- en Stralingsfysica)