



Contribution ID: 287

Type: **Talk**

## Fast luminous blue transients from newborn black holes

*Tuesday 15 December 2015 15:03 (21 minutes)*

Newborn black holes in collapsing massive stars can be accompanied by a fallback disc. The accretion rate is typically super-Eddington and strong disc outflows are expected. Such outflows could be directly observed in some failed explosions of compact (blue supergiants or Wolf-Rayet stars) progenitors, and may be more common than long-duration gamma-ray bursts. Using an analytical model, we show that the fallback disc outflows produce blue UV-optical transients with a peak bolometric luminosity of  $\sim 10^{42-43}$  erg s $^{-1}$  (peak R-band absolute AB magnitudes of -16 to -18) and an emission duration of  $\sim$  a few to  $\sim 10$  d. The spectra are likely dominated intermediate mass elements, but will lack much radioactive nuclei and iron-group elements. The above properties are broadly consistent with some of the rapid blue transients detected by Panoramic Survey Telescope & Rapid Response System and Palomar Transient Factory. This scenario can be distinguished from alternative models using radio observations within a few years after the optical peak.

**Primary author:** Dr KASHIYAMA, Kazumi (University of California, Berkeley)

**Presenter:** Dr KASHIYAMA, Kazumi (University of California, Berkeley)

**Session Classification:** 13 - Gamma-ray bursts