

# A Second-Quantization Approach to the Analytical Faraday Effect in Graphene

Thursday 21 May 2015 08:00 (3 hours)

We investigate coherent single photon traveling in the positive  $z$  direction passes through a graphene film subjected to a transverse magnetic field  $B$ . Analytical expressions are obtained for the transmission and reflection coefficients by full second-quantized form. Giant Faraday rotations in the infrared regime are generated and measurable Faraday rotation angles in the visible range become possible. For an example, this explains the departure from the semiclassical value for  $\theta_F$  observed in the right panel in  $B \approx 1$  T for  $\hbar\omega = 10$  meV ( $B \approx 0.5$  T for  $\hbar\omega = 30$  meV). If the magnetic field intensity is higher than a given value, we necessarily have  $N_F = 0$  (for 0.05 eV this value is about 1.9 T). In this case, the Hall conductivity at  $T = 0$  is fully determined by a single type of interband transition, and assuming  $E_1(B) \gg \hbar\omega$ , we obtain  $\theta_F \approx 3 \times 10^{-4}$ .

**Author:** Mr NUALPIJIT, Phusit (Department of Physics, Faculty of Science, Kasetsart University, Bangkok, THAILAND 10900)

**Co-author:** Mr BOONCHUI, Sooty (Department of Physics, Faculty of Science, Kasetsart University, Bangkok, THAILAND 10900)

**Presenter:** Mr NUALPIJIT, Phusit (Department of Physics, Faculty of Science, Kasetsart University, Bangkok, THAILAND 10900)

**Session Classification:** Poster-2

**Track Classification:** Condensed Matter Physics