



Search for heavy resonances  
decaying into a pair of boosted bosons  
in the  $\tau^- \tau^+ q \bar{q}$  final state at CMS

**Camilla Galloni**  
**University of Zurich**

**September 11th 2014**

# Theory Motivation

The **Standard Model** is a successful theory describing the fundamental interactions of matter constituents

However some issues still need to be addressed: hierarchy problem, unification of interactions, etc...

Many **New Physics** (NP) Scenarios have been proposed:

new symmetries and properties -> **new particles** have been introduced

- Standard Model extensions
- Extra Dimensions Model
- Beyond the Standard Model searches of new resonances:
  - Low mass searches (up to 1 TeV)
  - High mass searches (1 TeV - 2.5 TeV) -> **Boosted regime**


**First searches for high mass narrow resonances in the  $\tau\tau^+$  qq final state**



# ZH Search: Benchmark model

- New Physics searches adopting a Simplified Model strategy
- Heavy Vector Triplets model predicts 3 massive gauge bosons:  $Z'$  and  $W'^{\pm}$
- **Phenomenological Lagrangian** with only the relevant couplings and the mass parameters

arXiv:1402.4431




$$\begin{aligned}
 \mathcal{L}_V = & -\frac{1}{4}D_{[\mu}V_{\nu]}^a D^{[\mu}V^{\nu]}_a + \frac{m_V^2}{2}V_\mu^a V^{\mu a} \\
 & + i g_V c_H V_\mu^a H^\dagger \tau^a \overleftrightarrow{D}^\mu H + \frac{g^2}{g_V} c_F V_\mu^a J_F^{\mu a} \\
 & + \frac{g_V}{2} c_{VVV} \epsilon_{abc} V_\mu^a V_\nu^b D^{[\mu}V^{\nu]}_c + g_V^2 c_{VVHH} V_\mu^a V^{\mu a} H^\dagger H - \frac{g}{2} c_{VW} \epsilon_{abc} W^{\mu\nu a} V_\mu^b V_\nu^c
 \end{aligned}$$

- $g_V$  gauge coupling of the new interaction
- $c_H, c_L, c_q, c_3$ , couplings to the H, leptons, light and third family quarks
- $m_V$  resonance mass parameter



# ZH Signal topology




Final state with:


- two tau leptons from the Higgs decay
- two quarks from the Z boson decay

Heavy vector boson search starting from  $m_{Z'} > 1 \text{ TeV}$ :

- **H and Z very boosted**
- the final products can be really collimated
- H boson: boosted tau reconstruction requires **modification in the lepton ID**
- Z boson: reconstruction through **substructure study**



# ZH Signal topology




Final state with:

- two tau leptons from the Higgs decay
- two quarks from the Z boson decay



Six possible final states, depending on the  $\tau$  decay:

- Fully leptonic channel:
  - $\tau\tau \rightarrow \mu\mu$  4v,  $\tau\tau \rightarrow \mu e$  4v,  $\tau\tau \rightarrow ee$  4v
- Semi-leptonic channel:
  - $\tau\tau \rightarrow nh \mu$  3v,  $\tau\tau \rightarrow nh e$  3v
- All hadronic channel:
  - $\tau\tau \rightarrow 2 nh$  2v



# Object Identification

- Particle Flow Algorithm — all the information from the sub detectors are combined to reconstruct all particles in the collisions :
  - charged and neutral hadrons, photons, electrons, muons
- Particles are used to identify jets, tau and MET



**Missing Transverse Energy (MET)** for the presence of escaping neutrinos:

- Computed as the negative sum of all the reconstructed particles momenta in the event
- Various Corrections: Jet energy correction and X-Y corrections


# Object Identification: Z-jet

The **Z boson** is reconstructed as one single energetic massive fat jet ( $R = 0.8$ ) for the merging of the two jets coming from the quarks hadronization  $\rightarrow p_T > 400$  GeV to match the trigger requirements

**Pruning technique:** removes the soft and large angle emitted radiation inside the jet ( $70 \text{ GeV} < m(\text{pruned jet}) < 110 \text{ GeV}$ )

**Jet substructure:** jet track topology is analyzed to see the number of subjets



N-subjettiness  $\tau_{21} < 0.75$



# Object Identification: taus

**Hadronic taus:** starting from a PF jet



- Cut based and MultiVariate identification discriminants :
  - Decay mode finding: tau lepton decay signature
  - Isolation: used to reject QCD jets
  - Muon rejection
  - Electron rejection





# Boosted regime

- Higgs decay products very collimated
- Tracks coming from a tau decay may enter the other tau cone and be considered by PF as an additional decay product
- Challenge in the identification of the particles:
  - hadronic tau reconstruction
  - lepton isolation




# Hadronic tau reconstruction

Jet cleaning procedure:  $\longrightarrow$  PF Tau Producer


- electrons and muons identified with loose requirements inside the jet cone
- removed from the Jet constituents

CMS-EXO-13-007

CMS Work in Progress



CMS Work in Progress






$$\epsilon = \frac{DEN \ \& \ PassAllDMs}{\Delta R(\tau_{reco}, \tau_{gen}) < 0.4 \ \& \ pT(\tau_{reco}, gen) > 20}$$



# Lepton Isolation

## New algorithm for lepton isolation:

- hadronic tau in the lepton isolation cone
- tau is fully identified (IDs)
- the PF constituents removed from the lepton isolation deposits



# H $\rightarrow$ $\tau\tau$ reconstruction



Neutrinos in the final state

Challenge in kinematics reconstruction

**SVFit** tool: algorithm for taus invariant system reconstruction

arXiv:1401.5041

- Probability model:
  - measured METx and METy
  - the tau decay visible kinematics
  - the MET resolution
- Marginalized likelihood function on an event-by-event basis



Higgs mass reconstructed for all the resonance masses



# Event Selection

## Fully/Semi leptonic channel:

$$p_T(\tau_1 + \tau_2)_{\text{SVFit}} > 100 \text{ GeV}$$

## Fully hadronic channel:

$$\Delta\phi(\tau, \text{MET}) < 1.5$$

$$\Delta\phi(\text{Z-jet}, \text{MET}) > 2$$

## Z-jet side:

$$p_T > 400 \text{ GeV}$$

$$M_{\text{pruned}} \in [70, 110] \text{ GeV}$$

$$\text{N-subjettiness } \tau_{21} < 0.75$$

## Additional requirements:

**b-jet Veto:** b-jets tagged using CSV loose WP, jet is required to have  $\Delta R(\text{jet}, \text{lep}) > 0.5$  and  $\Delta R(\text{jet}, \text{Z-jet}) > 0.5$




Selection	$ee, \mu\mu, e\mu$ channels	$\mu\tau_{\text{had}}, e\tau_{\text{had}}$ channels	$\tau_{\text{had}}\tau_{\text{had}}$ channel
$E_T^{\text{miss}}$	$> 100 \text{ GeV}$	$> 50 \text{ GeV}$	$> 80 \text{ GeV}$
$p_{T,l}^{\text{leading}}$	-	$> 35 \text{ GeV}$	$> 50 \text{ GeV}$
$N_{\text{b-tagged jet}}$	$= 0$	$= 0$	-
$\Delta R_{\ell\ell}$	$< 1.0$	$< 1.0$	$< 1.0$
$m_{\tau\tau}$	-	-	105-180 GeV



# Background Estimation

Background estimation strategy varies accordingly to the tau decay channels


- Different background composition: Drell Yan, top pair production,  $W$ + jets..
- Data driven estimation procedure



# Results

No deviation from the background prediction is observed in data

Upper limits on the production cross section of a new resonance decaying into ZH bosons is set



	Mass	$\tau_e \tau_e$	$\tau_\mu \tau_\mu$	$\tau_e \tau_\mu$	$\tau_e \tau_{had}$	$\tau_\mu \tau_{had}$	$\tau_{had} \tau_{had}$
$\epsilon_{sig}(\%)$	1.0 TeV	$17 \pm 2$	$38 \pm 2$	$24 \pm 1$	$21.2 \pm 0.6$	$29.3 \pm 0.7$	$18.2 \pm 0.5$
	1.5 TeV	$30 \pm 2$	$53 \pm 2$	$42 \pm 2$	$29.2 \pm 0.8$	$38.1 \pm 0.9$	$29.0 \pm 0.7$
	2.0 TeV	$28 \pm 2$	$56 \pm 3$	$39 \pm 2$	$31.1 \pm 0.8$	$39.2 \pm 0.9$	$31.9 \pm 0.7$
	2.5 TeV	$27 \pm 2$	$42 \pm 2$	$37 \pm 2$	$26.8 \pm 0.8$	$37.0 \pm 0.8$	$30.0 \pm 0.7$
$N_{bkg}$	1.0 TeV	$1.2 \pm 1.2$	$2.0 \pm 0.9$	$1.7 \pm 1.0$	$9.5 \pm 3.5$	$7.6 \pm 2.2$	
	1.5 TeV	$0.4 \pm 0.4$	$0.9 \pm 0.4$	$0.07 \pm 0.04$	$4.3 \pm 1.8$	$2.6 \pm 0.9$	
	2.0 TeV	$< 0.5$ at 68% CL	$0.7 \pm 0.4$	$< 0.4$ at 68% CL	$0.1 \pm 0.1$	$< 0.4$ at 68% CL	$6.1^{+3.2}_{-2.5}$
	2.5 TeV	$< 2.1$ at 68% CL	$0.3 \pm 0.1$	$< 0.3$ at 68% CL	$0.18 \pm 0.05$	$< 0.5$ at 68% CL	
$N_{observed}$	1.0 TeV	2	5	2	2	13	
	1.5 TeV	0	1	0	2	5	8
	2.0 TeV	0	0	1	0	0	
	2.5 TeV	0	0	0	0	0	

Resonance production cross section in a range between 0.7 and 4.1 fb are excluded at 95% confidence level






# Other searches outlook: HH

Extend the search of heavy resonances also to other final states and theoretic models

## Warped extra-dimension model:

arXiv: 1303.6636


- Explains hierarchy between the Electroweak and the Planck scales
- Non trivial geometry a fifth extra dimension
- Higgs pair production by:
  - spin 2 particle (Kaluza-Klein (KK) Graviton)
  - scalar particle (Radion)






# Search strategy

- Radion or Graviton decaying into a pair of Higgs bosons




## Strategy

- *High statistics*
- *Purity of the signal*




# Search strategy

- Radion or Graviton decaying into a pair of Higgs bosons




## Strategy

- *High statistics*
  - High branching ratio:  $H \rightarrow b\bar{b}$  ( $\sim 60\%$ )
- *Purity of the signal*




# Search strategy

- Radion or Graviton decaying into a pair of Higgs bosons



## Strategy


- *High statistics*
  - High branching ratio:  $H \rightarrow b\bar{b}$  ( $\sim 60\%$ )
- *Purity of the signal*
  - $H \rightarrow \tau^-\tau^+$  ( $\sim 7\%$ )



**Similar final state wrt ZH analysis:  
lepton ID, jet substructure and b-tagging**



# Conclusions

- The first search for a high mass narrow resonances decaying into Z and H bosons has been performed
  - Heavy resonances  highly boosted bosons
    - Jets with merged tau leptons or b quark pairs are most common for Higgs decays, but experimentally very challenging
    - Algorithms have been developed for the physics object and the event reconstruction
- Searching for physics BSM predicting HH resonances, e.g. Composite Higgs or Extra Dimension models
  - First time we look for such high mass resonances in Higgs pairs

Analysis of 8 TeV data well underway!


Stay tuned for the next run at 13/14 TeV!



Back up

# Search strategy

- Radion or Graviton decay into a pair of Higgs bosons
- Channel  $HH \rightarrow \tau\tau bb$ 
  - High breaching ratio
  - The presence of tau leptons can help discriminate against QCD Multi-jet background.
- Many possible final states depending on the tau lepton decay mode:
  - Fully leptonic:  $\tau \rightarrow \mu \nu \nu$ ,  $\tau \rightarrow e \nu \nu$
  - Semileptonic:  $\tau \rightarrow \mu \nu \nu$ ,  $\tau \rightarrow h \nu$
  - All hadronic:  $\tau \rightarrow h \nu$






Decay channel	BR(%)
$\tau$	17.36
$\tau$	17.85
$\tau$	11.6
$\tau$	26.0
$\tau$	9.5
$\tau$	9.8
$\tau$	4.8
others	3.1



# LHC and CMS

- Heavy particles production require high energy.
- LHC 8 TeV of energy in the center of mass reference frame
- CMS detector for particle identification



# Higgs Pair Production

arXiv: 1303.6636

- Metric due to the fifth extra dimension

$$ds^2 = e^{-2ky} \eta_{\mu\nu} dx^\mu dx^\nu - dy^2$$

$$g_{\mu\nu} = e^{-2ky} \eta_{\mu\nu} \rightarrow e^{-2(ky+F(x,y))} (\eta_{\mu\nu} + G_{\mu\nu}(x,y))$$





$$F(x,y) \propto e^{2ky} \phi(x) \quad G_{\mu\nu}^{(1)}(x,y) \propto e^{2ky} J_2 \left( e^{2ky} \frac{m_G}{k} \right) G_{\mu\nu}^{(1)}(x)$$

- Localization of the fields
- Coupling to SM fields

$$\mathcal{L} = -\frac{c_i}{\Lambda_G} G^{\mu\nu(1)} T_{\mu\nu}^i - \frac{d_i}{\Lambda_\phi} \phi T_\mu^{\mu i}$$

RS1 scenario:  $c_H =$  all the other  $c_i \simeq \mathcal{O}(1)$   
 Bulk RS scenario:  $c_H \simeq c_{Z,W,t} \simeq \mathcal{O}(1) \simeq (kL) c_{\gamma,g} \gg c_{u,d,l,\dots}$


Profiles





# Search strategy

- Radion or Graviton decay into a pair of Higgs bosons
- Channel  $HH \rightarrow \tau\tau bb$ 
  - High branching ratio
  - The presence of tau leptons can help discriminate against QCD Multi-jet background.
- Many possible final states depending on the tau lepton decay mode:
  - Fully leptonic:  $\tau \rightarrow \mu \nu \nu$ ,  $\tau \rightarrow e \nu \nu$
  - Semileptonic:  $\tau \rightarrow \mu \nu \nu$ ,  $\tau \rightarrow h \nu$
  - All hadronic:  $\tau \rightarrow h \nu$





Decay channel	BR(%)
$\tau$	17.36
$\tau$	17.85
$\tau$	11.6
$\tau$	26.0
$\tau$	9.5
$\tau$	9.8
$\tau$	4.8
others	3.1



# PF particle Identification


- Particle Flow: all the information from the sub detectors are combined to reconstruct all particles in the collisions
- Particles are identified in mutually exclusive categories: charged and neutral hadrons, photons, electron and muons.
- Particles are used to identify jets, tau and MET



- Algorithms have been developed to reconstruct the tau hadron decay:
  - HPS

# PF Tau Identification

- Starting from a PF Jet, special attention is given to photon conversion in the tracker, since the bending of electron positron can broaden the photon signature in the  $\phi$
- A strip of  $\Delta\eta = 0.05$  and  $\Delta\phi = 0.2$  is centered around the most energetic electromagnetic particle in the jet
- Other electromagnetic particles in the strip are considered and added to the four momentum.
- Strips with  $p_T > 1$  GeV/c are combined to the hadrons to reconstruct the Tau hadronic decay mode



Strips and hadrons ( $\pi^\pm, K^\pm$ ) are combined to reconstruct the main decay topologies:

- Single hadron (for  $h^- \nu_\tau$  and  $h^- \pi^0 \nu_\tau$ );
- One hadron + 1 or 2 strips (for  $h^- \pi^0 \nu_\tau$ );
- Three hadrons (for  $h^- h^+ h^- \nu_\tau$ ).

The other decay modes are reconstructed via the previous topologies.

- All hadrons and strips have to be within a cone of  $\Delta R = (2.8 \text{ GeV}/c) / p_T^{(\text{Tau}_h)}$
- The tau 4-vector has to be in a cone of  $\Delta R = 0.1$  from the Jet axis
- The mass of the composite system has to be compatible with  $\rho$  (770 MeV) (2 hadrons) or  $a_1$  ( $\geq 3$  hadrons 1200 MeV)



# Searches at low mass (<1TeV)

CMS (CMS PAS HIG-13-032):

hh ->gamma gamma bb final state.

The search for a new particle X is performed in the range  $260 < m_X < 1100$  GeV.

Upper limits at 95%confidence-level are extracted on new particles production cross-section.

WED Radion is observed (expected) to be excluded with masses below 0.97 TeV (0.88 TeV).

ATLAS(arXiv:1406.5053):

hh ->gamma gamma bb final state.


A 95% confidence level upper limit on the cross section times branching ratio of non-resonant production is set at 2.2 pb, while the expected limit is 1.0 pb.

The corresponding limit observed for a narrow resonance ranges between 0.8 and 3.5 pb as a function of its mass.


ATLAS (ATLAS-CONF-2014-005):

$G^* \rightarrow hh \rightarrow b\bar{b}b\bar{b}$  final state.


arXiv:1406.5053






CMS PAS HIG-13-032



ATLAS-CONF-2014-005



# Results



Observed and expected limits for the fully leptonic, semi leptonic and fully hadronic channels



# Systematics

Source	$e\mu$ channel	$\mu\mu$ channel	$ee$ channel	$\mu\tau_h$ channel	$e\tau_h$ channel	$\tau_h\tau_h$ channel
Bkg estimation method (N of sideband events)	0-8	0-10	0-3	0-26	0-23	29
Bkg estimation method (alpha-ratio)	25%-54%	23%-43%	35%-53%	13%-24%	13%-96%	-
Bkg estimation method (normalization)	35.7%	25.4%	46.7%	17.7%	29.1%	-
Luminosity	2.6%	2.6%	2.6%	2.6%	2.6%	2.6%
Pile-up	0.2%-2.2%	1.0%-1.9%	0.4%-1.5%	0.2%-1.4%	0.7%-1.2%	0.1%-1.0%
V-tag	6.4%	6.4%	6.4%	6.4%	6.4%	6.4%
B-tag	1.8%-3.0%	2.3%-2.5%	1.9%-5.6%	2.3%-2.7%	2.4%-2.6%	-
Jet scale	< 0.5%-1.6%	0.7%-2.6%	< 0.5%-4.3%	0.5%-1.5%	< 0.5%-1.4%	1.0%-3.1%
Jet resolution	< 0.5%-0.9%	< 0.5%-1.6%	< 0.5%-2.2%	< 0.5%-1.8%	< 0.5%-1.7%	2.0%-2.9%
Electron ID	1.1%-1.6%	-	2.5%-3.5%	-	1.3%-1.8%	-
Electron resolution	0.1%-0.6%	-	0.1%-2.4%	-	0.2%-0.7%	-
Electron scale	0.4%	-	0.1%-0.9%	-	0.1%-0.4%	-
Muon ID	0.8%-1.0%	5.7%	-	0.8%-0.9%	-	-
Muon resolution	< 0.5%	< 0.5%-1.8%	-	< 0.5%	-	-
Muon scale	< 0.5%-1.1%	< 0.5%-1.9%	-	< 0.5%-0.8%	-	-
Lepton modified iso	-	-	-	1.2%-14.3%	3.5%-20.8%	-
Tau ID	-	-	-	8.9%-12.4%	8.5%-11.9%	17.9%-26.2%
Tau Scale	-	-	-	< 0.5%-1.1%	< 0.5%-2.4%	0.7%-4.2%
Tau - jet cleaning	-	-	-	0.4%-7.0%	0.5%-15.7%	-
Tau - subjet	-	-	-	-	-	10%
MET	Included in leptons and jet energy uncertainties					





# Jet reconstruction


- Reconstruction of the hadronically decaying boson
- Analysis of the substructure of the jets for discriminating wrt QCD jets
  - Main idea: set of requirements during the jet clustering algorithm to “prune” the jet, i.e. to remove constituents that are at large angles or soft.
  - N-subjettiness helps discriminate between a jet that has 2 subjets or a jet that doesn't have substructures

$$\tau_N = \frac{1}{d_0} \cdot \sum_L [p_{T,k} \cdot \min(\Delta R_{1,k}, \Delta R_{2,k}, \dots, \Delta R_{N,k})]$$


CMS Work in Progress



CMS Work in Progress



CMS Work in Progress



# Background evaluation Method

- **Fully leptonic channel:** DY is the main background source
- Alpha-ratio method:

$$N_{bkg}(M_{ZH}) = \mathcal{N} \cdot N_{sb}(M_{ZH}) \cdot \alpha(M_{ZH})$$

- where:
  - $\mathcal{N}$  = normalization found in data
  - $N_{sb}(M_{HZ})$  is the number of events in the sideband in data
  - $\alpha(M_{HZ})$  is ratio between the MC bkg events in the signal and sideband region
- Sideband defined:
  - ✓ events with  $\tau_{21} < 0.75$  and  $20 < M(\text{pruned jet}) < 70 \text{ GeV} \parallel M(\text{pruned jet}) > 140 \text{ GeV}$
  - ✓ events with  $\tau_{21} > 0.75$  and  $M(\text{pruned jet}) > 20 \text{ GeV}$






# Background evaluation Method

- Fully leptonic channel:

$$N_{bkg}(M_{ZH}) = \mathcal{N} \cdot N_{sb}(M_{ZH}) \cdot \alpha(M_{ZH})$$



- Fit the pruned jet mass distribution in the MC in [20,200]GeV with the the function

$$F(x) = e^{Ax} \cdot \frac{1 + \text{erf}((x - B)/C)}{2}$$


- Fit data in the range [20,70] + [140,200] with the shapes found before, leaving the normalization unconstrained
- Extrapolate the number of events in the signal region [70,100] GeV



# Background evaluation Method



- Fully leptonic channel:

$$N_{bkg}(M_{ZH}) = \mathcal{N} \cdot N_{sb}(M_{ZH}) \cdot \alpha(M_{ZH})$$



# Background evaluation Method


- Fully leptonic channel:



## RESULTS

CMS Work in Progress


Channel	$e\mu$	$\mu\mu$	$ee$
BKG prediction	$2.7 \pm 1.0$	$5.9 \pm 1.5$	$1.5 \pm 0.7$
BKG in MC	$3.4 \pm 0.5$	$6.1 \pm 0.8$	$1.7 \pm 0.4$
DATA	3	8	2



# Background evaluation Method

- **Semi leptonic channel:** DY is the main background source
- Alpha-ratio method as before but with the ttbar contribution fixed.
- Top events normalization estimated in a ttbar enriched control sample (at least a b-tagged jet-CSVM)
- Fit of the pruned mass distribution using the function

$$F_{t\bar{t}}(x) = N(\text{non-peaking}) \cdot e^{Ax} \cdot \frac{1 + \text{erf}((x - B)/C)}{2} + N(\text{peaking}) \cdot \text{Gauss}(D, E)$$





- The two SFs are found as the ratio between data and MC of the normalization of the two functions

	$\tau_{\mu}\tau_h$	$\tau_e\tau_h$
SF(peak)	$1.8 \pm 1.3$	$1.7 \pm 2.1$
SF(NO peak)	$0.9 \pm 0.1$	$0.9 \pm 0.1$




# Background evaluation Method

- Semi leptonic channel:



# Background evaluation Method

- Semi leptonic channel:



CMS Work in Progress


**RESULTS**

Channel	$\mu\tau_h$	$e\tau_h$
BKG prediction	$20.7 \pm 3.7$	$18.6 \pm 5.6$
BKG in MC	$18.3 \pm 1.4$	$19.3 \pm 1.5$
DATA	28	12



# Background evaluation Method


- Fully hadronic channel:
  - ABCD Method:
  - $M_{\text{pruned}}$  and  $M_{\tau\tau}$  from SVFit are used: uncorrelated variables
    - Number of background events evaluated through:



$$N(A) = N(D) \cdot \frac{N(B)}{N(C)}$$

# Background evaluation Method

- Fully hadronic channel:



## RESULTS

CMS Work in Progress

Regions	Events in Data - Original
NB	$9.0^{+4.1}_{-2.9}$
NC	$43^{+8}_{-7}$
ND	$29^{+6}_{-5}$
NA(estimate)	$6.1^{+3.2}_{-2.5}$
Observed A	8





# Systematics

- The largest systematics are the ones due to the background evaluation
- When the number of events is evaluated from a control sample the distribution for  $N_{bkg}$  is a Gamma function

$$n = \alpha \cdot N \quad \Gamma(n) = \frac{(n/\alpha)^N}{\alpha N!} e^{-n/\alpha}$$

CMS Work in Progress

	1.0 TeV	1.5 TeV	2.0 TeV	2.5 TeV
$\tau_e \tau_\mu$	8	1	0	0
$\tau_\mu \tau_\mu$	10	6	3	1
$\tau_e \tau_e$	3	1	0	0
$\tau_\mu \tau_h$	26	11	0	0
$\tau_e \tau_h$	23	11	2	1
$\tau_h \tau_h$	29			

Number of events in the sideband

The other two systematics associated to the background estimation method for the fully and semi leptonic channels are:


- ✓ statistical error on the alpha ratio
- ✓ statistical error on normalization factor



# Pruning

- Developed by **S.Ellis et al:** [Phys. Rev. n D80 (2009)]
- Main idea → add a set of requirements during the clustering algorithm
  - **to prune the jet:** remove constituents that are at large angles or soft
- **Starting point:** jets clustered with CA algorithm and distance parameter of 0.8
- Steps of the algorithm:
  1. **rerun** the clustering sequence
  2. **two more requirements** are asked:

- $z_{ij} \equiv \frac{\min(p_T^i, p_T^j)}{p_T^p} > z_{cut}$  → remove soft particles
- $\Delta R_{ij} < \alpha \cdot m_{jet}/p_T^{jet}$  → remove large angle particles



ref. CMS PAS JME-13-006

