

Where is sWaldo?

Searching for sTops amongst the SM crowd with the **ATLAS** Detector

arXiv [hep-ex]1407.0583

Benjamin Nachman
SLAC, Stanford University

October 7, 2014

Other SLAC people involved:
Till Eifert, Philippe Grenier, Qi Zeng



Motivation

A minimal SUSY solution to the hierarchy problem requires light stops



Gospel Church Choir Cats Cat Art By Taraflly



Copyright © 2011 DC Comics - from <http://en.wikipedia.org/wiki/File:Superman.jpg>, see copyright information there.

There is no (approximate) symmetry to protect the Higgs mass from Quantum Corrections





Image from Flip Tanedo (Quantum Diaries)

Naively, the mass receives quadratic corrections to highest mass scale

The largest (quantum) contribution comes from the **[fermionic]** top quark loop - can cancel with **[scaler]** stop loops



The parameter space of natural SUSY is huge


Technically, natural SUSY is a framework, not a model

The Minimal Supersymmetric SM has $\mathcal{O}(100)$ parameters

Simplifying assumptions


1. R -parity conservation
2. stop is the only light squark
3. lightest neutralino is the LSP

Three important parameters:




Target Models

We use simplified models: Leading order processes with 100% branching ratios



Most important parameter
- sets the cross section




Stop is (mostly) right-handed and the neutralino is (mostly) bino



How 'realistic' is simple?

There are regions of the MSSM with near unity branching ratios

1212.3526: G. Belanger et al.



Right-handed top




1404.7554: S. Dimopoulos, K. Howe, and J. March-Russell

The neutralino field content is important in part for the top polarization

→ impacts acceptance through the momentum of the top decay products

Another motivation: *Maximally Natural SUSY*; gluinos are heavy and it look likes simplified models

This is not the MSSM!




Not happy with simple?


We will also consider *asymmetric* decays where

$$BR(b+chargino) + BR(t+LSP) = 1$$

$$\text{but } BR(b+chargino) < 1 \text{ \& } BR(t+LSP) < 1$$



We also will study the impact of the stop mixing (impacts top polarization and acceptance)



Still not happy?

Just for you, we also considered a scan in the pMSSM*

(R-parity conserving MSSM subject to experimentally motivated constraints - 19 parameters)

*Thanks to the work of M. Cahill-Rowley, J.L. Hewett, A. Ismail, and T.G. Rizzo

Search Strategy

1 lepton channel: optimal mix of cross section and background rejection

(1) Preselection

- Reach the trigger plateau
- Remove most multijet events

(2) Discriminating Variables

- Robust techniques to isolate the signal

Many designed specifically for this search


- Combine variables to form signal regions (SRs)

(3) Background estimation

- For the dominant backgrounds, define control regions (CRs)
- Estimate systematic uncertainties

(4) Results

Top Pair Branching Fractions



Credit: D0 Collaboration: http://www-d0.fnal.gov/Run2Physics/top/top_public_web_pages/

Preselection

Trigger: (Single Isolated e or μ) or E_T^{miss}

- $E_T^{\text{miss}} > 100 \text{ GeV}$

$> 24 \text{ GeV @ HLT}$


$> 80 \text{ GeV @ HLT}$

- Exactly one isolated e or μ with $p_T > 25 \text{ GeV}$


- No other e or μ with $p_T > 10 \text{ GeV}$

- At least one b-jet @ 70% efficiency

- At least four jets (anti- k_t $R=0.4$)



SRs target particular regions of phase space




Two main discriminating variables

Transverse Mass

$$m_T = \sqrt{2 \cdot p_T^\ell \cdot E_T^{\text{miss}} (1 - \cos \Delta\phi(\vec{\ell}, \vec{p}_T^{\text{miss}}))}.$$


For semileptonic $t\bar{t}$, $m_T \leq m_W$



1L top 2L top


Even though our signal has 1L,
after m_T , background mostly 2L

Nearly uncorrelated



Main Backgrounds:

- (dileptonic) $t\bar{t}$
- W +jets



There is an m_{T2} for you!

Transverse mass is a powerful variable because it takes advantage of the targeted topology




There is a class of variables which generalize the transverse mass to multiple invisible particles

$$m_{T2} \equiv \min_{\vec{q}_{Ta} + \vec{q}_{Tb} = \vec{p}_T^{\text{miss}}} \{ \max(m_{Ta}, m_{Tb}) \},$$

$$m_{Ti}^2 = \left(\sqrt{p_{Ti}^2 + m_{p_i}^2} + \sqrt{q_{Ti}^2 + m_{q_i}^2} \right)^2 - (\vec{p}_{Ti} + \vec{q}_{Ti})^2$$


(usual transverse mass)




With the appropriate choices of m_{q_i} , these variables have endpoints for the background

m_{T2} for the stop 1L search



After E_T^{miss} and m_T requirements, dominant background has **two** leptons



am_{T2}



m_{T2}



Including Resolution Information: *Significance Variables*

Mis-measurement can induce large E_T^{miss}

This motivated the E_T^{miss} *significance*:


$$E_T^{\text{miss}} / \text{Uncertainty}(E_T^{\text{miss}})$$

It is common to approximate
the uncertainty as $\sim 0.5 \times \sqrt{H_T}$

We can improve this by using known η and
 p_T dependent resolution functions for jets

$$H_{T,\text{sig}}^{\text{miss}} = \frac{|\vec{H}_T^{\text{miss}}| - M}{\sigma_{|\vec{H}_T^{\text{miss}}|}},$$

where H_T^{miss} is the vector sum of all
measured identified objects



-Shameless self promotion-

*Generalizes to any
kinematic variable*

See 1303.7009
(BN and C. G. Lester)

Many other discriminating variables have been developed to suppress the two lepton background

Hadronic Top Mass

-dileptonic $t\bar{t}$ has no hadronic top -

$$m_{\text{had-top}} = \operatorname{argmin}_{m_{bjj}} \left\{ \frac{(m_{bjj} - m_{\text{top}})^2}{\sigma_{m_{bjj}}^2} + \frac{(m_{jj} - m_W)^2}{\sigma_{m_{jj}}^2} \right\}$$

Topness **1212.4495: M. Graesser and J. Shelton**

One lost lepton; reconstruct the event by minimizing a χ^2 kinematic compatibility with the dileptonic hypothesis

Hadronic Taus: reconstruct and veto


τ candidates: 1-3 tracks & $Q_\tau \times Q_{\text{lepton}} < 0$

A BDT based on track and calo orientations for rejecting QCD jets

Isolated Tracks $Q_{\text{track}} \times Q_{\text{lepton}} < 0$

Reject events with a well-isolated Hard Scatter (HS) track with $p_T > 10$ GeV

Isolation: no HS tracks with $p_T > 3$ GeV in $\Delta R < 0.4$





High stop mass = Boosted Tops

Hadronic Top Mass breaks down

When the p_T of the tops is high enough, jets begin to merge

$$\Delta R \sim 2m/p_T$$

For $p_T^{\text{top}} \sim m_{\text{stop}}/2$, $\Delta R \sim 1$ for $m_{\text{stop}} \sim 700 \text{ GeV}$



Use $R = 1.0$ anti- k_t trimmed **jet mass**
 ($f_{\text{cut}} = 0.05$ $R_{\text{sub}} = 0.3$)
 for a powerful hadronic top mass discriminant

The direction of the large radius jets is also powerful




For SM, second top is leptonic, which is close to the p_T^{miss}

Compressed Spectra: Low E_T^{miss}

Signal Resembles the Background

Recover sensitivity by fitting the **shapes** of kinematic variables



Bin the relevant distribution(s) with bins chosen to have a distribution of S/B

We have used 2D shape fits using \mathbf{m}_T and one of $\mathbf{E}_T^{\text{miss}}$ or \mathbf{am}_{T2}


t+Neutralino SRs

A highlight; for more details see the paper

Medium Kinematic Requirements

Tight Kinematic Requirements

**Shape fit in m_T and am_{T2}
(low am_{T2} most sensitive)**




b+Chargino SRs

A highlight; for more details see the paper

2 high p_T b-jets @ 80%
 am_{T2} most effective
at reducing top backgrounds

$am_{T2} \gg m_{top}$

Loose E_T^{miss}
Shape fit in m_T and am_{T2}



Signal Regions
 bCa_low,
 bCa_med

bCb_med1,
 bCb_med2,
 bCb_high

bCc_diag

bCd_bulk,
 bCd_high1,
 bCd_high2

2 high p_T b-jets @ 80%
Shape fit in m_T and am_{T2}

Require only 3 jets and veto b-jets
Loose E_T^{miss} and m_T

b+Chargino SRs

A highlight; for more details see the paper

Unlike other regions, there is a non-negligible QCD multijet background

For low chargino-neutralino mass splitting, use **soft leptons**: $p_T > 6(7)$ GeV for $e(\mu)$

leptons: $p_T > 6(7)$ GeV for $e(\mu)$

Veto extra jets beyond 2 leading
Loose E_T^{miss}


Medium E_T^{miss}

at least 2 jets
Tight E_T^{miss}

at least 3 jets
Medium E_T^{miss}

1st jet not b-tagged (ISR)
 $m_T \gtrsim m_W$
Shape fit in lepton p_T


at least 2 jets
2 b-tags @ 60%
Large m_{bb}
Shape fit in am_{T2}



Background Estimation

Background Estimation

Every SR has two dedicated Control Regions (CRs) for data-driven estimates of the dominant backgrounds



*Top CR: invert
 $m_T < m_W$*

*W+jets CR: invert
 $m_T < m_W$
b-veto instead of b-tag*

*QCD multijets estimated in the data
by loosening lepton isolation*


Shape Fits: Background Integrated into Fit

Every shape fit has regions which ‘act’ like CRs

In the fit, they are treated like all other bins

Can normalize per E_T^{miss} bin to maximize sensitivity



(Trade off syst for stat uncertainty)



Data-Driven Validation: $t\bar{t}Z(\rightarrow \nu\bar{\nu})$ from $t\bar{t}\gamma$

Except for neutrinos and their mass, the Feynman diagrams are identical

At high p_T , boson mass irrelevant so the yield of one can predict the other





Define new m_T and E_T^{miss} variables with the photon treated as invisible

Many theoretical and experimental uncertainties should cancel in the extrapolation to the SR

Additional Validation

Beyond LO radiation in dilepton top events

W+Heavy Flavour
Jet veto + $m_{bb} < 80$ GeV




Additional checks with taus and isolated tracks show generally good modeling of the data

Results




95% Confidence Level

*Reminder for reading
exclusion plots*

- Line: Observed Limit
- ==== Band: Signal Theory Uncertainties
- Line: Expected Limit
- Band: All other Uncertainties





t+Neutralino Results





'Stealth Stop'
 Difficult to reach with direct searches

Reach limited by cross section:
 10^{-5} x top cross section

t+Neutralino Results




t+Neutralino Results



*Probably not; **CMS** sees a deficit where we see an excess*



Stop Mixing

Nominal mixing: $\sim 70\%$ stop right $P_t \sim +1$



~ 50 GeV weaker limits for the left-handed stop
 (true for a range of neutralino mass hypotheses)

For a left handed stop, $P_t < 0$




b+Chargino Results

To show 2D exclusions, need a hypothesis on $m_{\text{chargino}} = f(m_{\text{neutralino}}, m_{\text{stop}})$

Soft b-jet p_T spectrum
3 jets (ISR) and veto b-jets

One common form is $f(x,y)=2x$
(Gaugino Universality)



Why $m_{\text{LSP}} > 50$ GeV?
LEP limit for charginos is ~ 100 GeV

W off-shell

$m_{\text{stop}} \sim m_{\text{top}}$

W off-shell, chargino acts as 'W'
Looks like SM top!

Exploit Kinematic Shapes
Shapes
Fit in m_T and am_{T2}



Hard b-jet p_T spectrum
Require 2 b-jets with high p_T
Tight am_{T2} threshold

b+Chargino Results


To show 2D exclusions, need a hypothesis on $m_{\text{chargino}} = f(m_{\text{neutralino}}, m_{\text{stop}})$

Soft b-jet p_T spectrum
3 jets and veto b-jets

One common form is $f(x,y)=2x$
(Gaugino Universality)



Hard b-jet p_T spectrum
Tight am_{T2} threshold




b+Chargino Results

To show 2D exclusions, need a hypothesis on $m_{\text{chargino}} = f(m_{\text{neutralino}}, m_{\text{stop}})$


Another choice, $f(x,y) = 150 \text{ GeV}$
(why 150? Because > LEP limit)

[Many other choices for f in backup]




bCa_low/med use ISR in the selection and so limited to lower cross sections (stop masses)

Phase-Space Only



$m_{bb} > 150 \text{ GeV}$ requirement
 Limits power at low m_{stop}



Results: Mixed Decays


Gaugino Universality:

$$m_{\text{chargino}} = 2 \times m_{\text{LSP}}$$

$\tilde{t}_1, \tilde{t}_1^*$ production, $\tilde{t}_1 \rightarrow t \tilde{\chi}_1^0 / b \tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow W^{(*)} + \tilde{\chi}_1^0, m_{\tilde{\chi}_1^\pm} = 2 m_{\tilde{\chi}_1^0}$

*Generate events with one stop decaying to **bC** and one to **tN***

*A superposition of the 100% BR models give the **mixed models***



b+Chargino
Exclusion

SR optimized for
BR = 50%

t+Neutralino
Exclusion

Continuous exclusion between simplified models

Results: pMSSM

M. Cahill-Rowley, J.L. Hewett, A. Ismail, and T.G. Rizzo produced a large scan of the 19 parameter pMSSM


Models required to get $m_h \sim 125$ GeV, saturate the DM relic density and have low FT.

~ 10 k models with a neutralino LSP

We take a subset of 27 in three small mass ranges


Excluded at 95% CL

**Less 'simplified',
less sensitivity**





8 TeV **ATLAS** Stop Summary

ATLAS has a comprehensive program in direct stop searches



Largest coverage with 1L





Stealth Stop from Top Properties



Prospects for 14 TeV

At the higher stop masses, we will gain from higher **cross sections** and **more data**

ATL-PHYS-PUB-2013-011




There will be new challenges, as the **top cross section** also increases; objects are more **boosted** and formerly **subdominant backgrounds** are now important

Conclusions


*We have searched extensively for a natural stop using the 8 TeV **ATLAS** dataset*

We have developed many new techniques to search for stops in the range $\sim 200\text{-}700$ GeV



Many regions of simplified and not-so-simplified parameter space excluded


No evidence (yet) of SUSY



Stay tuned for 13 TeV results next year!

BACKUP



Large Radius Jets




Jet mass squared = sum of the constituent 4-vectors, squared

Why low am_{T2} for 3body?

arXiv:03042226



3body more sensitive to the unbalanced solution



pMSSM Models

\tilde{t}_1	Mass [GeV]					Branching ratio $\tilde{t}_1 \rightarrow$					$[T_{11}]^2$	$[N_{11}]^2$
	$\tilde{\chi}_1^0$	$\tilde{\chi}_2^0$	$\tilde{\chi}_3^0$	$\tilde{\chi}_1^\pm$	$\tilde{\chi}_2^\pm$	$t\tilde{\chi}_1^0$	$t\tilde{\chi}_2^0$	$t\tilde{\chi}_3^0$	$b\tilde{\chi}_1^\pm$	$b\tilde{\chi}_2^\pm$		
404	40	221	230	220	1073	0.09	0.01	0.09	0.81	0.00	0.53	0.96
404	44	324	445	325	471	0.16	0.00	0.00	0.84	0.00	0.98	0.99
407	46	368	372	367	1515	0.74	0.00	0.00	0.26	0.00	0.02	0.98
408	49	187	207	188	376	0.02	0.31	0.23	0.41	0.04	0.97	0.95
409	39	211	212	206	1768	0.05	0.24	0.02	0.68	0.00	0.56	0.95
409	49	180	190	179	795	0.02	0.22	0.17	0.59	0.00	0.99	0.94
410	40	232	253	234	427	0.11	0.25	0.00	0.64	0.00	0.96	0.97
410	43	387	396	386	889	0.88	0.00	0.00	0.12	0.00	0.01	0.99
413	42	197	367	197	385	0.03	0.10	0.00	0.85	0.02	0.95	0.98
413	45	373	406	374	508	0.32	0.00	0.00	0.68	0.00	0.99	0.99
414	45	194	440	195	453	0.03	0.14	0.00	0.83	0.00	0.96	0.99
416	45	394	397	393	1975	0.90	0.00	0.00	0.10	0.00	0.99	0.99
417	46	333	350	335	573	0.65	0.00	0.00	0.35	0.00	0.96	0.98
418	39	206	209	202	1779	0.09	0.05	0.28	0.59	0.00	0.47	0.95
546	46	292	310	292	520	0.02	0.28	0.24	0.44	0.01	0.98	0.98
547	46	346	374	346	500	0.12	0.49	0.00	0.22	0.16	0.93	0.98
550	40	225	235	225	760	0.02	0.28	0.24	0.46	0.00	0.98	0.96
551	43	351	366	351	621	0.07	0.38	0.21	0.35	0.00	0.98	0.99
552	41	249	275	252	420	0.02	0.20	0.21	0.44	0.13	0.98	0.97
552	42	332	337	331	1496	0.05	0.47	0.35	0.13	0.00	0.99	0.98
552	43	346	350	344	1501	0.08	0.27	0.52	0.13	0.00	0.97	0.98
552	43	385	397	385	731	0.36	0.00	0.00	0.64	0.00	0.97	0.99
554	44	439	445	439	1007	0.21	0.00	0.00	0.79	0.00	0.99	0.99
555	47	279	287	280	933	0.04	0.54	0.38	0.04	0.00	0.97	0.97
553	147	169	444	168	455	0.31	0.12	0.00	0.27	0.30	0.07	0.93
554	151	195	207	191	1969	0.09	0.35	0.43	0.12	0.00	0.88	0.68
546	154	210	213	200	434	0.07	0.40	0.34	0.05	0.14	0.86	0.70

Table 1. Properties of the 27 selected pMSSM models. The table contains the masses of the stop, of neutralinos and of the charginos, the branching ratios of the stop decays, the \tilde{t}_L content of the \tilde{t}_1 ($[T_{11}]^2$, with T being the stop mixing matrix) and the bino content of the χ_1^0 ($[N_{11}]^2$, with N being the neutralino mixing matrix).

Soft Lepton Selections

	bCa_low	bCa_med	bCb_med1	bCb_high
Preselection	soft-lepton preselection, cf. table 3.			
Lepton	= 1 soft lepton		= 1 soft lepton with $p_T < 25$ GeV	
Jets	≥ 2 with $p_T > 180, 25$ GeV	≥ 3 with $p_T > 180, 25, 25$ GeV	≥ 2 with $p_T > 60, 60$ GeV	
Jet veto	–		$H_{T,2} < 50$ GeV	–
<i>b</i>-tagging	$\geq 1b$ -tag amongst sub-leading jets (70% eff.)		Leading two jets <i>b</i> -tagged (60% eff.)	
<i>b</i>-veto	1 st jet not <i>b</i> -tagged (70% eff.)		–	
m_{bb}	–		> 150 GeV	
E_T^{miss}	> 370 GeV	> 300 GeV	> 150 GeV	> 250 GeV
$E_T^{\text{miss}}/m_{\text{eff}}$	> 0.35	> 0.3	–	
m_T	> 90 GeV	> 100 GeV	–	
Exclusion setup: shape-fit				
	4 bins in lepton p_T range [6(7), 50] GeV		6 bins in am_{T2} range [0, 500]GeV	
Discovery setup				
	lepton $p_T < 25$ GeV		$am_{T2} > 170$ GeV	$am_{T2} > 200$ GeV

b+Chargino Lepton Selections

	bCb_med2	bCc_diag	bCd_bulk	bCd_high1	bCd_high2
Preselection	Default preselection criteria, cf. table 3.				
Lepton	= 1 lepton	= 1 lepton with $ \eta(\ell) < 1.2$	= 1 lepton		
Jets	≥ 4 with $p_T > 80, 60, 40, 25$ GeV	≥ 3 with $p_T > 80, 40, 30$ GeV	≥ 4 with $p_T > 80, 60, 40, 25$ GeV		≥ 4 with $p_T > 80, 80, 40, 25$ GeV
b-tagging / veto	≥ 2 (80% eff.) with $p_T > 140, 75$ GeV	= 0 (70% eff.) with $p_T > 25$ GeV	≥ 1 (70% eff.) with $p_T > 25$ GeV	≥ 2 (80% eff.) with $p_T > 75, 75$ GeV	≥ 2 (80% eff.) with $p_T > 170, 80$ GeV
E_T^{miss}	> 170 GeV	> 140 GeV	> 150 GeV		> 160 GeV
m_T	> 60 GeV	> 120 GeV	> 60 GeV	> 120 GeV	
$E_T^{\text{miss}}/\sqrt{H_T}$	> 6 GeV ^{1/2}	> 5 GeV ^{1/2}	> 7 GeV ^{1/2}	> 9 GeV ^{1/2}	> 8 GeV ^{1/2}
am_{T2}	> 80 GeV	–	> 80 GeV	> 200 GeV	> 250 GeV
Track, τ-veto	track & loose τ -veto	–	track & loose τ -veto		
$\Delta R(j_1, \ell)$	–	$\in [0.8, 2.4]$	–		
$\Delta\phi(\text{jet}, \vec{p}_T^{\text{miss}})$	> 0.8 (1 st and 2 nd jet)	> 2.0 (1 st jet), > 0.8 (2 nd jet)	> 0.8 (1 st and 2 nd jet)		
Exclusion setup	shape-fit in m_T and am_{T2} , cf. figure 9.	cut-and-count	shape-fit in m_T and am_{T2} , cf. figure 9.	cut-and-count	
Discovery setup	test signal-sensitive bins.	cut-and-count	test signal-sensitive bins.	cut-and-count	


3body and Mixed SRs

	tNbC_mix	3body
Preselection	Default preselection criteria, cf. table 3.	
Lepton	= 1 lepton	
Jets	≥ 4 jets with $p_T > 80, 70, 50, 25$ GeV	≥ 4 jets with $p_T > 80, 25, 25, 25$ GeV
<i>b</i>-tagging	≥ 1 <i>b</i> -tag (70% eff.) with $p_T > 60$ GeV	≥ 1 <i>b</i> -tag (70% eff.) with $p_T > 25$ GeV
E_T^{miss}	> 270 GeV	> 150 GeV
m_T	> 130 GeV	> 60 GeV
am_{T2}	> 190 GeV	> 80 GeV
<i>topness</i>	> 2	–
m_{jjj}	< 360 GeV	–
$E_T^{\text{miss}} / \sqrt{H_T}$	> 9 GeV ^{1/2}	> 5 GeV ^{1/2}
τ-veto	loose	
$\Delta\phi(\text{jet}_i, \vec{p}_T^{\text{miss}})$	> 0.6 ($i = 1, 2$)	> 0.2 ($i = 1, 2$)
$\Delta\phi(\ell, \vec{p}_T^{\text{miss}})$	> 0.6	> 1.2
$\Delta R(\ell, \text{jet}_i)$	< 2.75 ($i = 1$)	> 1.2 ($i = 1$), > 2.0 ($i = 2$)
$\Delta R(\ell, b\text{-jet})$	< 3.0	–
Exclusion setup	cut-and-count	shape-fit in m_T and am_{T2} , cf. figure 6.
Discovery setup	cut-and-count	test signal-sensitive bins one-by-one.

top+Neutralino SRs

	tN_diag	tN_med	tN_high	tN_boost
Preselection	Default preselection criteria, cf. table 3.			
Lepton	= 1 lepton			
Jets	≥ 4 with $p_T >$ 60, 60, 40, 25 GeV	≥ 4 with $p_T >$ 80, 60, 40, 25 GeV	≥ 4 with $p_T >$ 100, 80, 40, 25 GeV	≥ 4 with $p_T >$ 75, 65, 40, 25 GeV
<i>b</i>-tagging	≥ 1 <i>b</i> -tag (70% eff.) amongst four selected jets			
large-<i>R</i> jet	–			≥ 1 , $p_T > 270$ GeV and $m > 75$ GeV
$\Delta\phi(\text{jet}_2^{\text{large-}R}, \vec{p}_T^{\text{miss}})$	–			> 0.85
E_T^{miss}	> 100 GeV	> 200 GeV	> 320 GeV	> 315 GeV
m_T	> 60 GeV	> 140 GeV	> 200 GeV	> 175 GeV
am_{T2}	–	> 170 GeV	> 170 GeV	> 145 GeV
m_{T2}^τ	–	–	> 120 GeV	–
<i>topness</i>	–	–	–	> 7
$m_{\text{had-top}}$	$\in [130, 205]$ GeV	$\in [130, 195]$ GeV	$\in [130, 250]$ GeV	
τ-veto	tight	–	–	modified, see text.
$\Delta R(b\text{-jet}, \ell)$	< 2.5	–	< 3	< 2.6
$E_T^{\text{miss}} / \sqrt{H_T}$	> 5 GeV ^{1/2}	–		
$H_{T,\text{sig}}^{\text{miss}}$	–	> 12.5		> 10
$\Delta\phi(\text{jet}_i, \vec{p}_T^{\text{miss}})$	> 0.8 ($i = 1, 2$)	> 0.8 ($i = 2$)	–	$> 0.5, 0.3$ ($i = 1, 2$)
Exclusion setup	shape-fit in m_T and E_T^{miss} , cf. figure 6.	cut-and-count		
Discovery setup	test signal-sensitive bins one-by-one.	cut-and-count		

Validation Regions




Results

Region	Obs.	Exp. bkg.	p_0	$N_{\text{non-SM}}$		σ_{vis} [fb]	
				Obs.	Exp.	Obs.	Exp.
tN_med	12	13.0 ± 2.2	≥ 0.5	8.5	9.2	0.4	0.5
tN_high	5	5.0 ± 1.0	≥ 0.5	6.0	6.0	0.3	0.3
tN_boost	5	3.3 ± 0.7	0.17	7.0	5.3	0.3	0.3
bCa_low	11	6.5 ± 1.4	0.08	12.2	7.8	0.61	0.92
bCa_med	20	17 ± 4	0.33	14.4	12.3	0.72	0.68
bCb_med1	41	32 ± 5	0.12	23.5	16.0	1.17	0.88
bCb_high	7	9.8 ± 1.6	≥ 0.5	6.5	7.9	0.32	0.22
bCc_diag	493	470 ± 50	0.27	110.6	95.1	5.4	4.7
bCd_high1	16	11.0 ± 1.5	0.09	13.2	8.5	0.7	0.4
bCd_high2	5	4.4 ± 0.8	0.36	6.3	5.7	0.3	0.3
tNbC_mix	10	7.2 ± 1.0	0.13	9.7	7.0	0.5	0.3
tN_diag							
$125 < E_{\text{T}}^{\text{miss}} < 150 \text{ GeV}, 120 < m_{\text{T}} < 140 \text{ GeV}$	117	136 ± 22	≥ 0.5	42.1	55.7	2.1	2.7
$125 < E_{\text{T}}^{\text{miss}} < 150 \text{ GeV}, m_{\text{T}} > 140 \text{ GeV}$	163	152 ± 20	0.35	55.4	47.8	2.7	2.4
$E_{\text{T}}^{\text{miss}} > 150 \text{ GeV}, 120 < m_{\text{T}} < 140 \text{ GeV}$	101	98 ± 13	0.43	36.1	33.9	1.8	1.7
$E_{\text{T}}^{\text{miss}} > 150 \text{ GeV}, m_{\text{T}} > 140 \text{ GeV}$	217	236 ± 29	≥ 0.5	58.7	71.4	2.9	3.5
bCb_med2							
$175 < am_{\text{T}2} < 250 \text{ GeV}, 90 < m_{\text{T}} < 120 \text{ GeV}$	10	12.1 ± 2.0	≥ 0.5	7.3	8.8	0.4	0.4
$175 < am_{\text{T}2} < 250 \text{ GeV}, m_{\text{T}} > 120 \text{ GeV}$	10	7.4 ± 1.4	0.10	9.7	7.3	0.5	0.4
$am_{\text{T}2} > 250 \text{ GeV}, 90 < m_{\text{T}} < 120 \text{ GeV}$	16	21 ± 4	≥ 0.5	9.3	12.3	0.5	0.6
$am_{\text{T}2} > 250 \text{ GeV}, m_{\text{T}} > 120 \text{ GeV}$	9	9.1 ± 1.6	≥ 0.5	7.7	7.8	0.4	0.4
bCd_bulk							
$175 < am_{\text{T}2} < 250 \text{ GeV}, 90 < m_{\text{T}} < 120 \text{ GeV}$	144	133 ± 22	0.29	36.1	33.9	1.8	1.7
$175 < am_{\text{T}2} < 250 \text{ GeV}, m_{\text{T}} > 120 \text{ GeV}$	78	73 ± 8	0.34	58.7	71.4	2.9	3.5
$am_{\text{T}2} > 250 \text{ GeV}, 90 < m_{\text{T}} < 120 \text{ GeV}$	61	66 ± 6	≥ 0.5	17.5	20.9	0.9	1.0
$am_{\text{T}2} > 250 \text{ GeV}, m_{\text{T}} > 120 \text{ GeV}$	29	26.5 ± 2.6	0.34	14.8	12.6	0.7	0.6
3body							
$80 < am_{\text{T}2} < 90 \text{ GeV}, 90 < m_{\text{T}} < 120 \text{ GeV}$	12	16.9 ± 2.8	≥ 0.5	7.3	9.9	0.4	0.5
$80 < am_{\text{T}2} < 90 \text{ GeV}, m_{\text{T}} > 120 \text{ GeV}$	8	8.4 ± 2.2	≥ 0.5	7.9	7.8	0.4	0.4
$90 < am_{\text{T}2} < 100 \text{ GeV}, 90 < m_{\text{T}} < 120 \text{ GeV}$	29	35 ± 4	≥ 0.5	11.7	14.7	0.6	0.7
$90 < am_{\text{T}2} < 100 \text{ GeV}, m_{\text{T}} > 120 \text{ GeV}$	22	29 ± 5	≥ 0.5	55.4	47.8	2.7	2.4

b+Chargino Results

To show 2D exclusions, need a hypothesis on $m_{\text{chargino}} = f(m_{\text{neutralino}}, m_{\text{stop}})$


Another choice, $f(x,y)=106 \text{ GeV}$
(why 106? Because \sim LEP limit)



b+Chargino Results

To show 2D exclusions, need a hypothesis on $m_{\text{chargino}} = f(m_{\text{neutralino}}, m_{\text{stop}})$


Another choice, $f(x,y) = x + 5 \text{ GeV}$



b+Chargino Results

To show 2D exclusions, need a hypothesis on $m_{\text{chargino}} = f(m_{\text{neutralino}}, m_{\text{stop}})$


Another choice, $f(x,y) = x + 20 \text{ GeV}$



b+Chargino Results

To show 2D exclusions, need a hypothesis on $m_{\text{chargino}} = f(m_{\text{neutralino}}, m_{\text{stop}})$

Another choice, $f(x,y)=y-10 \text{ GeV}$



Can this explain the WW excess (and other excesses)? see arXiv: 1406.0858

b+Chargino Results

$m_{\text{stop}} = 300 \text{ GeV}$

