

DIS 2015

XXIII International Workshop on
Deep-Inelastic Scattering and
Related Subjects

Dallas, Texas
April 27 – May 1, 2015



Contribution ID: 219

Type: **not specified**

Production of top pair events with additional radiation using the ATLAS detector at the LHC

Wednesday 29 April 2015 09:45 (20 minutes)

The large centre-of-mass energy available at the proton-proton collider LHC allows for the copious production of top quark pairs in association with other final state particles at high transverse momentum. The ATLAS experiment has measured several final state observables that are sensitive to additional radiation in top anti-top quark final states. Results on the top production in association with W and Z bosons are presented along with measurements of the cross section for production with an associated isolated photon. Analyses probing the top pair production with additional QCD radiation include the multiplicity of jets for various transverse momentum thresholds or the probability to emit jets above a given threshold in a fixed rapidity region. These measurements are compared to modern Monte Carlo generators based on NLO QCD matrix element or LO multi-leg matrix elements. The data are able to constrain the uncertainty on the modelling of the top pair production mechanism. We also discuss the production of top quark pairs in association with additional b-jets.

Primary author: GREENWOOD JR, Dick (Louisiana Tech University (US))

Presenter: SCHWANENBERGER, Christian (Deutsches Elektronen-Synchrotron (DE))

Session Classification: WG5 Heavy Flavours

Track Classification: WG5 Heavy Flavours