




Diffractive processes in pp collisions at 7 TeV measured with the CMS experiment




Grzegorz Brona
(University of Warsaw)
on behalf of

CMS Collaboration

28.04.2015

DIS 2015
Dallas, USA






Outline

1

- CMS detector at forward rapidities
- Soft diffraction - topologies
- Events with forward gap
- Events with central gap
- Cross section of the single and double dissociation
- Forward rapidity gap cross section
- Summary

CMS at forward rapidities


2



- Tracker $|\eta| < 2.4$, $p_T > 100$ MeV
- Electromagnetic calorimeter ECAL
- Hadronic Calorimeter HCAL
- Muon chambers

CMS at forward rapidities

3




- Tracker $|\eta| < 2.4$, $p_T > 100$ MeV
- Electromagnetic calorimeter ECAL
- Hadronic Calorimeter HCAL
- Muon chambers
- **H**adronic **F**orward calorimeters (HF) $3 < |\eta| < 5$
- Very Forward Calorimeter (CASTOR) $5.2 < |\eta| < 6.6$
- **Z**ero **D**egree **C**alorimeter (ZDC)
- **B**eam **S**cintillator **C**ounters BSC: $3.2 < |\eta| < 4.7$
- **F**orward **S**hower **C**ounters FSC: $6 < |\eta| < 8$

+ Totem (T1/T2 tracking detectors and RP roman pots) separate experiment


Soft Diffraction

4


non-diffraction




single diffractive dissociation (SD)



double diffractive dissociation (DD)




central diffraction



send to PRD,
arXiv:1503.08689
CMS-PAS-FSQ-12-005

Selection:

- $\sim 20 \mu\text{b}^{-1}$ of low pile-up data ($\mu=0.14$), from 2010
- Online: activity in either of the BSC - Minimum Bias trigger
- No vertex requirement (low diffractive masses $12 < M_X < 100 \text{ GeV}$ accepted)
- Diffractive offline selection: Large Rapidity Gaps within $|\eta| < 4.7$



MC Models


5

Monte Carlo:


- PYTHIA8-4C
 - diffraction generated according to Schuler&Sjostrand model from PYTHIA6
 - SD and DD cross sections scaled down by 10% and 12%
 - PYTHIA8-MBR (Minimum Bias Rockefeller model)
 - diffraction generated based on renormalized Regge theory model
 - developed for CDF
 - linear parametrization of the Pomeron trajectory $\alpha(t) = 1 + \epsilon + \alpha' t$
 - $\alpha' = 0.25 \text{ GeV}^{-2}$, $\epsilon = 0.08$ or 0.104
 - DD cross section scaled down by 15%
 - PYTHIA6-Z2*
 - PHOJET
 - QGSJET-II
 - EPOS
- } cosmic rays MC

Diffractive event topologies

6




Three experimental topologies based on the position of the LRG without CASTOR information






Diffractive event topologies

7




- η_{\max} (η_{\min}) highest (lowest) η of the particle candidate with $|\eta| < 4.7$

- $\Delta\eta = \eta^0_{\max} - \eta^0_{\min}$



Forward gap

8



$$\log_{10} M_Y < 0.5$$

$$0.5 < \log_{10} M_Y < 1.1$$

$$M_X, M_Y \text{ in GeV}$$

- Variable ξ defined as:

$$\xi_X = \frac{M_X^2}{s}$$

- Reconstructed:

$$\xi_X^{rec} = \frac{\sum (E^i - p_z^i)}{\sqrt{s}}$$

- And corrected for undetected particles:
(PYTHIA 8 MBR)

$$\log_{10} \xi_X^{cal} = \log_{10} \xi_X^{rec} + C(\xi_X^{rec})$$

Forward gap

9


Detector level distributions

Cross section measured in bins of ξ_x

full sample

no-CASTOR tag

CASTOR tag





- Data unfolded (acceptance and migration corrections)
- PYTHIA 8 MBR
- Corrections for pile-up



Forward gap

10

no-CASTOR tag




Castor tag



- Data with tag favor $\epsilon = 0.08$
- P8-4C and P6-2Z* higher than data in no-CASTOR tag
- P8-4C and P6-2Z* predicts raising behavior in no-CASTOR tag
- PHOJET, QGSJET, EPOS cannot describe data in tag sample

Central gap

11



DD

Range:

$$\log_{10} M_X > 1.1$$

$$\log_{10} M_Y > 1.1$$

$$\Delta\eta = \eta_{\max}^0 - \eta_{\min}^0$$





- Reconstructed:

$$\Delta\eta_{\text{rec}}$$

- And corrected for detector effects:

$$\Delta\eta_{\text{cal}} = \Delta\eta_{\text{rec}} - C$$

- Unfolding (response matrix from PYTHIA MBR)



- P8-MBR describes the data
- P8-4C underestimates
- P6-Z2* overestimates
- PHOJET, QGSJET, EPOS underestimates

Integrated cross section measured for 3 samples:

- FG2, no-CASTOR tag $-5.5 < \log_{10} \xi_X < -2.5$ $\log_{10} M_Y < 0.5$
- FG2, CASTOR tag $-5.5 < \log_{10} \xi_X < -2.5$ $0.5 < \log_{10} M_Y < 1.1$
- CG $\Delta\eta > 3$ $\log_{10} M_X > 1.1$ $\log_{10} M_Y > 1.1$

Cross section	$\sigma_{\text{no-CASTOR}}(\text{mb})$ SD dominated	$\sigma_{\text{CASTOR}}(\text{mb})$ DD dominated	$\sigma_{\text{CG}}(\text{mb})$ DD dominated
Data	$2.99 \pm 0.02^{+0.32}_{-0.29}$	$1.18 \pm 0.02 \pm 0.13$	$0.58 \pm 0.01^{+0.13}_{-0.11}$
PYTHIA 8 MBR	3.05	1.24	0.54
PYTHIA 8 4C	3.31	1.10	0.40
PYTHIA 6 Z2*	3.86	1.52	0.78
PHOJET	3.06	0.63	0.32
QGSJET-II 03	2.63	0.48	0.22
QGSJET-II 04	1.70	0.78	0.37
EPOS	2.99	0.85	0.31

From no-CASTOR tag sample, visible SD cross section:

- Subtraction of DD component \rightarrow from PYTHIA 8 MBR
- $\sigma^{SDvis} = 4.06 \pm 0.04 (stat)_{-0.63}^{+0.69} (syst) mb$ $-5.5 < \log_{10} \xi_X < -2.5$

From CASTOR tag sample, visible DD cross section:

- $\sigma_{CASTOR}^{DDvis} = 1.06 \pm 0.02 (stat) \pm 0.12 (syst) mb$ $-5.5 < \log_{10} \xi_X < -2.5$
 $0.5 < \log_{10} M_Y < 1.1$


From CG sample, visible DD cross section:

- $\sigma_{CG}^{DDvis} = 0.56 \pm 0.01 (stat)_{-0.13}^{+0.15} (syst) mb$ $\Delta\eta > 3$
 $\log_{10} M_X > 1.1$
 $\log_{10} M_Y > 1.1$

Extrapolation to the not observed region: PYTHIA 8 MBR ($\epsilon = 0.08$)

$$\sigma^{SD} = 8.84 \pm 0.08 (stat)_{-1.38}^{+1.49} (syst)_{-0.37}^{+1.17} (extr) mb \quad \xi_{X(Y)} < 0.05$$


$$\sigma^{DD} = 5.17 \pm 0.08 (stat)_{-0.57}^{+0.55} (syst)_{-0.51}^{+1.62} (extr) mb \quad \Delta\eta > 3$$



CMS results consistent with MBR predictions - SD cross section weakly rising with energy

Cross section

15




CMS results consistent with MBR and KP model predictions - DD cross section weakly rising with energy

Pseudorapidity gap cross section

16

- Forward rapidity gap - the largest of the two empty regions extending from the edge of the detector acceptance ($|\eta|=4.7$) to the nearest particle candidate


$$\Delta\eta^F = \max(\eta_{\min} - (-4.7), \eta_{\max} - (4.7))$$




- Unfolding done with PYTHIA 8 MBR with demand of at least 1 stable particle with $p_T > 200$ MeV and $|\eta| < 4.7$

Pseudorapidity gap cross section

17



- Exponential falling non-diffractive contribution
- Diffractive plateau at $\Delta\eta_F > 3$
- mixture of SD and DD events
- Best described: PYTHIA8+MBR with 0.08 intercept.



Differences in acceptance

- CMS performed measurement of diffractive dissociation cross section in pp
- Extrapolation of the SD and DD to the regions $\xi < 0.05$ and $\Delta\eta > 3$ gave:


$$\sigma^{SD} = 8.84 \pm 0.08 (stat)_{-1.38}^{+1.49} (syst)_{-0.37}^{+1.17} (extr) mb$$

$$\sigma^{DD} = 5.17 \pm 0.08 (stat)_{-0.57}^{+0.55} (syst)_{-0.51}^{+1.62} (extr) mb$$

- PHOJET, QGSJET, EPOS predict too few DD events
- PYTHIA8-4C, PYTHIA6-Z2*, PHOJET, EPOS do not predict ξ_x dependence for SD
- PYTHIA8-MBR describes the data in all the measured regions
- Data are consistent with SD and DD cross sections weakly rising with energy

Spares

Extrapolation



i	MC model	f^{SD}	$f_{\text{MBR}}^{\text{SD}}$
1	PYTHIA 8 MBR ($\epsilon = 0.08$)	2.18 (1.00)	2.18 (1.00)
2	PYTHIA 8 4C	2.32 (1.06)	2.51 (1.15)
3	PYTHIA 6 Z2*	2.29 (1.06)	2.89 (1.34)
4	PHOJET	2.06 (0.95)	2.18 (1.00)
5	QGSJET-II 03	2.72 (1.25)	3.19 (1.46)
6	QGSJET-II 04	3.62 (1.66)	2.30 (1.06)
7	EPOS	3.44 (1.58)	2.15 (0.99)
8	PYTHIA 8 MBR ($\alpha' = 0.125$)	2.27 (1.04)	2.34 (1.07)
9	PYTHIA 8 MBR ($\epsilon = 0.104$)	2.23 (1.03)	2.42 (1.11)
10	PYTHIA 8 MBR ($\epsilon = 0.07$)	2.16 (0.99)	2.09 (0.96)

i	MC model	f^{DD}	$f_{\text{MBR}}^{\text{DD}}$
1	PYTHIA 8 MBR ($\epsilon = 0.08$)	1.92 (1.00)	1.92 (1.00)
2	PYTHIA 8 4C	2.52 (1.32)	1.86 (0.97)
3	PYTHIA 6 Z2*	2.39 (1.25)	2.15 (1.13)
4	PHOJET	1.80 (0.94)	0.60 (0.31)
5	QGSJET-II 03	—	—
6	QGSJET-II 04	2.04 (1.07)	0.94 (0.49)
7	EPOS	4.73 (2.47)	1.93 (1.01)
8	PYTHIA 8 MBR ($\alpha' = 0.125$)	1.97 (1.03)	2.32 (1.21)
9	PYTHIA 8 MBR ($\epsilon = 0.104$)	2.00 (1.04)	2.37 (1.24)
10	PYTHIA 8 MBR ($\epsilon = 0.07$)	1.88 (0.98)	1.73 (0.90)