


# Measurement of longitudinal single-spin asymmetries for $W^\pm$ boson production in polarized p+p collision at $\sqrt{s}=510$ GeV at RHIC



◆ Introduction / Theoretical Aspects


◆ Experimental Aspects

◆ Recent Results / Current Analysis




**DIS 2015**  
XXIII International Workshop on  
Deep-Inelastic Scattering and  
Related Subjects  
Dallas, Texas  
April 27 – May 1, 2015



**Devika Gunarathne**  
**(for the STAR collaboration)**  
**Temple University**



# Evolving Picture of Proton's Spin Structure



**Valance Quarks**



**Sea Quarks**

# Anti Quark Polarization

**Spin sum rule for longitudinally Polarized proton :**

$$\langle S_p \rangle = \frac{1}{2} = \frac{1}{2} \Delta\Sigma + \Delta G + L$$

Jeffe and Monahar, 1990


$$\Delta\Sigma = \int (\Delta u + \Delta d + \Delta s + \Delta\bar{u} + \Delta\bar{d} + \Delta\bar{s}) dx$$

**~30 % from polarized inclusive DIS**

**Helicity PDF**


$$\Delta f(x, Q^2) \equiv f^+(x, Q^2) - f^-(x, Q^2)$$

## DSSV Global Analysis




PRD 80, 034030(2009)

# W Boson Production



- ❖ **Direct Coupling** to the Quark and anti Quark of interest.
- ❖ Maximum **violation of parity** leads to perfect **spin separation**.



- ❖ **High** resolution scale ( $Q^2$ ) set by the **W** mass.
- ❖ **Easy detection** via the **leptonic decay channels**.

Parity violating longitudinal  
single spin asymmetry

$$A_L = \frac{\sigma^+ - \sigma^-}{\sigma^+ + \sigma^-}$$


# W A<sub>L</sub> : Theoretical Aspects

W A<sub>L</sub>, highly sensitive to individual polarizations at forward and backward decay lepton pseudo rapidity (η<sub>e</sub>)

$$\eta = -\ln\left(\tan\left(\frac{\theta}{2}\right)\right) \quad \langle x_{1,2} \rangle \sim \frac{M_W}{\sqrt{s}} e^{\pm\eta_e/2}$$

$$\eta \lll 0 \rightarrow x_1 \ll x_2, \theta \rightarrow \pi$$


$$A_L^{e^-} \approx \frac{\int_{\otimes(x_1, x_2)} [\Delta\bar{u}(x_1)d(x_2)(1 - \cos\theta)^2 - \Delta d(x_1)\bar{u}(x_2)(1 + \cos\theta)^2]}{\int_{\otimes(x_1, x_2)} [\bar{u}(x_1)d(x_2)(1 - \cos\theta)^2 + d(x_1)\bar{u}(x_2)(1 + \cos\theta)^2]}$$




$$\frac{\Delta\bar{u}}{\bar{u}}$$

$$\eta \ggg 0 \rightarrow x_1 \gg x_2, \theta \rightarrow 0$$


$$A_L^{e^+} \approx \frac{\int_{\otimes(x_1, x_2)} [\Delta\bar{d}(x_1)u(x_2)(1 + \cos\theta)^2 - \Delta u(x_1)\bar{d}(x_2)(1 - \cos\theta)^2]}{\int_{\otimes(x_1, x_2)} [\bar{d}(x_1)u(x_2)(1 + \cos\theta)^2 + u(x_1)\bar{d}(x_2)(1 - \cos\theta)^2]}$$




$$\frac{\Delta\bar{d}}{\bar{d}}$$



# RHIC : Relativistic Heavy Ion Collider



# STAR Detector Overview



$$\eta = -\ln(\tan(\theta/2))$$


**TPC: Charge particle tracking**  
**BEMC, EEMC: EM Calorimetry**

**TPC** :  $-1.3 < \eta < +1.3$   
**BEMC** :  $-1.0 < \eta < +1.0$   
**EEMC** :  $+1.1 < \eta < +2.0$   
**FGT** :  $+1.0 < \eta < +2.0$

Year 2012


# Mid-rapidity ( $|\eta_{el}| < 1$ ) W Selection

## W → e + ν candidate



Isolated tracks → isolated EMC cluster  
Large opposite missing “energy”

## $P_T > 10$ GeV




Transverse plane views

$$\vec{p}_T^{balance} = \vec{p}_T^e + \sum_{\Delta R > 0.7} \vec{p}_T^{jets}$$


$$signed P_T - balance = \frac{(\vec{p}_T^e \cdot \vec{p}_T^{balance})}{|\vec{p}_T^e|}$$

## QCD background candidate




Several tracks → several EMC towers  
No large missing “energy”


## $E_T$ distribution



## Sign- $P_T$ balance



# Mid-rapidity Background Estimation



PRL 113,72301 (2014)

## Primary Background

**QCD processes** where fragments of a jet satisfy candidate  $e^{+/-}$  isolation cuts while **all other jets escape detection** outside the acceptance.

♣ **Second Endcap**  $-2 < \eta < -1.09$

♣ **Data driven QCD**  $|\eta| > 2$

## Electroweak BG


♣  $W \rightarrow \tau \nu$  MC Embedded in Zero-bias events

♣  $Z \rightarrow e^+ e^-$

**Forward rapidity ( $1 < \eta_e < 1.4$ ) W selection** uses similar technique as mid rapidity and **Background Estimation** improve using additional **Endcap Shower Maximum Detector (ESMD)**

# Results

## $W A_L(\eta_e)$ 2012+2011



PRL 113,72301 (2014)

- ❖ **Profile Likelihood** method used to extract **Asymmetries** from combination of **2012** and **2011** data.

- ❖  **$A_L(W^-)$  is larger** than the DSSV Predictions.


- ◆ The enhancement at  $\eta_e < 0$ , in particular is sensitive to the  $\Delta\bar{u}$ , polarized **antiquark distribution**.

- ❖  **$A_L(W^+)$  is consistent** with **theoretical predictions** using the DSSV polarized PDFs.

- ❖ The **Systematic uncertainties** for  $A_L$  are well **under control** for  $|\eta_e| < 1.4$ .


# Impact on Recent Global Analysis

- ❖ Preliminary Global analysis (**DSSV++**) from DSSV group and recent **NNPDF** includes preliminary **STAR 2012 W A<sub>L</sub>** data.
- ❖ **Shift** in central value for  $\Delta\bar{u}$  (negative -> positive) and  $\Delta\bar{d}$  due to A<sub>L</sub> W from STAR .
- ❖ **STAR 2012 W** results provide **significant constraints** on **anti u** and **anti d** quark polarization.



arXiv: 1304.0079

arXiv: 1304.0079




arXiv: 1403.0440

# STAR 2013 W Analysis Status

## 2013 Data Sample

### Barrel EMC triggered Integrated Luminosity



	L (pb <sup>-1</sup> )	P	FOM (P <sup>2</sup> L (pb <sup>-1</sup> ))
Run 9	12	0.38	1.7
Run 11	9.4	0.49	2.3
Run 12	72	0.56	24
<b>Run 13</b>	<b>~ 300</b>	<b>0.54</b>	<b>~ 87</b>




In 2013 STAR collected total integrated luminosity of  $\sim 300 \text{ pb}^{-1}$  at  $\sqrt{s} = 500 \text{ GeV}$  with an average beam polarization of  $\sim 54\%$ .

# Detector Calibration


## ♣ TPC

♣ **Charge Sign reconstruction** is based on bending of TPC tracks in the presence of magnetic field.


♣ Challenging environment of **charge-sign discrimination at high-pT using TPC in high luminosity / pile-up operation** - **Very careful TPC calibration required!!!!**



global TPC tracks




W candidate tracks




# Detector Calibration

## ❖ BEMC



❖ BEMC is a sampling calorimeter and it is segmented into 4800 towers, each with  $\Delta\eta \times \Delta\phi = 0.05 \times 0.05$



The geometry of the calorimeter towers in a BEMC module



Calibration need to be done considering all possible Energy ranges !!!

- ❖ In the low energy range **relative Calibration of towers** is done using abundant **MIPs (high P hadrons)**
- ❖ **E/P from electrons** is used to obtain **absolute calibration**.
- ❖ Other **High Energy probes** (from rare processes) such as **Z invariant mass peak position**, **slope of the sharp drop of W jacobian peak** will be used to calibrate in the high energy range.


# Forward-rapidity Analysis Status :

## FGT (Forward Gem Tracker)

## W Charge-sign Separation using FGT



FGT( $2 \geq \text{hits}$ )+Vertex+EEMC+Prompt




Prompt = actual measurement using TPC internal Multi-Wire proportional chamber

**~2.5 sigma** separation with FGT+VTX+EEMC  
+PROMPT (~1/3 events)

# STAR 2013 W A<sub>L</sub> Projections


## STAR W A<sub>L</sub> Projections



Higher precision results are expected from much larger statistics

Extension of forward and backward acceptance enhances sensitivity to  $\bar{u}$  and  $\bar{d}$  quark polarization

## Impact on antiquark polarization



arXiv: 1304.0079

# Summary / Outlook

- ◆ **The Production of W boson in polarized P+P collisions provides a new means to study the spin and flavor asymmetries of the proton sea quark distribution.**
- ◆ **STAR has measured the parity violating single spin asymmetry  $A_L$  for  $|\eta_e| < 1.4$  from 2012 and 2011 data, providing the first detailed look at the asymmetries  $\eta_e$  dependence.**
- ◆ **STAR 2012 W  $A_L$  results provide significant constraints on anti u and anti d quark polarization.**
- ◆ **Large statistics of STAR Run 2013 is being analyzed now in mid-rapidity region ( $|\eta| < 1$ ) and will use the 510 GeV Barrel EM calorimeter calibration which is in progress.**
- ◆ **Run 2013 analysis will be extended up to  $\eta = 1.4$  utilizing the EEMC in the forward region and then very forward region using the FGT as the tracking device requiring completion of careful calibration of EEMC and FGT.**
- ◆ **High precision results from 2013 will improve the constraints on the sea quark polarization.**


# Backup

# systematic uncertainties


- Beam polarization uncertainty: correlated scale 3.4%
- Relative luminosity uncertainty: correlated offset  $\Delta A_L = 0.007$
- Background estimation: less than 10% of statistical error

# unpolarized BG $\beta$


betaP\_2011




betaN\_2011



betaP\_2012




betaN\_2012



$$W^+ \beta: \sim 0.95, W^- \beta: \sim 0.9$$

where  $\beta = S/(S + B)$ ,  $S$  and  $B$  are the number of signal and background events in [25, 50] GeV

# STAR 2009 W Results



STAR pp500 Longitudinal		
Run	$L$ ( $pb^{-1}$ )	$W^+(W^-)$ raw yield
2009	12	462 (192)
2011	9	342 (103)
2012	77	2417 (734)

PRL 106, 062002 (2011)




# Recent results - Quark / Anti-quark pol. program

## Impact of new DSSV result

D. de Florian et al., Phys. Rev. Lett. 101 (2008)

RHIC Spin Collaboration (2012)

072001




○ From recent DSSV  
++ result incl.

STAR  $A_L$  data:  
 $\int_{0.05}^1 \Delta\bar{u}(x, Q^2) dx \approx 0.02$


$$\int_{0.05}^1 \Delta\bar{d}(x, Q^2) dx \approx -0.05$$

$M_W/\sqrt{s} = 0.16$

# Mid-Rapidity charge sign separation



- Charge sign reconstruction based on TPC track bending
- Estimate wrong sign contamination by fitting  $Q * E_T / p_T$  with Gaussian.



# Extracting Asymmetries using Profile Likelihood Method

- Profile Likelihood method used in extracting Asymmetries from combination of run 2012 and run 2011 data [simple gaussian uncertainties breakdown particularly for small 2011 data sample ]
- Define likelihood function for 8 spin-dependent yields from pair of symmetric  $\eta$  region of STAR

$$L = \prod_i^4 p(M_i^a | \mu_i^a) p(M_i^b | \mu_i^b) g(\beta^a) g(\beta^b)$$

$p(M_i | \mu_i)$  - Poisson probability, for measured spin sorted yield  $M_i$  in the expected value  $\mu_i$  given by:


$$\mu_{++}^a = I_{++} N(1 + P_1 \beta A_L^{+\eta_e} + P_2 \beta A_L^{-\eta_e} + P_1 P_2 \beta A_{LL})$$

$$\mu_{+-}^a = I_{+-} N(1 + P_1 \beta A_L^{+\eta_e} - P_2 \beta A_L^{-\eta_e} - P_1 P_2 \beta A_{LL})$$

$$\mu_{-+}^a = I_{-+} N(1 - P_1 \beta A_L^{+\eta_e} + P_2 \beta A_L^{-\eta_e} - P_1 P_2 \beta A_{LL})$$

$$\mu_{--}^a = I_{--} N(1 - P_1 \beta A_L^{+\eta_e} - P_2 \beta A_L^{-\eta_e} + P_1 P_2 \beta A_{LL})$$



$P_1, P_2$  - beam polarization  $A_L^{+\eta_e} (A_L^{-\eta_e})$  - single spin asymmetry  
 $A_{LL}$  - double spin asymmetry  $N$  - spin averaged yield  $I_{\pm\pm}$  - relative luminosity



$g(\beta)$  - Gaussian probability for estimated dilution background  $\beta$







Extract asymmetries from likelihood function  $L_{2011} \times L_{2012}$

# Run 12 Preliminary results compare to Final





PRL 113,72301 (2014)

# Endcap W Selection




# ESMD CUTS



$$R_{ESMD} = \frac{\sum_{i=-3}^{+3} E_i^U + E_i^V}{\sum_{i=-20}^{+20} E_i^U + E_i^V}$$


# Forward Rapidity Background Estimation and charge sign separation




PRL 113,72301 (2014)

# Run 12 ALL and Z AL results


$$A_{LL} = \frac{(\sigma^{++} + \sigma^{--}) - (\sigma^{+-} + \sigma^{-+})}{(\sigma^{++} + \sigma^{--}) + (\sigma^{+-} + \sigma^{-+})}$$



PRL 113,72301 (2014)



## Z invariant mass



PRL 113,72301 (2014)

- Probes different combination of quark polarizations

$$A_{LL} = \frac{\sigma^{++} - \sigma^{+-}}{\sigma^{++} + \sigma^{+-}} \quad A_{LL}^{W^+} \sim \frac{\Delta u}{u} \frac{\Delta \bar{d}}{\bar{d}} \quad A_{LL}^{W^-} \sim \frac{\Delta d}{d} \frac{\Delta \bar{u}}{\bar{u}}$$

- Asymmetries expected to be smaller, and first measurement consistent with predictions from DSSV

# W production: more details

Helicity structure can see in the differential cross section of W

$$\frac{d\sigma_{W^+}}{d\cos\theta} \propto \bar{d}(x_1)u(x_2)(1 + \cos\theta)^2 + u(x_1)\bar{d}(x_2)(1 - \cos\theta)^2$$

$$\frac{d\sigma_{W^-}}{d\cos\theta} \propto \bar{u}(x_1)d(x_2)(1 - \cos\theta)^2 + d(x_1)\bar{u}(x_2)(1 + \cos\theta)^2,$$

W tends to boost direction of the valance quark traveling

Helicity structure of the interaction causes lepton to emit parallel (antiparallel) to W-(W+)



higher (lower) x parton in the collision is most likely quark (antiquark) . And quark is very likely to come from valance region



W longitudinal momentum

e decay kinametics in lab frame related to W boost direction


$$p_{L,W} = \frac{\sqrt{s}}{2} (x_1 - x_2)$$

$$p_{L,e}^{lab} = \frac{1}{\gamma} p_{L,e}^* + \beta E_e^{lab}, \quad p_{L,e}^* = \cos\theta \cdot M_W/2 \quad (p_T^e = \sin\theta \cdot M_W/2),$$

# W, di-Jet and Z type events



**simulated W->e event**




**simulated di-jet event**

# Mid-rapidity ( $|\eta_e| < 1$ ) $W$ Selection


❖ Match  $P_T > 10$  GeV TPC tracks to EMC cluster

$$E_T^e / E_T^{4 \times 4} > 0.95$$




❖ Isolate from QCD di-jet type event

$$E_T^e / E_T^{\Delta R < 0.7} > 0.88$$






❖ Use Larger imbalance of transverse momentum

$$\vec{P}_T^{balance} = \vec{P}_T^e + \sum_{\Delta R > 0.7} \vec{P}_T^{jets}$$




❖  $e^+$  and  $e^-$  Charge sign Separation



# FGT

## Illustration of charge-sign discrimination




The TPC prompt hits are 'hits' using the ANODE wires located at large Z values at 209.5 cm for inner subsectors and 209.7 cm for outer subsectors. The electrons from a charge drift to the pad planes. The 'first signal in time' is then used to define a prompt hit.


# FGT track reconstruction

- Comparison of data / fast MC: Track reconstruction

$R$  ISOLUTION




$p_T$  [GeV/c]



Run13 Data (FGT+VTX  
+EEMC, no prompt)

PYTHIA QCD charged  
hadrons

$E$  [GeV]



PseudoRapidity

50GeV/c

PYTHIA W

PYTHIA QCD electrons

PYTHIA QCD photons

100GeV

0

3

The central values are both from the theoretical predictions. So the central values of the polarized anti-u and anti-d quarks should be same. The the uncertainties are estimated from the W yields. For the old version, it maybe estimated from the run9 W efficiency. But for the new version, the W yields are estimated from run13 W efficiency. Due to the higher  $\langle z_{dc} \rangle$  rate , the later one should be a little smaller. I roughly compared the error bars in these two projection plots, the differences is very slight, something like  $<10\%$ . But, for the new version, we indeed don't have the corresponding polarized anti-u and anti-d distribution.

**[simple gaussian uncertainties breakdown particularly for small 2011 data sample ]**

