DIS 2015 - XXIII. International Workshop on Deep-Inelastic Scattering and Related Subjects



Contribution ID: 230

Type: not specified

A dedicated eRHIC Detector and Interaction Region design

Tuesday 28 April 2015 14:50 (25 minutes)

Construction of an Electron-Ion Collider with luminosities exceeding $10^{33} cm^{-1}s^{-1}$ is becoming a project of highest priority for the US Nuclear Physics community. The main physics topics to be explored at this new facility are (i) the polarized sea quark and gluon distributions in the nucleon, (ii) QCD dynamics of the low-x, high density gluon regime, (iii) hadronization and energy loss in the nuclear medium [1]. One of the considered construction options is the addition of a high-energy polarized electron beam to the existing RHIC hadron machine, converting it into an Electron-Ion Collider (eRHIC) [2]. A dedicated eRHIC detector, designed to efficiently register and identify deep inelastic electron scattering (DIS) processes in a wide range of center-of-mass energies available with the new collider is one of the key elements of such an upgrade. The progress on the detector and interaction region design work will be shown, and the simulation results presented.

 A. Accardi et al., "Electron Ion Collider: The Next QCD Frontier - Understanding the glue that binds us all" (EIC White Paper), arXiv:1212.1701v3 (2014).

[2] E.C. Aschenauer et al., "eRHIC design study (An Electron-Ion Collider at BNL)", arXiv:1409.1633 (2014).

Author: Dr KISELEV, Alexander (Brookhaven National Laboratory)

Presenter: Dr KISELEV, Alexander (Brookhaven National Laboratory)

Session Classification: WG7 Future experiments

Track Classification: WG7 Future experiments